

SPECIAL VARIETIES AND CLASSIFICATION THEORY

FRÉDÉRIC CAMPANA

Abstract: We describe two structure theorems in the birational geometry of complex projective manifolds, analogues of the structure theorems for Lie algebras, reducing these first to semi-simple and solvable ones, the solvable being iterated extensions of abelian ones. The roles of semi-simple, solvable and abelian are respectively played here by the orbifolds of general type, special manifolds, and third: manifolds either rationally connected or with $\kappa = 0$.

A complex projective manifold X , is said to be **special** if it has no map onto an orbifold of general type, the orbifold structure on the base being given by the divisor of multiple fibres (suitably defined). We show that manifolds either rationally connected or with $\kappa = 0$ are special. We also show that X is special if and only if it has no Bogomolov sheaf, which are rank one coherent subsheaves of maximal Kodaira dimension p of Ω_X^p , for some $p > 0$.

An orbifold version of Iitaka's $C_{n,m}$ additivity conjecture implies that special manifolds have a canonical decomposition as towers of fibrations with generic (orbifold) fibres either rationally connected or with zero Kodaira dimension. This decomposition is obtained by alternatively iterating orbifold Iitaka fibrations and suitably defined rational quotients.

For any complex projective manifold X , we next construct a functorial fibration $c_X : X \rightarrow C(X)$ called its **core**¹, such that its general fibre is special, and contains any special subvariety of X meeting it. The core is thus the constant (resp. identity) map iff X is special (resp. of general type).

As a consequence of the special case of the orbifold version of Iitaka conjecture $C_{n,m}$ when the base orbifold is of general type, we show that the core has an orbifold base of general type if X is not special.

The core thus canonically decomposes any X into its “components” in the dichotomy special (the fibres) versus general type (in the orbifold sense: the base). Special manifolds being (conjecturally) orbifold combinations of the first two types of the fundamental trichotomy (rationally connected, $\kappa = 0$, general type) of classification theory.

This decomposition is deeply linked with other aspects of classification theory: we indeed conjecture, and show in some few cases, that special manifolds have an almost abelian fundamental group, and are exactly the ones having zero Kobayashi metric, or a potentially dense set of rational points over any field of definition finitely generated over $\mathbb{Q}$. This immediately leads to a natural extension of Lang's conjectures to arbitrary X 's (and even to orbifolds).

This decomposition gives a simple synthetic view of the structure of arbitrary X 's, and indicates that the natural frame of classification theory is the category of orbifolds, to which our observations should be extended.

The present observations actually work for any compact Kähler X , which are considered here. Most of them can also be applied to other algebraically closed fields (with the usual problems of separability in positive characteristics).

¹ From the french “coeur”. A case to which A. Dumas's statement applies well: “Comme l'anglais n'est que du français mal prononcé...” (in “Vingt ans après”, Chap LXVII: “Londres”)

CONTENTS

Introduction	4
1. Orbifold Base of a Fibration.	8
1.1. Fibrations	8
1.2. Orbifolds.	10
1.3. The Kodaira Dimension of a Fibration	11

1.4. The Sheaf of Differential Forms determined by a fibration	16
1.5. Semi-continuity of the Kodaira Dimension	19
1.6. Composition of Fibrations	19
2. Special Fibrations and General Type Fibrations	21
2.1. Special or General Type Fibrations	21
2.2. Special Fibrations Dominate General Type Fibrations	22
2.3. General Type Fibrations and Bogomolov Sheaves	26
2.4. General Type Reduction	28
2.5. The Algebraic Reduction	30
2.6. The Category of Orbifolds	31
3. The Core	32
3.1. The Core: Functoriality Properties	32
3.2. Construction of the Core as the Lowest Special Fibration	34
3.3. Rationally connected manifolds	36
3.4. The Rational Quotient and The Core	37
3.5. Surfaces	38
3.6. Higher Kodaira Dimensions	40
3.7. Threefolds	41
3.8. Orbifold Surfaces	43
4. Orbifold Additivity	45
4.1. Orbifold Conjecture $C_{n,m}$	45
4.2. Reduction to Weak Positivity	47
4.3. Orbifold Weak-Positivity	48
4.4. Twisted Weak Positivity	51
5. Geometric Consequences of Additivity	52
5.1. Varieties with $\kappa=0$	52
5.2. The Albanese Map	53
5.3. Varieties of General Type	53
5.4. The Decomposition Theorem	55
5.5. Finite étale Covers	56
5.6. Essential and Bogomolov Dimensions	56
5.7. Fibre Products	57
5.8. Construction of the Core as the Highest General Type Fibration	57
6. Geometric Conjectures	57
6.1. Orbifold Rational Connectedness	58
6.2. Orbifold Iitaka-Moishezon Fibration	59
6.3. Orbifold Rational quotient and Iitaka Fibration of a Special fibration	60
6.4. The Core as an Iterated Orbifold Rational Quotients and Iitaka Fibrations	61
6.5. Deformation invariance	62
6.6. Weakly-Special Manifolds	63
6.7. The Weak Version of the Core	64
6.8. Classical gcd -Multiplicities	65
6.9. Irreducible Special Manifolds	66

6.10. Manifolds with nef anticanonical bundle	66
7. The Fundamental group	67
7.1. The Abelianity conjecture	67
7.2. Linear and Solvable Quotients	68
7.3. Manifolds Covered by $\mathbb{C}^n$	69
8. An Orbifold generalisation of Kobayashi-Ochiai's Extension theorem	70
8.1. Statements	70
8.2. Sketch of Proof of the Kobayashi-Ochiai's Extension Theorem.	72
8.3. Pseudo-Volume and Ricci Forms	73
8.4. Pseudo-Volume forms on Varieties of General Type	74
8.5. The Lemma of Ahlfors-Schwarz	75
8.6. Proof of the Orbifold Version	76
8.7. Extension of Pluri-Canonical Meromorphic Forms	78
8.8. Proof that H' -Specialness Implies Log-Rational Connectedness	79
9. The Core and the Kobayashi Metric	80
9.1. The Kobayashi Pseudo-Metric	80
9.2. Hyperbolically Special Manifolds	81
9.3. Similarities	82
9.4. The Origin of Conjecture III _H	82
9.5. Surfaces	83
9.6. Threefolds	83
9.7. The Fundamental group	84
9.8. The Albanese map	84
9.9. Geometric variants of H -speciality	85
9.10. The Core and the Splitting of the Kobayashi Metric	86
10. Arithmetic Aspects	87
10.1. Fields of Definition	87
10.2. Arithmetically Special Manifolds	88
10.3. A Related Conjecture of Abramovitch and Colliot-Thelene	90
11. The Function Field Case	90
11.1. The (Complex) Function Field Case	90
11.2. Split and "Dominated" Fibrations: the Rational Quotient	92
12. Appendix: Geometric quotients	94
12.1. Covering Families	95
12.2. Construction of the Quotient for an Irreducible Covering Family	96
12.3. Reduction to the case where T is irreducible	100
12.4. Zariski Regularity	101
12.5. Stability	104
Références	104

Introduction

The structure of arbitrary complex Lie algebras is described to a large extent by the following two theorems: the first one says that any such Lie algebra is in a unique way an extension of a semi-simple Lie algebra by a solvable one. Next, this extension can further be decomposed, also in a canonical functorial way, as a composition of extensions with abelian kernels.

The objective of this paper is to establish two similar structure results for arbitrary complex projective varieties X , in birational geometry.

The first theorem asserts the existence and uniqueness of a fibration $c_X : X \dashrightarrow C(X)$, called the core of X , such that its general fibres are special, while its orbifold base is of general type. The second result (only conjectural) asserts that the core can be obtained in a functorial canonical way as a composition of orbifold rational quotients and Iitaka fibrations. This statement depends on an orbifold version of Iitaka's conjecture $C_{n,m}$.

One thus gets a canonical decomposition of X , the elementary components of which are the varieties of pure type in the fundamental trichotomy of algebraic geometry (rationally connected, $\kappa = 0$, general type). The orbifold structures seem to be unavoidable, and to lie at the heart of classification theory, which should thus deal with this category of objects.

We now describe the content of the different sections.

In §1, we introduce the notion of orbifold base of a holomorphic fibration $f : X \rightarrow Y$ (f is thus surjective, and its fibres are connected) between compact Kähler manifolds. This is simply (1.1.4) a $\mathbb{Q}$ -divisor on Y , suitably defined by the multiple fibres of f . This orbifold structure carries naturally a canonical bundle, a Kodaira dimension and a fundamental group.

The Kodaira dimension $\kappa(Y, f)$ is the minimum of the Kodaira dimensions of the orbifold bases, when f' runs over all models of f , which are the fibrations bimeromorphically equivalent to f . This minimum is achieved by f itself when $\kappa(Y) \geq 0$ (by 1.12), but may be smaller in general.

We next show (1.7) that $\kappa(Y, f)$ behaves as in the “classical” case (where $f = id_X$) under composition with generically finite maps $u : X' \dashrightarrow X$ and Stein factorisation.

In the next section, we define canonically a rank one subsheaf F_f of Ω_X^p by saturating the inverse image of K_Y . This sheaf is an intrinsic invariant of the equivalence class of fibration. Its Kodaira dimension is $\kappa(Y, f)$. As an application, the orbifold base of f has Kodaira dimension $\kappa(Y, f)$ if f is what we call neat (see 1.2).

We next show (1.25) the countable upper semi-continuity of $\kappa(Y, f)$ in families.

Finally, we show how to compute, on suitable models, the orbifold structure on the base of a composition $g \circ f$ of two fibrations, from the orbifold bases of g and f . This computation plays an essential role in the proof of the shown case of the orbifold additivity conjecture in §4. It is also the clue to the definition of the orbifold base of a fibration between orbifolds. See section 2.6.

In §2, we first define general type fibrations $f : X \dashrightarrow Y$ as the ones having $\kappa(Y, f) = \dim(Y) > 0$, special manifolds X are defined as the ones having no f of general type, and special fibrations as

the ones having special general fibres. We list without proof some examples of special manifolds. The two most important ones are the rationally connected manifolds, and manifolds with $\kappa = 0$. Notice however that being special does not give any restriction on the Kodaira dimension, except for the top one. For example, any elliptic surface with base elliptic or rational is special if it has a section. Somewhat unexpectedly maybe, the moduli of fibrations do not play any role in our considerations.

We next show various geometric properties of special and general type fibrations. The most important one (2.12) is that any special fibration $f : X \dashrightarrow Y$ dominates any general type fibration $g : X \dashrightarrow Z$, in the sense that there exists $\phi : Y \dashrightarrow Z$ such that $g = \phi \circ f$.

From which one concludes that on any X there exists at most one fibration both special and of general type. One of the main results of the present paper is the **existence** of such a fibration. We get it by the two possible approaches: either from “above”, as the “lowest special” fibration on X , obtained by geometric means (see section 3), or from “below”, as the “highest general type fibration” on X (see 5.17). The second approach is less geometric and hides the motivations and origins of the constructions made, but has technical advantages similar to those of the axiomatic method, turning theorems into definitions, once the foundations have been laid. (See footnote 2 below) ¹

The next subsection shows the bijective correspondance between Bogomolov sheaves and general type fibrations (2.25). From which we conclude that special manifolds are characterised by the absence of such sheaves. Applying a result from [C95], we obtain a first simple proof that X is special if either rationally connected or with $c_1(X) = 0$.

We next show (2.37), among other things, that the general fibre X_s of a fibration $h : X \rightarrow S$ is special if it is special for any s in a set E not contained in a countable union of Zariski closed subsets with empty interior of Z . This property, which we call Zariski regularity (for specialness) is, together with 2.12, one of the main ingredients in the proof that the core of a manifold is a special fibration (3.3).

This section ends with a brief sketch of the extension of the considerations of the present paper to the orbifold category. It seems that most of them extend without any difficulty to the orbifold context, including the notions of orbifold modification and of differential form on an orbifold (Y/D) with smooth Y and the support of D of normal crossings. In particular the notion of Bogomolov sheaf and the sheaf F_f associated to a fibration between orbifolds make sense. The geometric aspect seems more delicate to handle (see section 6.1 for some subtle problems on surface orbifolds which are classical in the non-orbifold context).

The notion of orbifold differential form we introduce interpolates between the usual one when D is empty, and the classical notion of $\Omega_Y(\log(D))$ when D is reduced, which corresponds to the limiting case of infinite multiplicities. It seems that this topic deserves by itself further developments.

In §3, we construct and start to study the core $c_X : X \dashrightarrow C(X)$ of a manifold. We first show (3.3) that its general fibres are special (this fails for general singular varieties, and does not follow from the original definition, obtained applying the construction of meromorphic quotients recalled in

¹B.Russell once compared these advantages to the ones of stealing over honest work

section 12). From this result, we immediately get (3.21) that rationally connected manifolds are special (simply because $\mathbb{P}^1$ is special). The notions around rational connectedness are recalled. We next deduce in (3.25) from [G-H-S] that if $R(X)$ is the rational quotient of X (see 3.22 for this notion), then $C(R(X)) = C(X)$.

We next describe (3.30 and 3.37) the core and list the special manifolds in dimensions 2 and 3, after having introduced (see §3.6) the “higher Kodaira dimensions” of a (compact complex) manifold. From the description so obtained, we deduce that in these cases the core is a fibration of general type (when X is not special). The “decomposition theorem” (5.8) asserts that this is true in any dimension.

The next §4 states (see 4.3) the orbifold version of Iitaka conjecture $C_{n,m}$, and shows (4.4) the special case when its orbifold base is of general type. This result is one of the main technical tools of the present paper. The proof consists in extending the classical weak positivity results of Fujita, Kawamata and Viehweg for direct image sheaves of pluricanonical forms to the orbifold situation, by suitably introducing the orbifold divisor into the proofs.

§5 contains the geometric consequences of the orbifold additivity result (theorem 4.4): Manifolds with $\kappa = 0$ are special (5.1), the Albanese map of a special manifold is surjective, connected, and has no multiple fibres in codimension one (5.3). We also show (5.5 (resp. 5.7)), that the essential dimension $\text{ess}(X) := \dim(C(X))$ of X is equal to its dimension $\dim(X) = n$ (resp. to $(n-1)$) iff X is of general type (resp. iff X has a fibration of general type with generic fibre a curve either rational or elliptic).

These are two special cases of the “decomposition theorem” already stated above. This theorem is established in (5.8). We also show that the decomposition theorem implies the invariance of the essential dimension under finite étale covers, in particular that such covers of a special manifold are still special. We finally observe (see subsection 5.6) that the decomposition theorem asserts that any nonspecial X has a (unique) fibration both special and of general type, and a unique maximum Bogomolov sheaf.

We then show that the Stein factorised product of two fibrations of general type is itself of general type ² This gives, together with another application of 4.4, the second construction of the core “from below”, as the “highest general type” fibration on a given X .

The §6 presents various conjectures related to the core. We first start with defining and comparing some orbifold variants of the notion of rational connectedness. One of them (the weakest, possibly equivalent to the strongest of these notions) turns out to give a good substitute of the notion of rational quotient in the orbifold context, provided the orbifold version of conjecture $C_{n,m}$ holds. This implies (see section 6.4) that the core can be canonically and functorially decomposed into a

²Maybe surprisingly, it was not noticed, since [Bo78] that one could so naturally define a fibre product of two Bogomolov sheaves, although the techniques used in the present paper are available since two decades. This is because the question is obvious if one starts from “above”: the geometry of orbifolds and general type fibrations, but very hidden and unnatural if one starts from the sheaves.

succession of orbifold Iitaka-Moishezon³ fibrations and rational quotients (defined in section 6.3). See §6 for more details. This construction thus “only” depends on $C_{n,m}$ in its orbifold version. So that the core actually gives directly the final step of this decomposition, without producing the intermediate steps.

In subsection 6.5, we conjecture the deformation invariance of all the invariants introduced (in particular: $ess(X)$ and the higher Kodaira dimensions. So that in particular, deformations of special manifolds should be special). The next subsections deal with two other similar theories built on related notions of specialness and general type fibrations? They seem to both differ from the one presented here, but presently no example is known that distinguishes them.

In §7, we consider the fundamental group, and conjecture (“abelianity conjecture” 7.1) that a special manifold should have an almost abelian fundamental group. This conjecture is supported and motivated by the fact that rationally connected manifolds are simply connected, that this conjecture is standard for the case $\kappa = 0$, and the preceding conjectural orbifold decomposition of any special manifold as a tower of fibrations with fibres of these two types. This conjecture seems to sum up all conjectures with the same conclusion.

We show that this conjecture is true for linear and torsionfree solvable representations of the fundamental group, as an immediate consequence of previous result by various authors. As usual, we extend this conjecture to the orbifold case, anyway necessary to solve the non-orbifold one in higher dimensions.

In §8, we establish and apply an orbifold version (see (8.2)) of the famous extension theorem of Kobayashi-Ochiai, asserting roughly that meromorphic nondegenerate (ie: somewhere submersive) maps ψ from a dense Zariski open subset U of a complex manifold V to a variety of general type Y do not have essential singularities along the boundary divisor $D := (V - U)$. Our version essentially says this holds true if ψ factorises as $\psi = f \circ \phi$, with $f : X \dashrightarrow Y$ is a fibration of general type, and $\phi : U \dashrightarrow X$ meromorphic.

The proof is an orbifold version of the proof of Kobayashi-Ochiai. This result implies among other more general results that a manifold X is special if there is a meromorphic nondegenerate map $\phi : \mathbb{C}^n \dashrightarrow X$. This is a special instance where the Kobayashi pseudometric of X vanishes identically if ϕ has dense image in the metric topology.

The next §9 discusses two conjectures concerning the Kobayashi pseudometric of a compact Kähler X . The first one (9.2) states that special manifolds are exactly the ones having zero Kobayashi pseudometric d_X . The second one (9.10) states that for any X , d_X is the lift by the core $c_X : X \dashrightarrow C(X)$ of the orbifold Kobayashi pseudometric of the base orbifold of c_X , defined in 9.23, and that this orbifold pseudometric is a metric outside some algebraic subset S of $C(X)$. Recall that the orbifold base of c_X is of general type (if not a point), so this second part of the conjecture 9.10 is the orbifold version of Lang’s hyperbolicity conjecture.

The next §10 (resp. §11) is the exact analog of §9 for arithmetic geometry (resp. the function field version of §10)). We do not give a more detailed description, here, because our discussion rests only on known results by many authors.

³Usually called Iitaka fibration. The construction obtained independently by B.Moishezon [Mois] remained apparently unnoticed

This paper ends with the §12, which is a technical appendix exposing (with proofs) the construction of meromorphic quotients of equivalence relations of geometric origin (12.4) identifying points connected by chains with irreducible components in a given family of subvarieties.

This construction appeared it seems for the first time in [C81]. We follow here the original argument, in a slightly simplified form. (This construction was exposed also in [Ko 96] by a different approach, corrected and exposed in a detailed way in [De]).

We complete this construction by introducing the notions of Zariski regularity (section 12.4) and stability (section 12.5), of constant use in this context. In particular, the notion of stability permits to give a criterion (12.31) for the quotient to have fibres in the class as subvarieties used to construct the chains providing the equivalence relation.

The present version is the sixth one. It differs from the original one in two aspects: first by the definition of multiple fibres (based on $\inf$, no longer $\gcd$. See 1.1.4), second, and more importantly, by the proof that the base orbifold of the core is of general type, which was only conjectured in the preceding versions. The first point was also observed, independently of me, by S.Lu, after he read the first versions of the present paper: one can replace the $\gcd$ -multiplicities by the present ones without changing any statement or proof (except for a small refinement in the proof of the former version of proposition 4.20). The new definition is better behaved with respect to hyperbolicity questions and Bogomolov sheaves (but seemingly not with respect to the fundamental group, for which $\gcd$ -multiplicities work very well).

Aknowlegements: I would like to thank F. Bogomolov, J.P. Demailly, and C. Voisin for discussions which permitted me to improve sections 7 and 10. P.Eyssidieux asked me about the function field version of §10, which lead to §11. F.Bogomolov noticed that general type fibrations might be linked with what is called here Bogomolov sheaves; this link is established in 2.25 (One direction was in fact shown in the first version in the transcendental context of section 8). Moreover, after a first version of this work was written, I got further stimulating comments and references included here from D.Abramovitch, J.L. Colliot-Thélène and B.Totaro (who independently suggested me to include Conjecture IV_A).

I thank heartily thank L.Bonavero, S.Druel, J.P.Demailly, L.Manivel, C.Mourougane, M.Paun and E.Peyre for their carefull reading of §8,9,12,14,12,12 and 7 of the previous version, (which are now §2,4,8,12,8,8 and 10), which they exposed in Grenoble in July of 2002. And also for pointing many mistakes, gaps and inaccuracies. They also suggested me to reorganise the exposition according to the technical order of proofs. The present version thus differs considerably from the preceding ones in this respect. Further, L.Bonavero and S.Druel provided me with considerable help in the manipulation of Latex.

Special thanks are due to E.Viehweg, who gave me a decisive hint in the proof of the crucial Lemma 4.22.

1. Orbifold Base of a Fibration.

1.1. **Fibrations.** Before giving definitions, let us start with

1.1.1. **A motivating example.**

Example 1.1. Let $X_0 := E \times \mathbb{P}_1(\mathbb{C})$, where E is an elliptic curve. Let $X := \tilde{X}/j$, where $\tilde{X} := E \times C$, where C is a hyperelliptic curve and $j = t \times h$ is the involution acting diagonally on $\tilde{X}$ by a translation t of order 2 on E , and by the hyperelliptic involution h on C . Then both X_0 and X have natural fibrations on $\mathbb{P}_1(\mathbb{C})$ with generic fibre E .

They cannot be distinguished by this information, although they differ fundamentally at the levels of Kodaira dimension, fundamental group, Kobayashi pseudo metric and arithmetics (for appropriate choice of E). If, however, we take into account the multiple fibres of the fibrations onto $\mathbb{P}_1(\mathbb{C})$, we see that in the second case (of X), the base is not really $\mathbb{P}_1(\mathbb{C})$, but rather the orbifold of general type C/h .

This we shall now generalise.

1.1.2. Fibrations. A **fibration** $f : X \dashrightarrow Y$ is a surjective (*i.e.* dominant) meromorphic map with connected fibres between normal irreducible compact complex analytic spaces X and Y (see below for the precise definition of the fibres of a meromorphic map).

Of course, this fibration is said to be holomorphic (or regular) if so is the map f .

Another fibration $f' : X' \dashrightarrow Y'$ is said to be **equivalent** to f if there exists bimeromorphic maps $u : X \dashrightarrow X'$ and $v : Y \dashrightarrow Y'$ such that $f' \circ u = v \circ f$. We denote by F or X_y the generic fibre of f . We also say that f' is a **model** or a **representative** of f (we shall not distinguish between f and its equivalence class).

A fibration $f : X \dashrightarrow Y$ canonically defines (see [C85]) a meromorphic map $\phi_f : Y \dashrightarrow \mathcal{C}(X)$, with $\mathcal{C}(X)$ the Chow-Scheme of X , by sending the generic $y \in Y$ to the point of $\mathcal{C}(X)$ parametrising the reduced fibre X_y of f over y . Let $\Phi_f \subset \mathcal{C}(X)$ be the image of Y by ϕ_f : it is a compact irreducible analytic subset of $\mathcal{C}(X)$ bimeromorphic to Y such that its incidence graph is bimeromorphic to Y . The above correspondance induces a bijective map between equivalence classes of fibrations and compact irreducible analytic subsets of $\mathcal{C}(X)$ with incidence graph bimeromorphic to X .

Definition 1.2. Assume that $f : X \rightarrow Y$ is a holomorphic fibration. An irreducible Weil divisor D on X is said to be **f -exceptional** if $f(D)$ has codimension 2 or more in Y . We say that $f : X \rightarrow Y$ is **neat** if f is holomorphic, X, Y are smooth, and if there moreover exists a bimeromorphic holomorphic map $u : X \rightarrow X'$ with X' smooth such that each f -exceptional divisor of X is also u -exceptional.

By Raynaud and Hironaka flattening theorems, any fibration $f_0 : X_0 \dashrightarrow Y_0$ has a neat model, in which X' may be chosen arbitrarily (bimeromorphic to the domain X_0 of the initial fibration f_0). Indeed: first blow-up X' in such a way that $f_0 \circ b : X' \dashrightarrow Y_0$ is holomorphic, where $b : X' \dashrightarrow X_0$ is bimeromorphic; then flatten $(f_0 \circ b)$ by blowing-up Y_0 to get a smooth Y . Finally, take a smooth model X of $X' \times_{Y_0} Y$.

Assume the fibration f is holomorphic. We say that the holomorphic fibration $f' : X' \rightarrow Y'$ with X', Y' smooth **dominates** f if there exists a commutative diagram in which u, v are bimeromorphic; obviously, f' is then equivalent to f :

$$\begin{array}{ccc}
X' & \xrightarrow{u} & X \\
f' \downarrow & & \downarrow f \\
Y' & \xrightarrow{v} & Y
\end{array}$$

Observe that any f' dominating a neat fibration is itself neat.

More generally, we say that the fibration $f' : X' \dashrightarrow Y'$ dominates the fibration $f : X \dashrightarrow Y$ if there exists a diagram as above with u, v meromorphic and bimeromorphic. This notion is well-defined on equivalence classes.

Let $f : X \rightarrow Y$ be a holomorphic fibration with X, Y smooth. We say (as in [Vi82]) that f is **prepared** if the locus $Y^* \subset Y$ of points y with a smooth f -fibre X_y has a complement contained in a normal crossing divisor $D \subset Y$ such that, moreover, $f^{-1}(D)$ is also a divisor of normal crossings in X .

Any fibration is dominated by a prepared model, again by an immediate application of Hironaka's results. The models of fibrations considered can thus always be assumed to be prepared.

1.1.3. Multiplicities. Assume now X, Y to be smooth and f to be a holomorphic fibration. Let $|\Delta| \subset Y$ be the union of all codimension one irreducible components of the locus of y 's $\in Y$ such that the scheme-theoretic fibre of f over y is not smooth. For each component Δ_i of $|\Delta|$, let $D_i := \sum_{j \in J} m_{i,j} D_{i,j}$ be the union of all components (counted with their scheme-theoretic multiplicities) of $f^*(\Delta_i)$ which are mapped surjectively onto Δ_i by f . Define $m_i = m(f, \Delta_i) =: \inf(m_{i,j}, j \in J)$: it is called the **multiplicity of f along Δ_i** .

1.1.4. Multiplicity Divisor of a Fibration. The $\mathbb{Q}$ -divisor $\Delta(f) = \sum_{i \in I} (1 - 1/m_i) \Delta_i$ on Y will be called the **multiplicity divisor of f** . The motivation for its introduction comes from the examples above and in Example 1.3 below.

Observe we can also write: $\Delta(f) = \sum_{D \subset Y} (1 - 1/m(f, D)) \cdot D$, where D ranges over the set of all irreducible divisors of Y , and $m(f, D)$ is the **multiplicity of f along D** , defined as $m(f, D) := \inf(m_j, j \in J(f, D))$, where $f^*(D) := \sum_{j \in J(f, D)} m_j D_j + R$, and $J(f, D)$ is the set of all components of $f^*(D)$ which are mapped surjectively onto D by f , while R is f -exceptional. Indeed, $m(f, D) = 1$ if D is not a component of $|\Delta|$.

1.2. Orbifolds.

1.2.1. Notion of Orbifold. An **orbifold** (Y/Δ) is a pair consisting of a compact irreducible complex space Y together with a Weil $\mathbb{Q}$ -divisor of the form: $\Delta =: \sum_{i \in I} (1 - 1/m_i) \Delta_i$ for distinct prime divisors Δ_i of Y , and positive integers m_i . We also say that Δ is an orbifold structure on Y . We write $|\Delta|$ for the support of Δ , in which the coefficient of each Δ_i is one. Or equivalently: each $m_i = +\infty$.

If Y is smooth, and if the support of Δ is a simple normal crossings divisor, such pairs (Y/Δ) were introduced and used by V.Shokurov in [Sk92] under the name of “standard pairs”. We shall say that the orbifold (Y/Δ) is a **klt-orbifold** if the pair (Y, Δ) is klt (this is an abbreviation for “Kawamata-Log-terminal”). This seems to be the right category to consider, morphisms being the obvious ones.

The next example shows why such pairs are rather called orbifolds here. We also note that this term is commonly used in similar situations either in differential geometry, or when Y is a curve (in [F-M] or [L], for example).

Example 1.3. *Let $f : X \rightarrow Y$ be a finite (ramified) Galois cover between manifolds. Let its ramification divisor be: $\Delta =: \sum_{i \in I} (1 - 1/m_i) \Delta_i$, the order of ramification above the generic point of Δ_i being m_i .*

This preceding example occurs in the construction of fibrations with multiple fibres, as in 1.1 for example.

1.2.2. Orbifold Base of a Fibration.

Definition 1.4. *Let f be a holomorphic fibration as in 1.1.2, and $\Delta =: \Delta(f)$ be the multiplicity divisor of f . We call (Y/Δ) the **orbifold base** of f .*

1.2.3. Orbifold invariants and fibrations.

Definition 1.5. *Let (Y/Δ) be an orbifold. We define its **canonical bundle** as the $\mathbb{Q}$ -divisor $K_{Y/\Delta} =: K_Y + \Delta$ on Y and its **Kodaira dimension** as $\kappa(Y/\Delta) =: \kappa(Y, K_Y + \Delta)$.*

One can also associate to an orbifold its fundamental group, considered below.

The term “orbifold” arises in this context from the following reason: first from the special case when f is the quotient of a fibration $f' : X' \rightarrow Y'$ by a finite group G acting on some Galois cover X' of X , the action of G preserving the fibration f' , and such that f' has no multiple fibre in codimension 1. In this case, $(Y/\Delta(f))$ is precisely the orbifold (Y'/G) . We then say that the orbifold $Y/\Delta(f)$ has the **unfolding** (Y', G) . When Y is a curve, such an unfolding exists, except when $Y = \mathbb{P}^1$ and Δ consists of one or two punctures with different multiplicities. In the general case, such unfoldings exist locally on Y (but not globally in general. M.Kapovitch explained me a very beautiful construction in dimension two for orbifolds of general type).

1.3. The Kodaira Dimension of a Fibration.

1.3.1. Kodaira Dimension. Define now in general, for $f : X \dashrightarrow Y$ a fibration between irreducible compact complex spaces X and Y :

$$\kappa(Y, f) := \inf \{ \kappa(\bar{Y}/\Delta(\bar{f})) \},$$

where $\bar{f} : \bar{X} \rightarrow \bar{Y}$ ranges over all holomorphic fibrations $\bar{f}$ between manifolds $\bar{X}$ and $\bar{Y}$ which are equivalent to f .

We call $\kappa(Y, f)$ the **Kodaira dimension of the fibration f** . Notice indeed that this notion depends only on the equivalence class of f .

Definition 1.6. *We shall say that $f : X \rightarrow Y$ is **admissible** if it is holomorphic, with X, Y smooth, and if $\kappa(Y, f) = \kappa(Y, K_Y + \Delta(f))$.*

We shall see in section 1.4 that neat fibrations are admissible. In 1.12, we shall see that any $f : X \rightarrow Y$ is admissible if $\kappa(Y) \geq 0$.

Notice that it follows from 1.7 right below that if f' dominates f , and if f is admissible, so is f' .

1.3.2. Generically Finite Maps. We consider a commutative diagram of holomorphic surjective maps between compact irreducible normal complex spaces :

$$\begin{array}{ccc} X' & \xrightarrow{u} & X \\ f' \downarrow & & \downarrow f \\ Y' & \xrightarrow{v} & Y \end{array}$$

We assume moreover that f, f' are holomorphic fibrations and u, v generically finite (and such that $f \circ u = v \circ f'$).

Theorem 1.7. *Let $f : X \rightarrow Y$ and $f' : X' \rightarrow Y'$ be two holomorphic fibrations and let holomorphic maps $u : X' \rightarrow X$ and $v : Y' \rightarrow Y$ such that $f \circ u = v \circ f'$ be given.*

1. *Assume also that u, v are bimeromorphic, then*
 - a. $\kappa(Y/\Delta(f \circ u)) = \kappa(Y/\Delta(f))$ and $\kappa(Y'/\Delta(f')) \leq \kappa(Y/\Delta(f))$,
 - b. *If, moreover, $\kappa(Y) \geq 0$, then equality holds, and $\kappa(Y/\Delta(f)) = \kappa(Y, f)$.*
2. *Assume that u and v are generically finite and surjective. Then:*
 - a. $\kappa(Y', f') \geq \kappa(Y, f)$,
 - b. $\kappa(Y', f') = \kappa(Y, f)$ *if u is étale, and X, X' are smooth.*

We shall prove Theorem 1.7 as a consequence of several lemmas, some of independent interest. A different, shorter, proof of 1.7 will be given in section 1.4 below.

Lemma 1.8. *Assume u, v bimeromorphic. Then:*

- a. $\Delta(f \circ u) = \Delta(f)$ and so $\kappa(Y/\Delta(f \circ u)) = \kappa(Y/\Delta(f))$,
- b. $\Delta(f') = v^*(\Delta(f)) + E$, *for some $\mathbb{Q}$ -divisor E supported on the exceptional locus of v .*
Therefore $v_(\Delta(f')) = \Delta(f)$ and $\kappa(Y'/\Delta(f')) \leq \kappa(Y/\Delta(f))$.*

Proof:

(a) For any $D \subset Y$, using notations of 1.1.4, we have:

$$(f \circ u)^*(D) = u^*(f^*(D)) = u^*(\sum_{j \in J(D, f)} m_j \cdot D_j + R) = \sum_{j \in J(D, f)} u^*(m_j \cdot (\overline{D}_j + R_j)) + u^*(R),$$

where $\overline{D}_j$ is the strict transform by u of D_j , and R_j is the u -exceptional part of $u^*(D_j)$.

Obviously, $u^*(R)$ is $(f \circ u)$ -exceptional. For each component R_{jk} of R_j , its multiplicity in $(f \circ u)^*(D)$ is divisible by m_j , by the above equalities (and the factoriality of the smooth X). Thus $m(D, f \circ u) = m(D, f)$, and $\Delta(f \circ u) = \Delta(f)$, as claimed. (Observe that R_{jk} does not need to be $(f \circ u)$ -exceptional, in general).

(b) By (a), we can and shall assume that $X = X'$, to ease notations. Let $D' \subset Y'$ be an irreducible divisor, and let $D := v(D')$. We just need to show that $m(D, f) = m(D', f')$ if D' is not u -exceptional, that is: if D is a divisor of Y . Then: $v^*(D) = D' + E$, with E an effective v -exceptional divisor of Y' . Thus $\sum_{j \in J(D, f)} m_j \cdot D_j + R = f^*(D) = (f')^*(D') + (f')^*(E)$. Now observe that $(f')^*(E)$ is f -exceptional, and that each D_j is mapped surjectively onto D by f , and so must be surjectively mapped to D' by f' . Thus: $\sum_{j \in J(D, f)} m_j \cdot D_j = \sum_{j \in J(D', f')} m_j \cdot D'_j$, and we get the claim.

We thus get the first assertion of (b). The others easily follow from it: write $E = E^+ - E^-$, with E^+ and E^- effective and v -exceptional. We thus get: $\kappa(Y'/\Delta(f')) \leq \kappa(Y', K_{Y'} + v^*(\Delta) + E^+) = \kappa(Y'/v^*(\Delta)) = \kappa(Y/\Delta)$, and so the conclusion.

Remark 1.9. *Modifications of X thus don't alter $\kappa(Y/\Delta(f))$, but modifications of the base may let it decrease. See the example below, which shows that strict inequality can actually occur in Theorem 1.7*

Example 1.10. *Probably the simplest example is when $Y = \mathbb{P}^2$ and Δ_{red} is a union of $2k \geq 6$ distinct lines meeting at one point $a \in \mathbb{P}^2$. Corresponding fibrations are easily constructed, as follows: Let $p : Y^+ \rightarrow Y$ be the double cover branched exactly along Δ_{red} , and h the involution of Y^+ exchanging the sheets of p . Let E be an elliptic curve and t a translation of order 2 of E . Let $X^+ := E \times Y^+$, $j := t \times h$ the diagonal involution of X^+ , and $X_0 := X^+/G$, where G is the group of order 2 generated by j . Let $f_0 : X_0 \rightarrow Y$ and $f^+ : X^+ \rightarrow Y^+$ be the induced holomorphic fibrations. Let $d : X \rightarrow X_0$ be a desingularisation (induced by a desingularisation of Y^+ , for example), and $f := d \circ f_0 : X \rightarrow Y$.*

Then $\Delta(f) = (1/2)\Delta_{red}$, and so $\kappa(Y/\Delta(f)) = -\infty$ (resp. 0; resp. 2) if $k \leq 2$ (resp. $k = 3$; resp. $k \geq 4$). On the other hand, let $v : Y' \rightarrow Y$ be the blow-up of $a \in \mathbb{P}^2$, let E be its exceptional divisor, and $f' : X' \rightarrow Y'$ be the lifting of the meromorphic map $v^{-1} \circ f : X \rightarrow Y'$ to a suitable modification X' of X . An easy local computation (for example) shows that $\Delta(f') = \bar{\Delta}$, the strict transform of $\Delta(f)$ by v . (This is just because the normalisation of $(Y^+ \times_Y Y')$ does not ramify over the generic point of E). We thus conclude that $\kappa(Y'/\Delta(f')) = -\infty$, whatever $k \geq 1$. (Because $(K_{Y'} + \Delta(f')) = (k-3)(v^*(H) - E) - 2E$, if H is the hyperplane line bundle on Y , and $v^*(H) - E$ defines the unique ruling of Y'). In particular, $\kappa(Y'/\Delta(f')) < \kappa(Y/\Delta(f))$ if $k \geq 3$. Finally, we observe that the same construction does not lead to this last (strict) inequality when $\Delta(f)$ is a normal crossing divisor (but such other examples should exist).

Corollary 1.11. *Let $f : X \dashrightarrow Y$ a fibration between irreducible compact complex spaces X and Y .*

1. *there exists $\bar{f} : \bar{X} \rightarrow \bar{Y}$ an **admissible** holomorphic model of f between manifolds $\bar{X}$ and $\bar{Y}$, and bimeromorphic maps $u : \bar{X} \rightarrow X$ and $v : \bar{Y} \rightarrow Y$ such that $f \circ u = v \circ \bar{f}$,*
2. *for any X' bimeromorphic to X , there exists $u : X'' \rightarrow X'$ bimeromorphic and $f'' : X'' \rightarrow Y''$, a holomorphic **admissible and prepared** model of f between manifolds X'' and Y'' such that any effective divisor $B'' \subset X''$ which is f'' -exceptional, is also u -exceptional.*

Proof: It is an easy application of Hironaka smoothing and Hironaka-Raynaud Flattening theorems.

(1) First take an admissible model $\bar{f} : \bar{X} \rightarrow \bar{Y}$ of f . Modify $\bar{Y}$ to Y' , smooth and dominating Y . Making base change by Y' over Y , and smoothing $X \times_{\bar{Y}} Y'$, we get the fibration f' , with the desired properties.

(2) Start with the previous fibration f' . We can first modify X' so that the new X' dominates any given bimeromorphic model of X . Then flatten the new f' by modifying Y' , to get Y'' , smooth and the non-smooth locus of f' contained in a divisor of normal crossings. As before, make base change of Y'' over Y' , and smooth $X' \times_{Y'} Y''$ to obtain a fibration $f'' : X'' \rightarrow Y''$ enjoying the claimed properties $\square$

Proposition 1.12. *Assume that $\kappa(Y) \geq 0$ in Lemma 1.8 above. Then:*

$$\kappa(Y/\Delta(f)) = \kappa(Y'/\Delta(f')) = \kappa(Y, f).$$

(In other words: any holomorphic model of f is then admissible).

Proof: Because Y is smooth, using the notations of the proof of Lemma 1.8, we have: $v^*(\Delta(f)) - bE \leq \overline{\Delta(f)} \leq \Delta(f')$, for some nonnegative rational number b , with $\overline{\Delta(f)}$ denoting the strict transform of $\Delta(f)$ by v . Also, here E denotes the reduced exceptional divisor of the map v , and the inequality: $D \leq D'$ between two $\mathbb{Q}$ -divisors D, D' on Y' means that their difference $(D' - D)$ is an effective $\mathbb{Q}$ -divisor.

Moreover, we have: $K_{Y'} \geq v^*(K_Y) + aE$, for some strictly positive rational number a (here we use the smoothness of Y , but Y having just terminal singularities would be sufficient).

We can thus write, as $\mathbb{Q}$ -divisors:

$$K_{Y'} + \Delta(f') \geq v^*(K_Y) + aE + v^*(\Delta(f)) - bE = v^*(K_Y + \Delta(f)) + (a - b)E.$$

We are thus finished if $(a - b) \geq 0$. So we now assume the contrary: $(a/b) < 1$.

From: $bE \geq v^*(\Delta(f)) - \overline{\Delta(f)}$, we get first: $aE = (a/b)bE \geq (a/b)(v^*(\Delta(f)) - \overline{\Delta(f)})$, and then: $\overline{\Delta(f)} + aE \geq (a/b)v^*(\Delta(f)) + (1 - (a/b))\overline{\Delta(f)}$.

From which we deduce: $(K_{Y'} + \Delta(f')) \geq v^*(K_Y) + \overline{\Delta(f)} + aE \geq D$, with:

$D := (a/b)v^*(K_Y + \Delta(f)) + (1 - a/b)(v^*(K_Y)) + \overline{\Delta(f)} \geq (a/b)v^*(K_Y + \Delta(f))$. Where the last inequality follows from our assumption that $\kappa(Y) \geq 0$ $\square$

Lemma 1.13. *Let the situation be as in Theorem 1.7 above. Assume u and v are generically finite and surjective. Then: $\kappa(Y', f') \geq \kappa(Y, f)$.*

Proof: We need only to show that for any admissible f' as above, we can find a holomorphic representative $f_0 : X_0 \rightarrow Y_0$ of f with: $\kappa(Y_0/\Delta(f_0)) \leq \kappa(Y'/\Delta(f'))$, because then:

$$\kappa(Y, f) \leq \kappa(Y_0/\Delta(f_0)) \leq \kappa(Y'/\Delta(f')) = \kappa(Y', f').$$

Modifying X', X, Y and Y' , we can and shall thus assume f' to be admissible.

Let $v = v'' \circ v'$ be the Stein factorisation of v , with $v' : Y' \rightarrow Y''$ connected (hence bimeromorphic), $v'' : Y'' \rightarrow Y$ finite, and Y'' normal.

We also denote here with $\Delta := \Delta(f)$, $\Delta' := \Delta(f')$, $\Delta'' := \Delta(f'')$, $\Delta_0 := \Delta(f_0), \dots$ the relevant multiplicity divisors for the corresponding fibrations $f, f', f'' := v' \circ f', f_0, \dots$

Let now $w : Y'' \rightarrow Y'_0$ be a modification, and define $f'_0 := w \circ f'' := w \circ v' \circ f'$, $w_o := w \circ v' : Y' \rightarrow Y'_0$.

The relevant diagram is the following:

$$\begin{array}{ccccc}
 X' & \xrightarrow{u} & X & & \\
 \downarrow f' & \searrow f'' & \downarrow f & & \\
 Y' & \xrightarrow{v'} & Y'' & \xrightarrow{v''} & Y \\
 & \searrow w_o & \downarrow w & \nearrow & \\
 & & Y'_0 & &
 \end{array}$$

(Note: A curved arrow labeled f'_0 goes from X' to Y'_0 .)

Lemma 1.14. *We can and shall further assume, modifying Y, X and X', Y' if needed, that such a w exists, with f'_0 admissible equivalent to f' .*

Proof Indeed: this existence follows from the next lemma, applied to our initial Y'_0, Y_0 in place of V, U , where f'_0 dominates f_0 (which means: there exists g, h such that $f_0 \circ h = g \circ f'_0$, and $f_0 : X_0 \rightarrow Y_0$ is equivalent to f , with $g : Y'_0 \rightarrow Y_0$ and $h : X'_0 \rightarrow X_0$ generically finite):

Lemma 1.15. *Let $r : V \rightarrow U$ be generically finite surjective, between irreducible normal compact complex spaces. There exist modifications $n : V' \rightarrow V$ and $m : U' \rightarrow U$ such that:*

(1) $s := m^{-1} \circ r \circ n : V' \rightarrow U'$ is holomorphic

(2) if $s = r'' \circ r'$ is the Stein factorisation of s , then there exists a (holomorphic) factorisation $n'' : V'' \rightarrow V$ of $m \circ r'' : V'' \rightarrow U$ through r (ie: $r \circ n'' = m \circ r''$). Here $r' : V' \rightarrow V''$ is connected and $r'' : V'' \rightarrow U'$ is finite $\square$

We can assume V' and U' to be smooth.

The relevant diagram is the following:

$$\begin{array}{ccccc}
 & & n & & \\
 & \nearrow & & \searrow & \\
 V' & \xrightarrow{r'} & V'' & \xrightarrow{n''} & V \\
 & \searrow s & \downarrow r'' & & \downarrow r \\
 & & U' & \xrightarrow{m} & U
 \end{array}$$

Proof: Just flatten the map r by suitably modifying U $\square$

To complete the proof of 1.13, we shall next need the following two properties **(a)**, **(b)**:

(a) We then have (because v'' is finite):

(*a) $K_{Y''} + \Delta'' = (v'')^*(K_Y + \Delta) + R''$, an equality between Weil Divisors, and R'' effective.

To see this, observe we may first assume that f' is deduced from f by base change by v and smoothing of a main component (this is simply because $\Delta(f \circ u) \geq \Delta(f)$ if $Y' = Y$, and u is generically finite. The argument is the same as in 1.8).

Next, a simple computation in local analytic coordinates over the generic point y'' of Y'' shows that: if r is the ramification order of v'' at y' and m is the multiplicity of the fibre of f over $y := v''(y'')$, then the multiplicity of the f'' -fibre over y'' is $m' := (m/d)$, if $d := \gcd(r, m)$ and $r' := r/d$

For this, we need to choose y'' lying outside the codimension two analytic subset of Y'' consisting of points y'' which are either singular on Y'' , or mapped by v'' to a point $y := v''(y'')$ either singular on $\Delta(f) \cup S$, or with f -fibre of nongeneric multiplicity, where $S := v''(S')$, with S' the Weil divisor of points of Y'' at which v'' ramifies.

From this we see that near y'' on Y'' , we have:

$K_{Y''} + \Delta'' = (v'')^*(K_Y + (1 - 1/r).R) + (1 - 1/m').(1/r).D$, with $R := v''(R^+)$ ($S'' = R^+$ the reduced ramification divisor of v'' near y''), and D the (unique) reduced component of Δ near y . If R^+ or D is empty, our equality (*a) is obvious. Otherwise, $R = D$, by our choice of y'' , and (*a) follows from the inequality: $(1 - 1/m) \leq (1 - 1/m.r') = (1 - 1/m'.r) = (1 - 1/r) + (1 - 1/m').1/r$.

(b) Moreover, we also have (quite generally): $K_{Y''} + \Delta'' = (v')_*(K_{Y'} + \Delta')$. This is an immediate check.

From the initial constructions made, we thus know that f'_0 is admissible, and dominates f_0 , admissible (in the sense (**)) just before the lemma).

We then get: $K_{Y''} + \Delta'' = (v')_*(K_{Y'} + \Delta') = (v'')^*(K_Y + \Delta) + R''$.

Hence: $K_{Y'_0} + \Delta'_0 = (w)_*(K_{Y''} + \Delta'') = (w)_*((v'')^*(K_Y + \Delta) + R'') = (w_0)_*(v^*(K_Y + \Delta) + R')$.

And so: $\kappa(Y', f') = \kappa(Y'_0, K_{Y'_0} + \Delta'_0) \geq \kappa(Y'_0, (w_0)_*(v^*(K_Y + \Delta))) \geq \kappa(Y', v^*(K_Y + \Delta)) = \kappa(Y, K_Y + \Delta) \geq \kappa(Y, f)$, as claimed in 1.13, the proof of which is thus complete $\square$

The last property asserted in Theorem 1.7 is:

Lemma 1.16. *Let the situation be as in Theorem 1.7 above. Assume now that $u : X' \rightarrow X$ is étale (ie: unramified). Then: $\kappa(Y', f') = \kappa(Y, f)$.*

Proof: By Lemma 1.13, we need only to show that $\kappa(Y', f') \leq \kappa(Y, f)$ for any f, f' as above.

We can assume f to be admissible, and still u to be étale: indeed, just replace X' by $X'_1 := X' \times_X X_1$, if $m : X_1 \rightarrow X$ is a modification having an admissible fibration $f_1 : X_1 \rightarrow Y_1$ equivalent to f . Then X'_1 is smooth, and $u_1 : X'_1 \rightarrow X_1$ is étale.

The Stein factorisation $f'_1 : X'_1 \rightarrow Y'_1$ of $f_1 \circ u_1 : X'_1 \rightarrow Y_1$ does not need to have Y_1 smooth. But a modification $m' : X'_2 \rightarrow X'_1$ of X'_1 allows to assume this to be true (accordingly modifying Y_1). Now the birational invariance of the fundamental group for complex manifolds shows that X'_2 may be assumed (after further modification) to be of the form $X' \times_X X_2$ for some modification X_2 of X . Let us check this.

We can indeed assume that $u : X'_1 := X' \rightarrow X$ is Galois, of group G , and that f' is the (connected part of the) Stein factorisation of $(f \circ u)$. The map $v : Y' \rightarrow Y$ is thus G -equivariant as well. But then, the modification $m' : X'_2 \rightarrow X'_1$ can be assumed to be also G -equivariant, which shows that X'_2 is obtained from u by base-change over a modification of X . This establishes our preliminary claim (that f can be chosen admissible).

Next, by Stein factorisation of $v = v'' \circ v'$, we see as in the above proof of Lemma 1.7, that it is sufficient to show that:

(*) $K_{Y''} + \Delta'' = (v'')^*(K_Y + \Delta)$ (that is: R'' is empty).

Because then $K_{Y''} + \Delta'' = (v')_*(K_{Y'} + \Delta')$, and so $K_{Y'} + \Delta' = (v)^*(K_Y + \Delta) + E'$, for some v -exceptional (not necessarily effective) $\mathbb{Q}$ -divisor E' of Y' , which implies the desired reverse inequality.

By the finiteness of the map v'' , the equality (*) has only to be shown near any point $y \in Y$ lying outside a codimension two subset S of Y . We can thus assume that y is a generic point on some component of the support of $\Delta(f)$. Let us now cut Y by divisors in general position through y : we are reduced to the case when Y is a curve (the argument being local on Y in the analytic topology, we don't need any algebraicity assumption of Y). But then an easy local computation (see [C98], for example) shows that for any $y'' \in Y''$, the order of ramification r of v'' at y'' divides the multiplicity m of the fibre X_y of f over $y := v''(y'')$. The multiplicity m'' of the fibre $X'_{y''}$ of f' over y'' is thus $m' := m/r$, since u is étale.

We now compute $K_{Y''} + \Delta''$ near y'' : $K_{Y''} + \Delta'' = (v'')^*(K_Y + (1 - 1/r)[y]) + (1 - 1/m')[y''] = (v'')^*(K_Y + ((1 - 1/r) + (1/r) \cdot (1 - 1/m))[y]) = (v'')^*(K_Y + (1 - 1/m)[y])$, as claimed. (Here $[y]$ is the reduced divisor on the curve Y supported by the point y) $\square$

1.4. The Sheaf of Differential Forms determined by a fibration.

Definition 1.17. *Let $X \in \mathcal{C}$ be smooth, and let $f : X \dashrightarrow Y$ be a meromorphic fibration, with Y reduced but not necessarily smooth. The rank one coherent subsheaf F_f of $\Omega_{X,p}^p := \dim(Y)$ is defined as the saturation in Ω_X^p of $f^*(K_{Y_0})$, if Y_0 is the smooth locus of Y .*

Let us remark that the subsheaf F_f so defined is a bimeromorphic invariant: it is preserved not only under modifications of X , but also under modifications of Y . In other words: F_f depends only on the equivalence class of f .

We define by $\kappa(f)$ its Kodaira dimension.

Let us make more precise what is understood by $\kappa(X, F_f)$: for $m > 0$, define $H^0(X, F_f^m)$ to be the complex vector space of sections of the subsheaf of $(\Omega_X^p)^{\otimes m}$ which coincides with $F_f^{\otimes m}$ over the Zariski open subset of X (with codimension two or more complement) over which F_f is locally free. Then define κ in the usual way.

To see the usual property of κ (being an integer, or $-\infty$), notice that the data are bimeromorphically invariant on X . So that F_f can be considered as the injective image of some locally free rank one sheaf L on X (after some suitable modification). The claimed property is then obvious, since it holds for L . We can then always implicitly assume the existence of L in the sequel, and also the holomorphic character of any meromorphic map defined on X (such as, for example, the ones defined by linear systems $L^{\otimes m}$).

We shall now describe F_f in more detail.

Definition 1.18. *In the preceding situation, define $F(f) := f^*(K_X) \otimes \mathcal{O}_X(\lceil f^*(\Delta(f)) \rceil)$, where the symbol used is the usual round-up (defined as $\lceil * \rceil := -[-(*)]$, applied to the coefficients of the irreducible components of the effective $\mathbb{Q}$ -divisor under consideration. Here $\lceil * \rceil$ is the integral part, of course).*

Definition 1.19. *Let $f : X \rightarrow Y$ be a holomorphic fibration, and S an effective divisor on X . We say that S is **partially supported on the fibres of f** if: $f(S) \neq Y$, and if T is any irreducible component of $f(S)$ of codimension one in Y , then $f^{-1}(T)$ contains an irreducible component mapped on T by f , but not contained in the support of S .*

Observe that if S is partially supported on the fibres of f , so are its positive multiples.

The introduction of this notion is due to the following:

Lemma 1.20. *Let $f : X \rightarrow Y$ be a fibration, and S a divisor of X partially supported on the fibres of f . Let L be a line bundle on Y . The natural injection of sheaves $L \subset f_*(f^*(L) + S)$ is an isomorphism.*

Proof: The assertion is of local nature on Y . So we can assume that L is trivial. We then just need to show that $f_*(\mathcal{O}_X(S)) \cong \mathcal{O}_Y$. We assume that $S \subset f^*(\mathcal{O}_Y(T))$, for some effective divisor T on Y . Local sections of the sheaf on the right are of the form $f^*(u/t)$, where u is holomorphic on Y , while t is a local equation of T . The sections of the sheaf on the left are meromorphic functions on X of the same form, but with poles contained in S . Because S is partially supported on the fibres of f , we get the claim (t divides u) $\square$

We apply this to the following situation:

Proposition 1.21. *Let $f : X \rightarrow Y$ be a holomorphic fibration between manifolds. There exists a Zariski closed subset $A \subset Y$ of codimension at least 2 such that $F(f) + S$ and F_f are naturally isomorphic over $(X - B) := f^{-1}(Y - A)$, where $S \subset X$ is an effective divisor partially supported on the fibres of f .*

Proof: Let A be the union of the singular set of the support of $\Delta(f)$ and of images of all f -exceptional divisors on X . Let us remark that the above natural isomorphism is immediate outside of Δ , because if $T \subset Y$ is a one-codimensional component of the locus of non-smooth fibres of f , then $f^{-1}(T)$ contains a reduced component at the generic point of which f is smooth. So we consider the situation near a smooth point of some Δ_i not lying in A .

In suitable local coordinates at the generic point of D_{ij} , in the notations of the lines preceding 1.1.4, we have:

$(x) = (x_1, \dots, x_n)$, and $(y) = (y_1, \dots, y_p)$, with: $f(x) = (y_1 := x_1^{m_{ij}}, y_2 := x_2, \dots, y_p := x_p)$. And so: $[f^*(K_Y + \Delta(f))]$ is generated by $x_1^{[(m_{ij}/m_i)-1]} \cdot d(x')$, with $d(x') := dx_1 \wedge \dots \wedge dx_p$. A simple check shows that this is exactly the claim. (One may even observe that the divisor S has the same description as the one given to define $F(f)$, by adding to Δ the one-codimensional components of the locus of non-smooth fibres of f) $\square$

Corollary 1.22. *Let $f : X \rightarrow Y$ be a fibration as in 1.21 above. Let $m > 0$ be a sufficiently divisible integer. Then:*

- (1) *The natural isomorphism between $F(f) + S$ and F_f over $(X - B)$ extends to a natural injection of $H^0(X, F_f^{\otimes m})$ into $H^0(X, m \cdot (F(f) + S)) \cong H^0(Y, m \cdot (K_Y + \Delta(f)))$.*
- (2) *If f is neat, this injection is bijective.*

Proof: (1) We start by observing that, by 1.20, the bijection:

$$H^0(X, m \cdot (F(f) + S)) \cong H^0(Y, m \cdot (K_Y + \Delta(f))) \text{ actually holds.}$$

The natural map at the level of sections of m -th powers induces an isomorphism over a codimension two subset of Y . Because $m \cdot (K_Y + \Delta(f))$ is locally free on Y , the said isomorphism thus extends as an injection, by Hartog's theorem.

(2) This is because B is mapped to a codimension two or more Zariski closed subset of X' if $u : X \rightarrow X'$ is a modification with X' smooth and sending the f -exceptional divisors of X in codimension 2 or more in X' . Then the sections of $m \cdot (F(f) + S)$ over $(X - B)$ extend to sections of $F_f^{\otimes m}$, as claimed $\square$

We now have the following easy but important consequence:

Proposition 1.23. *Let $f : X \rightarrow Y$ be a holomorphic fibration, with X, Y smooth. Then:*

- (1) $\kappa(f) \leq \kappa(Y/\Delta(f))$
- (2) $\kappa(f) = \kappa(Y/\Delta(f))$ if f is neat (see 1.2 for this notion).
- (3) $\kappa(f) = \kappa(Y, f)$

Proof: (1) and (2) are simply restatements of 1.22. We deduce (3) by choosing a neat admissible model f' of f . Then: $\kappa(f) = \kappa(Y'/\Delta(f')) = \kappa(Y, f)$ $\square$

This allows us to give a short proof of the basic properties shown in Theorem 1.7:

Corollary 1.24. *The statements of Theorem 1.7 hold.*

Proof: Using the notations there, we have a natural inclusion: $u^*(F_f) \subset F_{f'}$, which is an equality if u is étale. the conclusions follow from the standard properties of the Kodaira dimension $\square$

Remark that one can also use the sheaves F_f to simplify some of the geometric proofs given in section 2.2

1.5. Semi-continuity of the Kodaira Dimension.

Proposition 1.25. *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be holomorphic fibrations, with X a connected compact complex manifold, and $h := g \circ f$. Let Z^* be the Zariski open subset of Z over which g and h are smooth. Let, for $z \in Z$, denote by $f_z : X_z \rightarrow Y_z$ the restriction of f to the z -fiber X_z of h , mapped by f to the z -fibre Y_z of g . Then:*

1. *There exist modifications $\mu : X' \rightarrow X$ and $\nu : Y' \rightarrow Y$ such that f'_z is admissible, for z general in Z , with $f' := \nu^{-1} \circ f \circ \mu$ holomorphic.*
2. *Let $d := \inf(\{\kappa(Y_z, f_z), z \in Z^*\})$. (So that $d \in \{-\infty, 0, 1, \dots, \dim(Y) - \dim(Z)\}$). Then: $d = \kappa(Y_z, f_z)$, for z general in Z .*
3. *Let A be the set of points z of Z^* such that $\kappa(Y_z, f_z) = \dim(Y) - \dim(Z)$. Then: either A contains the general point of Z , or A is contained in a countable union of closed proper analytic subsets of Z .*

Proof: We don't mention modifications of X , since they don't change the Kodaira dimensions of fibrations with X as domain. For any modification $\nu : Y' \rightarrow Y$ and $d \in \{-\infty, 0, 1, \dots, \dim(Y) - \dim(Z)\}$, let $S_d^*(\nu) := \{z \in Z^* \text{ such that: } \kappa(Y_z, \Delta(f'_z)) \geq d\}$. From [Gr60], and as in 5L-S], we deduce that $S_d^*(\nu) = S_d(\nu) \cap Z^*$, where $S_d(\nu)$ is a countable union of Zariski closed subsets of Z . Obviously, $S_{d+1}(\nu) \subset S_d(\nu)$ for any ν and d (this on Z^* at least, which is sufficient for our purposes).

If $\nu' : Y'' \rightarrow Y$ dominates ν in the sense that there exists a $\nu'' : Y'' \rightarrow Y'$ with $\nu' = \nu'' \circ \nu$, then obviously (by Theorem 1.7): $S_d(\nu') \subset S_d(\nu)$.

We can thus define, for any d , $S_d \subset Z$ as the intersection of all $S_d(\nu)$'s: it is again a countable intersection of Zariski closed subsets of Z .

Define now $d := \max\{d' \text{ such that } S_{d'} = Z\}$. There thus exists some ν such that $S_{d+1}(\nu) \neq Z$. Both claims then follow immediately from the constructions just made.

The third assertion is an immediate consequence of the second.

1.6. Composition of Fibrations.

This section will not be used before section 4.

Assume now X, Y, Z to be smooth and $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ to be holomorphic fibrations.

Our aim is to define, if H is an orbifold structure on Y (ie: an effective $\mathbb{Q}$ -divisor on Y with components having multiplicities of the form $(1 - 1/m)$, for m integer), an orbifold structure $\Delta(g, H)$ on Z in such a way that we have the equality: $\Delta(g, H) = \Delta(g \circ f)$ when $H = \Delta(f)$, if $f : X \rightarrow Y$ is sufficiently "high", in a sense defined in section 4 below.

Let us recall how we defined multiplicities: for any irreducible divisor $D \subset Z$, the multiplicity $m(g, D)$ of g along D is: $m(g, D) =: \inf\{m_j, j \in J(g, D)\}$. Where $g^+(D) := (\sum_{j \in J(g, D)} m_j \cdot D_j)$ is the union of all components (counted with their scheme-theoretic multiplicities) of $g^*(D)$ which are mapped surjectively onto D by g , and $J(g, D)$ is the set of these components.

Let, if $D' \subset Y$ is an irreducible divisor of Y mapped by g to a divisor $D = g(D') \subset Z$, $m_*(g, D')$ be the multiplicity of D' in $g^*(D)$. Thus $m(g, D) = (\sum_{j \in J(g, D)} m_*(g, D_j) \cdot D_j)$.

We shall now define the notion of **orbifold base of a fibration** $g : (Y/H) \rightarrow Z$, when the domain of the fibration is itself an orbifold (Y/H) . But $g : Y \rightarrow Z$ is just a usual fibration, with Z smooth.

Writing $H := \sum_{i \in I} (1 - 1/m_i) \cdot H_i$, define, for any irreducible reduced divisor $D' \subset Y$ its multiplicity $m(H; D')$ in H as being m_i if $D' = H_i$, and being 1 otherwise (ie: if D' is not a component of the support of H).

For any irreducible divisor $D \subset Z$, with Z smooth, define next the **multiplicity** $m(g, H; D)$ of $g : (Y/H) \rightarrow Z$ **along** D by: $m(g, H; D) := \inf_{j \in J(g, D)} \{m_*(g, D_j) \cdot m(H; D_j)\}$.

And, finally:

Definition 1.26. Let $g : Y \rightarrow Z$ be a fibration, with Z smooth. Let H be an orbifold structure on Y . We define the **orbifold base** $\Delta(g, H)$ of the fibration $g : (Y/H) \rightarrow Z$ by:

$$\Delta(g, H) := \sum_{D \subset Z} (1 - 1/m(g, H; D)) \cdot D.$$

In general, it is not true that $\Delta(g, H) = \Delta(g \circ f)$ if $H = \Delta(f)$. But the following results at least are available:

Proposition 1.27. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be two holomorphic fibrations, with X, Y, Z smooth. Then: $\Delta(g \circ f) \leq \Delta(g, \Delta(f))$.

(Recall that, for two $\mathbb{Q}$ -divisors A, B on a variety Z , we write $A \leq B$ if $(B - A)$ is effective.)

Proof: We easily check that we have, for any prime divisor $D \subset Z$:

$m(g, \Delta(f); D) = \inf_{j \in J(g, D)} \{m_*(g, D_j) \cdot m(\Delta(f); D_j)\}$, by definition; and:

$m(g \circ f, D) = \inf_{k \in J(g \circ f, D)} \{m(g \circ f, D_k)\}$, also by definition.

Now, if $f(D_k) = D_j$ is a divisor, then: $m_*(g \circ f, D_k) = m_*(g, D_j) \cdot m(\Delta(f); D_j)$, by an easy check.

Observe that the minimum of these values, taken over $J(g, D)$, is precisely $m(g, \Delta(f); D)$.

Thus: $m(g \circ f, D) = \inf \{m(g, \Delta(f); D), m(f, g)\}$, where: $m(f, g) := \inf_{k \in J'(g \circ f, D)} \{m_*(g \circ f, D_k)\}$, and where $J'(g \circ f, D)$ is the set of irreducible components of $(g \circ f)^*(D)$ which are surjectively mapped to D by $(g \circ f)$, but are f -exceptional. From this the claim follows.

Observe that strict inequality may thus occur. However:

Proposition 1.28. Let f, g be two fibrations, as above. Let $f' : X' \rightarrow Y'$ be a modification of f (with modifications $u : X' \rightarrow X$ and $v : Y' \rightarrow Y$ such that: $f \circ u = v \circ f'$). Then:

1. $\Delta(g' \circ f') = \Delta(g \circ f)$, with $g' := g \circ v$.
2. $\Delta(g', \Delta(f')) \leq \Delta(g, \Delta(f))$
3. There exists a modification f' of f such that: $\Delta(g', \Delta(f')) \leq \Delta(g'', \Delta(f''))$, for any modification f'' of f' .
4. If f' is a modification as in (3) above, then: $\Delta(g' \circ f') = \Delta(g', \Delta(f'))$.
5. In (3) above, we can choose f' such that, moreover, f', g' and $(g' \circ f')$ are prepared, admissible and high.

Proof:

(1) $g' \circ f' = (g \circ f) \circ u$. So, the claim follows from 1.7.

(2) We can and shall thus assume that $X' = X$ and $f = v \circ f$. Let $D \subset Z$ be a prime divisor. Then $J(g, D) \subset J(g', D)$, the difference consisting of the v -exceptional components of $(g')^*(D)$. Moreover, for each $j \in J(g, D)$, we have: $m(g, D_j) = m(g', \overline{D_j})$, with $\overline{D_j}$ the strict transform of D_j by v . Finally: $m(f, D_j) = m(f', \overline{D_j})$, by 1.7. This implies the claim.

(3) This is simply because there are only finitely many orbifold divisors on Z lying between $\Delta(g \circ f)$ and $\Delta(g, \Delta(f))$, and because of (1) and (2) above, which show that the first (resp. second) term is invariant (resp. decreases) under a modification.

(4) Assume indeed that f (rather than f' to ease notations) enjoys the property stated in (3). Assume we have $\Delta(g \circ f) < \Delta(g, \Delta(f))$. This means that there exists an f -exceptional prime divisor $D' \subset X$ such that $g(f(D')) := D \subset Z$ is a divisor in Z , and that the multiplicity m' of D' in $(g \circ f)^*(D)$ is equal to $m(g \circ f, D)$, and is so strictly less than $m(g, \Delta(f); D)$.

Take a modification f' of f such that the strict transform of D' in X' is no longer f' -exceptional. The multiplicity of D'' , the strict transform of D' in X' is then m' (by an easy check). Thus: $m(g', \Delta(f'); D) \leq m' < m(g, \Delta(f); D)$. But this contradicts the property (3), supposed to hold for f . Contradiction; the claim follows.

(5) By modifying Z , we can assume that $(g \circ f)$ and g are admissible, high, and moreover that the non-smooth loci of these two fibrations are contained in a normal crossings divisor. By modifying next Y , we can assume that g is prepared, f admissible and high, and that the non-smooth locus of f is contained in a normal crossings divisor of Y . Finally modify X to get the remaining stated properties $\square$

2. Special Fibrations and General Type Fibrations

2.1. Special or General Type Fibrations.

Definition 2.1. Let $f : X \dashrightarrow Y$ be a fibration, with X, Y compact irreducible.

1. The fibration $f : X \dashrightarrow Y$ is said to be of **general type** if $\kappa(Y, f) = \dim(Y) > 0$.
2. The variety X is said to be **special** if $X \in \mathcal{C}$ and if there is no meromorphic fibration $f : X \dashrightarrow Y$ of general type, for any Y .
3. The fibration $f : X \dashrightarrow Y$ is said to be **special** if $X \in \mathcal{C}$, and if its general fibre is special.

Recall that here $\mathcal{C}$ is the class of compact complex spaces X which are bimeromorphic to (or, equivalently: dominated by) some compact Kähler manifold X' (depending on X). This class was introduced by A.Fujiki.

Recall (see [C81]) also that a point of a complex space Y is said to be **general** if it lies outside of a countable union of closed analytic subsets of Y , none of which containing any irreducible component of Y . Similarly, if $f : X \rightarrow Y$ is a fibration, one of its fibres X_y is general if it lies above a general point y of Y .

Example 2.2.

We list (proofs need tools developed below, and so are given later) some examples of special manifolds.

0. A variety of general type (and positive dimension) is **not** special. (Consider its identity map: it is a fibration of general type).
1. A **curve** is special iff its genus is 0 or 1, iff its Kodaira dimension is at most zero, iff its fundamental group is abelian, iff it is not hyperbolic. This because a curve has only the two trivial fibrations (constant, and identity).

The two fundamental examples of special Manifolds are direct generalisations:

2. A manifold which is **rationally connected** is special. See Theorem 3.21 for a geometric proof and definition of the notions involved. Another shorter (but more abstract) proof of the specialness of rationally connected manifolds is given 2.27 below.
3. A manifold X with **vanishing Kodaira dimension** (ie: $\kappa(X) = 0$) is special. See Theorem 5.1 for the proof. Another proof is given in 2.27 in the special case where $c_1(X) = 0$.
4. More generally, special manifolds are conjectured to be build up from manifolds either rationally connected, or with Kodaira Dimension zero by suitable compositions of fibrations with fibres of these two types. See section 6.4 for a precise formulation.
5. For any $d > 0$ and $k \in \{-\infty, 0, \dots, d-1\}$, there exists projective manifolds of dimension d and Kodaira dimension k . See 2.18 for such examples.
6. A manifold $X \in \mathcal{C}$ is special if there exists a nondegenerate meromorphic map from $\mathbb{C}^n$ to X , where nondegenerate means: submersive at some point where it is holomorphic. For example, a complex torus, or a projective space are special (this follows also from [2,3] above as well). See theorem 8.2 for the proof of a more general version.
- 7 Kähler manifolds with nef anticanonical bundle are conjectured to be special. This conjecture implies most usual conjectures concerning these manifolds. See [D-P-S],[Zh],[Pa].
8. A manifold X of **algebraic dimension zero** (denoted $a(X) = 0$, to mean that all meromorphic functions on X are constant, so that meromorphic maps from X to projective varieties are constant) is also special (simply because any meromorphic map from X onto a projective manifold is constant).
9. More generally (see [U],Chap. 12 for the notions used):

Theorem 2.3. *Let $a_X : X \rightarrow \text{Alg}(X)$ be the algebraic reduction of $X \in \mathcal{C}$. The generic fibre of a_X is special.*

Recall that the algebraic dimension of X , denoted $a(X)$, is the dimension of $\text{Alg}(X)$, and also the transcendence degree over $\mathbb{C}$ of the field of meromorphic functions on X . One says that X is **Moishezon** if $a(X) = \dim(X)$. This also means that X has a modification which is projective.

The proof of the preceding result is given in 2.38 below.

Question 2.4. *Two important stability properties of the class of special manifolds are expected to hold, but are not proved in the present paper: the stability under deformation and specialisation. Do these hold?*

2.2. Special Fibrations Dominate General Type Fibrations. The geometric study of special manifolds is based on Theorem 2.11, stated and proved below. Its proof rests on several preliminary results of independent interest that we now give.

Lemma 2.5. *Let $g : X' \rightarrow X$ be meromorphic surjective (ie: dominant). Assume that X' is special. Then X is special, too.*

Proof: Assume first that g is connected (ie: a fibration). Let, if any, $f : X \rightarrow Y$ be a fibration of general type. We can assume that $f \circ g : X' \rightarrow Y$ is admissible. Then obviously, $\Delta(f \circ g) = \Delta(f) + E$, for some effective divisor $E \subset Y$ (because $m(f \circ g, D) \geq m(f, D)$, for any irreducible divisor

$D \subset Y$). Thus $f \circ g$ is of general type, too. A contradiction. No such f does exist, which is what was claimed.

In the general case, Stein factorise g and use the first part to reduce to the case where g is generically finite. If f as in the first part exists, then we deduce from 1.7 that (the fibration part of) the Stein factorisation of $f \circ g$ also is of general type. Hence again a contradiction $\square$

Proposition 2.6. *Let $f : X \rightarrow Y$ be a fibration of general type. Let $j : Z \rightarrow X$ be holomorphic such that $f \circ j : Z \rightarrow Y$ is surjective. Let $f \circ j = g \circ h$ be the Stein factorisation of $f \circ j$, with $h : Z \rightarrow Y'$ connected and $g : Y' \rightarrow Y$ finite. Then: h is a fibration of general type.*

In particular, if $\dim(Z) = \dim(Y)$, we get:

Corollary 2.7. *Let $f : X \rightarrow Y$ be a fibration of general type. Let $j : Z \rightarrow X$ be meromorphic, and such that $f \circ j : Z \rightarrow Y$ surjective. Then Z is a variety of general type.*

Proof (of 2.6 and 2.7): Assume first that $f \circ j : Z \rightarrow Y$ is connected and admissible (as we can, then). For any component Δ_i of $\Delta := \Delta(f)$, we have (restricting to components surjectively mapped onto Δ_i by $(f \circ j)$): $(f \circ j)^*(\Delta_i) = j^*(f^*(\Delta_i)) = j^*(m_i \cdot D_i) = m_i \cdot j^*(D_i)$.

Thus: $\Delta(f \circ j) = \Delta(f) + E$, for some effective $\mathbb{Q}$ -divisor E of Y . And $(f \circ j)$ is thus of general type in this case.

We now consider the general case: define $f' : X' := (\widetilde{X \times_Y Y'}) \rightarrow Y'$ be deduced from the base change by g and smoothing of the fibre product. The map j lifts meromorphically to $j' : Z \rightarrow Y'$, by construction, because $f \circ j$ is surjective. But now $f' \circ j' : Z \rightarrow Y'$ is a fibration. Applying the first part, we get the claim.

For 2.7, notice that in this situation, h is bimeromorphic and of general type by 2.6. Thus Z is itself of general type, as claimed $\square$

Example 2.8. *We can now give two elementary examples of special varieties:*

1. $\mathbb{P}_n(\mathbb{C})$ is special.
2. A product of special varieties is special.

Proof: Indeed (for (1)): let $f : \mathbb{P}_n(\mathbb{C}) \dashrightarrow Y$ be any general type fibration, if any. Let $m := \dim(Y) > 0$. Choose $j : Z = \mathbb{P}_m(\mathbb{C}) \subset \mathbb{P}_n(\mathbb{C})$ such that $f \circ j$ is surjective to contradict 2.6.

The proof of (2) is similar. (We shall prove more general results in section 3).

Proposition 2.9. *Let $f : X \rightarrow Y$ and $k : Y \rightarrow W$ be fibrations. Assume that f is of general type. Then $f_w : X_w \rightarrow Y_w$ is also of general type, for $w \in W$ general.*

Proof: Let $w \in W$ be general, and recall that $f_w : X_w \rightarrow Y_w$ is nothing, but the restriction of f to X_w . But then, $\Delta(f_w) = \Delta(f)|_{Y_w} + E_w$, with E_w effective and empty for generic w in W . Moreover, $K_{Y_w} = K_Y|_{Y_w}$, by adjunction.

Thus $K_{Y_w} + \Delta(f_w) = (K_Y + \Delta(f))|_{Y_w}$ for general w . Now $(K_Y + \Delta(f))$ is big. Thus so is its restriction to Y_w . By modifying adequately X and Y , we can assume that f_w is admissible by the following lemma. We thus get the claimed property $\square$

Lemma 2.10. *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be fibrations. There exists representatives of f and g (also denoted f and g) such that f_z is admissible, for $z \in Z$ general.*

Proof: This is a special case of 1.25 $\square$

The main result in this section is:

Theorem 2.11. *Let $h : V \dashrightarrow Z$ and $f : X \dashrightarrow Y$ be fibrations with f of general type and h having general fibres which are special. Let $g : V \dashrightarrow X$ be meromorphic surjective. Then, there exists $k : Z \rightarrow Y$ such that $f \circ g = k \circ h$.*

The situation is described in the following commutative diagram:

$$\begin{array}{ccc} V & \xrightarrow{g} & X \\ h=\text{special} \downarrow & & \downarrow f=\text{gen.type} \\ Z & \xrightarrow{k} & Y \end{array}$$

The special case $V = X$ deserves special mention:

Theorem 2.12. *Let $h : X \dashrightarrow Z$ and $f : X \rightarrow Y$ be fibrations with f of general type and h **special**. Then, there exists $k : Z \rightarrow Y$ such that $f \circ g = k \circ h$ (We say that h **dominates** f).*

The corresponding diagram is:

$$\begin{array}{ccc} X & & \\ h=\text{special} \downarrow & \searrow f=\text{gen.type} & \\ Z & \xrightarrow{k} & Y \end{array}$$

Remark 2.13. *The special case where $X = Y$ is of general type is obvious, by the easy addition theorem, because the covering family of Y by the subvarieties $h(X_z)$ has a generic member of general type. These subvarieties must be points.*

Proof (of 2.11): This is a direct consequence of 2.14 below. Indeed, if such a map k does not exist, then $f \circ g(V_z)$ is positive-dimensional, for generic $z \in Z$. But the Stein factorisation of $f \circ g_z$ is then of general type, by 2.14. This contradicts the assumption that g is special $\square$

Proposition 2.14. *Let $f : X \rightarrow Y$ and $h : V \rightarrow Z$ be fibrations. Assume f is of general type. Let $g : V \rightarrow X$ be a surjective meromorphic map. Let $g_z : V_z \rightarrow Y_z$, be the restriction of $f \circ g$ to V_z , with $Y_z := h_z(V_z)$.*

Assume Y_z is positive dimensional. Then: the Stein factorisation of g_z is of general type for z general in Z .

Proof: By 2.9, we can then replace X by V and f by $(f \circ g)$ without losing the hypothesis that $(g \circ g)$ is of general type both when g is generically finite or is a fibration. We thus see that the (connected part of the) Stein factorisation f' of $f \circ g$ is a fibration of general type. Replace X by V and f by f' , so that we are reduced to the case where $X = V$, and $f = (f \circ g)$, which we now treat.

We then notice that we can replace Z by any subvariety $Z' \subset Z$ going through a general point of Z , and X by $X' \subset X$, defined by $X' := g^{-1}(Z')$, provided $f(X') = Y$. This is because of 2.9, which shows that the (Stein factorisation of the) restriction of f to X' is still of general type.

We shall then construct an appropriate $Z' \subset Z$.

Let $c : Z \rightarrow \mathcal{C}(Y)$ be the meromorphic map sending a generic $z \in Z$ to the reduced cycle of Y supported on Y_z . Here $\mathcal{C}(Y)$ denotes the Chow Scheme of Y . Observe next that f being of

general type, Y is Moishezon. By modifying suitably Y , we shall assume that Y is projective. Let $W \in \mathcal{C}(X)$ be the image of Z by c . Thus W is projective, too.

We next choose $W' \subset W$ to be an intersection of generic members of any very ample linear system on W , in such a way that the incidence graph $Y' \subset W' \times Y$ of the algebraic family of cycles of Y parametrised by W' is generically finite over Y . This means that if $p : Y' \rightarrow Y$ and $q : Y' \rightarrow W'$ are induced by the natural projections, then p is generically finite surjective. More concretely, the generic point of Y is contained in only finitely many of the Y_w 's, for $w \in W'$. Define now $Z' := c^{-1}(W')$. (Remark that when Z is Moishezon, we don't need to consider c , and can just take intersections of ample divisors of a projective modification of Z directly).

We now replace X, Z, g, f respectively by $X', Z',$ and their restrictions to X' .

Lemma 2.15. *$p : Y' \rightarrow Y$ is bimeromorphic.*

Proof: Let c' be the restriction of the above map c to Z' . Then $Y' \subset (W' \times Y)$ is the image of $(c' \circ h) \times f : X' \rightarrow W' \times Y$. Thus $f : X' \rightarrow Y$ lifts to $f' : X' \rightarrow Y'$ such that $p \circ f' = f$. Because p is generically finite and f connected, we see that p is bimeromorphic. $\square$

We have, by construction, $q \circ f = c' \circ h : X \rightarrow W'$. We now can conclude by applying successively 2.9 and 2.7 to X_w , for w general in W' . Indeed: from 2.9 we learn that the restriction $f_w : X_w \rightarrow Y_w$ of f to X_w is of general type. Further, for z generic in $c'^{-1}(w) := Z'_w$, we know that $f(X_z) = f(X_w) = Y_w$, and so by 2.7, the Stein factorisation of the restriction of f to X_z is of general type, as claimed $\square$

Corollary 2.16. *Let $f : X \dashrightarrow Y$ be a special fibration, and let $j : Z \rightarrow X$ be such that $f \circ j : Z \dashrightarrow Y$ is onto. Assume that Z is special. Then X is special.*

Proof: Let $h : X \dashrightarrow W$ be a fibration of general type (if any). By 2.11 with $V = X$, there is a factorisation $k : Y \rightarrow W$ such that $h = k \circ f$. Thus: $j \circ h : Z \rightarrow W$ is onto. We can thus apply 2.7, which says that the Stein factorisation $h' : Z \rightarrow W'$ of $h \circ j : Z \rightarrow W$ is of general type. But Z being special by assumption, this is a contradiction and X is special.

Remark 2.17. *It is not true in general that X is special if it admits a special fibration $f : X \rightarrow Y$ with Y special (see 1.1, for example). But in some cases (if the fibres are for example, rationally connected), this is true (see 3.28).*

Example 2.18. *For any $d > 0$ and $k \in \{-\infty, 0, \dots, d-1\}$, there exists projective manifolds of dimension d and Kodaira dimension k . In particular, it is not true that a special manifold has nonpositive Kodaira dimension.*

To get examples with $k \geq 0$, just take indeed a general member of the linear system:

$|\mathcal{O}_P(d-k+2, m)|$ on $P := \mathbb{P}^{d-k+1} \times \mathbb{P}^k$ for large m , and such that this member has a section over the base $\mathbb{P}^k$.

If $k = -\infty$, just take $\mathbb{P}^d$.

We now come to a very important consequence of 2.12

Corollary 2.19. *Let $X \in \mathcal{C}$. There is **at most** one fibration defined on X which is both special and of general type. If it exists, this fibration special and is dominated by any other special*

fibration defined on X . It is also, moreover, of general type and dominates any other general type fibration defined on X .

The proof is immediate, from 2.12. In other words, such a fibration is the “lowest special” and the “highest of general type” on X . The existence of such a fibration on any X is the main result of this paper.

These two descriptions provide us with two means of construction: by consideration of chains of special subvarieties, one geometrically constructs the “lowest special” fibration on any X . This is the way used in section 3. Dually, by making fibre products of general type fibrations, one constructs the “highest general type” fibration on X . This is the approach followed in section 5. In both cases, to show that the fibration so constructed has the missing property (special if general type, and conversely), we need the orbifold additivity result 4.4.

Recall now:

Definition 2.20. *Let $f : X \rightarrow Y$ be a surjective meromorphic map between normal compact irreducible analytic complex spaces.*

*We say that f is **almost holomorphic** if $f(J) \neq Y$, where J is the indeterminacy locus of f .*

More precisely: if $X' \subset X \times Y$ is the graph of f , and $f' : X' \rightarrow Y$ the (restriction of the) second projection, then $f(J) := f'(J')$, with J' being the set of all $x' \in X'$ such that $p^{-1}(x)$ does not reduce to x' , or equivalently: is positive dimensional. (Here $p : X' \rightarrow X$ is the first projection (which is a proper modification), and $x := p(x')$).

Theorem 2.21. *Let $f : X \rightarrow Y$ be a meromorphic fibration of general type, with $X \in \mathcal{C}$ smooth. Then f is almost holomorphic.*

Remark 2.22. *The smoothness assumption is essential, as shown by the example of the cone X over a projective manifold of general type Y . The conclusion of the preceding theorem should however hold if the singularities of X are terminal, or even canonical.*

Proof: Resolve the indeterminacies of f by a sequence of smooth blow-ups $u : X' \rightarrow X$ (by H. Hironaka’s results), with $f' := f \circ u : X' \rightarrow Y$ holomorphic. If f is not almost holomorphic, some irreducible component V of the exceptional divisor of u is mapped surjectively onto Y by f' , in such a way that the fibres V_z of the restriction u' of u to V are mapped to positive-dimensional subvarieties $f(V_z)$ of Y . This contradicts 2.14, because V has two maps $u' : V \rightarrow Z := u(D) \subset X$, and $f'' : V \rightarrow Y$, the restriction of f' to V . Now, by smoothness of X , u' is special (its generic fibre is a rational variety, so apply 2.8. Moreover, by 2.7, f'' is of general type (possibly after Stein factorisation).

From 2.14, we thus have a factorisation $\phi : Z \rightarrow Y$ with $f'' = \phi \circ u'$. But this precisely contradicts $\dim(f(V_z)) > 0$, and we get the claim $\square$

2.3. General Type Fibrations and Bogomolov Sheaves.

Definition 2.23. *Let $X \in \mathcal{C}$. A rank one coherent subsheaf F of $\Omega_{X,p}^p$, $p > 0$ is said to be a (p -dimensional) **Bogomolov sheaf** on X if $\kappa(X, F) = p$.*

The properties of these Kodaira dimensions have been discussed in section 1.4 to which we thus refer.

By the results of that section, any (equivalence class of a) general type fibration f defined on X canonically determines a Bogomolov sheaf F_f on X . We shall now see the converse direction. By the results of [Bo], any p -dimensional Bogomolov sheaf determines a meromorphic fibration $f_F : X \rightarrow Y_F$ with $\dim(Y_F) = p$, and such that $F = f_F^*(K_{Y_F})$ at the generic point of Y_F . The proof given there applies only to X projective (because of the argument of cutting by transversal hyperplane sections), but (as is well-known) can be easily modified to apply to any X compact Kähler (or in $\mathcal{C}$), as follows:

Theorem 2.24. [Bo79] *Let $F \subset \Omega_X^p, p > 0$, be a Bogomolov sheaf on $X \in \mathcal{C}$. Let $f_F : X \rightarrow Y_F$ be the fibration defined by the linear system $|L^{\otimes m}|$ for $m > 0$ sufficiently large and divisible. We can assume that f is holomorphic. Then $F = f_F^*(K_{Y_F})$ at the generic point of Y_F .*

Proof: We can, using the covering trick argument of [Bo79], reduce to the case when $m = 1$, which we now treat.

We can thus select $(p + 1)$ sections $s_i, i = 0, 1, \dots, p$ of F which are analytically independent (ie: the linear system they define is f_F up to stein factorisation, and so has p -dimensional image). Because F has rank one, there exists meromorphic functions $y_i, i = 1, \dots, p$ such that $s_i = y_i \cdot s_0$. By Hodge theory (X being Kähler, or even just in $\mathcal{C}$), the holomorphic p -forms $s_i, i = 0, \dots, p$ on X are closed. From which we get: $ds_0 = dy_i \wedge s_0 = 0, i = 1, \dots, p$.

The last equality shows by simple algebraic arguments the existence of a meromorphic function g on X such that $s_0 = g \cdot (dy_1 \wedge \dots \wedge dy_p)$. The first equality shows that $g = f^*(h)$, for some meromorphic function h on Y , and so the claim, since the argument applies to $i > 0$ as well $\square$

We can thus sum up the preceding observations as follows:

Theorem 2.25. *Notations being as above, for any $X \in \mathcal{C}$, there are inverse bijective correspondances between Bogomolov sheaves F on X and (equivalence classes of) general type fibrations f defined on X . These correspondances are defined as follows:*

1. *If f is of general type, then F_f is a Bogomolov sheaf on X .*
2. *If F is a Bogomolov sheaf on X , then f_F is a fibration of general type.*

A direct application (and motivation) is:

Theorem 2.26. *The manifold $X \in \mathcal{C}$ is special if and only if there is no Bogomolov sheaf on X .*

Proof: The Bogomolov subsheaves on X correspond bijectively to fibrations of general type with domain X $\square$

Corollary 2.27. *The manifold X is special in the following two cases:*

1. *X is rationally connected (see section 3.3 for this notion).*
2. *X is a compact Kähler manifold with trivial (or torsion) canonical bundle, or even, more generally, if $c_1(X) = 0$.*

Proof: In both cases, it is shown in [C95] that $\kappa^+(X) \leq 0$, which means (in particular) that a coherent rank one subsheaf of $\Omega_X^p, p > 0$ has Kodaira dimension negative or zero. Thus X has no Bogomolov subsheaf. It is thus special, by 2.26. The result in [C95] depends on Calabi-Yau's Theorem. But in the projective case, one can get algebro-geometric proofs using Miyaoka's generic semi-positivity Theorem $\square$

We shall see later that the weaker condition $\kappa(X) = 0$ is actually sufficient for X to be special. Notice that the property shown in [C95] in the above two cases is considerably stronger than the absence of Bogomolov sheaves. This is not surprising, in view of the fact that these manifolds are the building blocks of the class of special manifolds, but do not exhaust this class, by far.

2.4. General Type Reduction.

2.4.1. Ordering of Fibrations. Recall from 1.1.2 that a meromorphic fibration $f : X \dashrightarrow Y$ canonically defines (see [C85]) a meromorphic map $\phi_f : Y \rightarrow \mathcal{C}(X)$, with $\mathcal{C}(X)$ the Chow-Scheme of X , by sending the generic $y \in Y$ to the point of $\mathcal{C}(X)$ parametrising the reduced fibre X_y of f over y . Let $\Phi_f \subset \mathcal{C}(X)$ be the image of Y by ϕ_f : it is a compact irreducible analytic subset of $\mathcal{C}(X)$ bimeromorphic to Y such that its incidence graph is bimeromorphic to Y .

The above correspondance induces a bijective map between equivalence classes of fibrations and compact irreducible analytic subsets of $\mathcal{C}(X)$ with incidence graph bimeromorphic to X .

Recall also we said that f is **almost holomorphic** if the image by f (naturally defined using the graph of f) of the indeterminacy locus $I(f)$ of f is not all of Y (see 2.20 for details). It is easy to show that Φ_f is an irreducible component of $\mathcal{C}(X)$ if f is almost holomorphic (see [C85]).

We now introduce an order on the set of (equivalence classes of) fibrations with domain X .

We say that f **dominates** the fibration $g : X \rightarrow Z$ if there exists a meromorphic fibration $\phi : Y \dashrightarrow Z$ such that $g = \phi \circ f$. Equivalently: each fibre of f is contained in some fibre of g . This binary relation defines an ordering on the set $\mathcal{F}(X)$ of all equivalence classes of fibrations (seen as a subset of $\mathcal{C}(\mathcal{C}(X))$).

There is now an easy

Lemma 2.28. *If $\Lambda \subset \mathcal{F}(X)$ is any subset, it has in the ordered set $\mathcal{F}(X)$ a least upper bound, denoted Λ^+ .*

Proof: Λ^+ is so constructed: let $\Lambda_0 := \{\lambda_1, \dots, \lambda_N\} \subset \Lambda$ be finite such that the product map $f := f_{\lambda_1} \times \dots \times f_{\lambda_N}$ has an image of maximal dimension. Then take for Λ^+ the (fibration part of the) Stein factorisation of f . $\square$

Finally, the preceding construction shows that if any element of Λ is almost holomorphic, so is the least upper bound Λ^+ of the family.

Example 2.29. *If Λ consists of fibrations onto varieties of general type, then Λ^+ is also a fibration onto a variety of general type.*

This is easily reduced to the case when Λ has two elements, and then reduces to showing that if $Z \subset Y \times Y'$ is a subvariety of a product of two varieties of general type, then Z itself is of general type if it is mapped surjectively to Y, Y' by the first and second projections. This results easily from the additivity theorem for fibrations with base of general type (generalised orbifold versions will be proved in section 4 below) $\square$

Definition 2.30. *For any $X \in \mathcal{C}$, let $gt_X : X \dashrightarrow GT(X)$ be the least upper bound in $\mathcal{F}(X)$ of the family Λ_X of all (equivalence classes of) fibrations of general type $f_j : X \dashrightarrow Y_j$.*

(if X is special, we just take for gt_X the constant fibration).

*We call gt_X the **general type reduction** of X .*

From 2.4.1 above, we deduce:

Proposition 2.31. *For any $X \in \mathcal{C}$, smooth, the map $gt_X : X \rightarrow GT(X)$ is almost holomorphic.*

Question 2.32. *Let $X \in \mathcal{C}$. Is gt_X either constant, or a fibration of general type?*

This question has a positive answer if so does the next one:

Question 2.33. *Let $u : X \rightarrow U$ and $v : X \rightarrow V$ be fibrations of general type. Is then $f : X \rightarrow W$ of general type if so are u and v ? (Here f is the Stein factorisation of the product map $(u \times v) : X \rightarrow W'$, with $W' := (u \times v)(X) \subset U \times V$).*

It is obvious that if 2.33 has a positive answer, so does 2.32, by the construction of gt_X . We shall see in 5.17 that these two questions have an affirmative answer. But this answer rests on the very different techniques of section 4.

Theorem 2.34. *Let $X \in \mathcal{C}$, and $f : X \dashrightarrow Y$ be any fibration. Then f admits a **relative gt-reduction**.*

This means that there exists a unique factorisation $f = h \circ g$ of f by fibrations $h : Z \dashrightarrow Y$ and $g : X \dashrightarrow Z$ such that for y general in Y , the restriction $g_y : X_y \dashrightarrow Z_y$ of g to X_y is the gt -reduction of X_y .

Proof: This construction is actually in essence already in [C80] to which we refer for more details. We can and shall assume that X is smooth and f holomorphic, due to the bimeromorphic invariance of the notions involved.

We shall actually show a more precise version:

Lemma 2.35. *Let $f : X \rightarrow Y$ be a fibration, with $X \in \mathcal{C}$.*

After a generically base change $v : Y' \rightarrow Y$ and proper modifications that we notationally ignore, there exists finitely many factorisations $f = h_i \circ g_i$, $i = 1, 2, \dots, N$, with $g_i : X \rightarrow Z_i$, $h_i : Z_i \rightarrow Y$, such that:

1. *The restriction $g_{i,y} : X_y \rightarrow (Z_i)_y$ of each g_i to the general fibre X_y of f is of general type.*
2. *if $g : X \rightarrow Z$ is the (fibration part of) the Stein factorisation of the product map $g_1 \times \dots \times g_N : X \rightarrow Z_1 \times_Y Z_2 \times \dots \times Z_{N-1} \times_Y Z_N$, then:
the restriction $g|_{X_y} : X_y \rightarrow Z_y$ of g to the general fibre X_y of f coincides with the gt -reduction $gt_{X_y} : X_y \rightarrow G(X_y)$ of that fibre.*

Proof: Let, for $y \in Y^*$, $gt_y : X_y \rightarrow Z_y$ be the gt -reduction of X_y , where Y^* is the Zariski open subset of Z over which f is smooth. For such a y , let Z'_y be the family of fibres of gt_y , defined as the image of the meromorphic map from Z_y to $\mathcal{C}(X_y)$ sending a generic point of Z_y to the point in $\mathcal{C}(X_y)$ parametrising its reduced gt_y -fibre in X_y .

Because gt_y is an almost holomorphic map by 2.21, Z'_y is an irreducible component of $\mathcal{C}(X_y)$.

Consider now the Zariski closed subset $\mathcal{C}(X/Y)$ of $\mathcal{C}(X)$ consisting of cycles contained in some fibre of f . It is naturally equipped with the holomorphic map $f_y : \mathcal{C}(X/Y) \rightarrow Y$ sending such a cycle to the fibre containing it. (Strictly speaking, one may need to weakly normalise first, to make f_* holomorphic, but this does not change the argument).

Because $X \in \mathcal{C}$, the irreducible components of $\mathcal{C}(X/Y)$ are compact.

Assume the fibre of f is not special, for y in a subset of Y which is of second category, in Baire's terminology. (Being of second category means: not contained in a countable union of closed subsets with empty interior. As we shall see later, the right topology here in our context is the Zariski topology, not the metric topology).

Because $X \in \mathcal{C}$, the irreducible components of $\mathcal{C}(X/Y)$ are compact. By the countability at infinity of $\mathcal{C}(X/Y)$, there is an irreducible component Γ' of $\mathcal{C}(X/Y)$ mapped surjectively onto Y by f_* , and such that the f_* -fibre Γ'_y of Γ' over y has a component equal to $Z'_y \subset \mathcal{C}(X_y)$, the family of fibres of a fibration of general type $g_{i,y} : X_y \dashrightarrow Z'_y$. This map is almost holomorphic, by 2.21. The Stein factorisation of f_* restricted to Γ' gives a finite base change for Y . This base changes we shall notationally ignore, here, because they are irrelevant to our problem. So we deal as if the generic fibres of $f_{*|\Gamma'}$ were irreducible.

Thus, for some $y \in Y^*$, the fibre Γ'_y of $f_{*|\Gamma'}$ is the family of fibres of some almost holomorphic meromorphic fibration $g_{i,y} = \gamma_y : X_y \rightarrow Z_{y,\gamma}$ on X_y . By the (obvious) openness of almost holomorphicity, one deduces the existence of such a γ_y for the generic $y \in Y$. And so, using the graph of the family, we get a factorisation $f = \delta \circ \gamma$, with fibrations $\gamma : X \rightarrow Z_\gamma$, and $\delta : Z_\gamma \rightarrow Y$.

By our assumption, γ_y is a fibration of general type for y in $S \subset Y$ of second Baire category in Y . From 1.25, we conclude that γ_y is still of general type for y general in Y .

The construction of the g_i 's is now obvious, by observing that if the map g resulting from a finite family of g_i 's, $i = 1, 2, \dots, N$, does not induce gt_{X_y} on the general X_y , there exists, by the same argument as above, a component Γ' , inducing a general type fibration on the general X_y , and such that its (Stein factorised) fibre product over Y with the preceding ones will increase the dimension of the resulting Z . Contradiction.

This shows the lemma, and so 2.34 $\square$

Definition 2.36. *A subset $A \subset V$ of a complex analytic space is said to be of **second Zariski category in V** if it is not contained in a countable union of Zariski closed subsets with empty interior of V . (Notice that the definition makes sense in the algebraico-geometric context as well).*

From the proof 2.34, we immediately get:

Corollary 2.37. *Let $f : X \dashrightarrow Y$ be a fibration, with $X \in \mathcal{C}$.*

Assume that $\dim(GT(X_y)) = d$, for $y \in A$, where A is of second Zariski category in Y . Then, this equality holds for the general point y of Y .

In particular, if X_y is special for y in a subset of second Zariski category in Y , the general fibre of f is special.

Proof: Let $f = h \circ g$ be the gt -reduction of f . By assumption, $\dim(GT(X_y)) = d$ for $y \in A$. But also $\dim(g(X_y)) = \dim(GT(X_y))$ for y general in Y , and $\dim(g(X_y)) = d$ for y generic in Y . Thus $\dim(GT(X_y)) = d$ for $y \in Y$ general $\square$

2.5. The Algebraic Reduction. As an application of the preceding arguments, we show Theorem 2.3:

Theorem 2.38. *Let $a_X : X \dashrightarrow \text{Alg}(X)$ be the algebraic reduction of $X \in \mathcal{C}$. The generic fibre of a_X is special.*

Proof: Assume not. By lemma 2.35 above, after a suitable finite base change over $\text{Alg}(X)$ (which we notationally ignore because it preserves the algebraic reduction and dimension), there exists a non-trivial factorisation $a_X = h \circ g$ with g a fibration inducing a fibration of general type over the general fibre of a_X . Write it: $a_X = h \circ g$, with $g : X \rightarrow Z$ and $h : Z \rightarrow \text{Alg}(X)$. Then: $\dim(Z) > \dim(\text{Alg}(X))$.

By construction, the line bundle $(K_Y + \Delta(g))$ over Z is thus h -big. Thus Z is Moishezon, as one sees considering the line bundle $L := (h^*(k.H) + (K_Z + \Delta(g)))$ on Z , which is big for k a large and positive integer, and H an ample line bundle on $\text{Alg}(X)$, which we obviously can choose to be projective. (See for example, the Proof of [U], Theorem (12.1)). But this contradicts the definition of a_X , and proves the claim $\square$

2.6. The Category of Orbifolds. We very briefly discuss without proofs the extension of part of our considerations to orbifolds, restricting here to prepared orbifolds (Y/Δ) with Y smooth and the support of Δ an s.n.c divisor of Y (but ultimately, one needs to consider *klt* orbifolds). One of the main point is to define bimeromorphic equivalence. The right notion seems to be derived from **terminal modifications**.

Definition 2.39. *The bimeromorphic holomorphic map $v : Y' \rightarrow Y$ is said to induce a bimeromorphic map: $v : (Y'/\Delta') \rightarrow (Y/\Delta)$ if it is **terminal** with respect to the orbifold structures, that is if: $K_{Y'} + \Delta' = v^*(K_Y + \Delta) + \sum_{j \in J} a_j.E_j$, where (as usual) the a_j are all positive, and J is the collection of v -exceptional divisors on Y' . (One might also define similarly the notion of **canonical modification**, of course).*

Notice that the orbifold Kodaira dimension is invariant under bimeromorphic equivalence of orbifold, which is the one generated by terminal modifications.

One can define for any fibration $g : (Y/\Delta) \rightarrow Z$ its orbifold base, as in 1.6. One can extend this notion to the case of meromorphic g , by first resolving the indeterminacies of g by a terminal modification.

The Kodaira dimension of this fibration is then the minimum of the Kodaira dimensions of the orbifolds bases of fibrations equivalent to g , these being defined on orbifolds (Y'/Δ') bimeromorphically equivalent to (Y/Δ) .

Definition 2.40. *The fibration $g : (Y/\Delta) \rightarrow Z$ is of **general type** if its Kodaira dimension is $\dim(Z) > 0$.*

*The orbifold (Y/Δ) is **special** if it has no fibration of general type.*

Fundamental tools for the study of orbifolds are the locally free sheaves $\Omega_Y^p(\log(\Delta))$ of logarithmic forms along Δ (classically known when the multiplicities are infinite, or said differently, when Δ is reduced). We shall not give the definition here, but simply say that sections of this sheaf can be symbolically written locally in the standard normal crossing coordinates for fixed q as linear combinations of expressions of the form:

$h.(dy_{j_1}/(y_{j_1}^{(1-1/m_{j_1})}) \wedge \dots \wedge (dy_{j_s}/y_{j_s}^{(1-1/m_{j_s})}) \wedge dy_{j_{s+1}} \wedge \dots \wedge dy_{j_q}$, with h holomorphic, and:

$1 \leq j_1 < \dots < j_s \leq r < j_{s+1} < \dots < j_q$ if $y_1 \dots y_r$ is a local equation of $|\Delta|$, the multiplicities being given by the m_i 's.

A section of $\Omega_Y^q(\log(\Delta))$ is thus an m -th root of a well-defined holomorphic tensor, when m is an integer divisible by each of the m_i 's.

More precisely, a section s of this sheaf is defined as a pair (F, s) , where F is a rank one coherent subsheaf of $\Omega_Y^q(\log |\Delta|)$, and s is a holomorphic section of $F^{\otimes m}$, for some m divisible by all the m_i 's, and such that $s \in (\Omega_Y^q(\log |\Delta|))^{\otimes m}(-m.\Delta^*)$, where Δ^* is the $\mathbb{Q}$ -divisor on Y defined by: $\Delta^* := |\Delta| - \Delta = \sum_{i \in I} (1/m_i) \cdot \Delta_i$.

By lifting to a Δ -nice covering (see section 4), these sections become standard p -forms. From which one deduces the important property of d -closedness of such $\log((\Delta))$ -forms.

As we did above, one can then also define directly the Kodaira dimension of a fibration by introducing the saturation of the differential sheaf defined by g in $\Omega_Y(\log(\Delta))$. Because we may only consider high and divisible multiples to define the Kodaira dimensions, one does not need to define precisely $\Omega_Y(\log(\Delta))$ to define this orbifold Kodaira dimension, and directly look at rank one subsheaves F of $\Omega_Y(\log |\Delta|)$, and define as usual the Kodaira dimension of $(F^{\otimes m}(-m\Delta^*))$.

Their relevance to our topic is that special orbifolds (Y/Δ) are characterised by the absence of Bogomolov sheaves on (Y/Δ) , defined as in 2.23 above when Δ is empty, just replacing there Ω_Y^p by $\Omega_Y^p(\log \Delta)$.

The correspondance between Bogomolov sheaves on (Y/Δ) and general type fibrations on this orbifold extends to this orbifold context. The orbifold additivity theorem then applies in this context.

The construction of the core for an orbifold can then be made by the second approach we followed (as the highest special fibration). The geometric approach seems more delicate than for varieties, and certainly needs some extra arguments, because one needs to take into account the order of contact of the subvarieties with the orbifold divisor.

3. The Core

3.1. The Core: Functoriality Properties. We use the notations of section 12.

Definition 3.1. *Let $X \in \mathcal{C}$ be normal. Let $A := A(X) \subset \mathcal{C}(X)$ be the family of special subvarieties of X . It is Z -regular. Let then $T(A)$ be the family of its components (see proposition 12.22), and let $c_X : X \dashrightarrow C(X)$ be the $T(A)$ quotient of X . This almost holomorphic fibration will be called **the core of X** . (see section 12)*

In general, not much can be said about the fibres of c_X (except that two of their points are joined by a chain of special subvarieties of X , at least if question 12.25 has a positive answer).

Example 3.2. *Let X be the cone over a variety of general type V . Then c_X is the constant map. But X is by no means special, since it has a $\mathbb{P}^1$ -fibration over V . Note that this fibration is not almost holomorphic.*

This example shows the role of singularities. Indeed:

Theorem 3.3. *Let $X \in \mathcal{C}$ be **smooth**. Let $c_X : X \rightarrow C(X)$ be the core of X . Then:*

1. *The general fibre of c_X is special.*
2. *If F is a general fibre of c_X , and if $Z \subset X$ is a special subvariety of X meeting F , then Z is contained in F (such a fibre F will be said c_X -general).*
3. *c_X is almost holomorphic.*

Remark 3.4. *The above result should hold true when X is singular, provided it has at most canonical singularities. Notice that c_X is not a bimeromorphic invariant, in general (as the preceding example shows). But it is easily seen from 3.3 to be so if X is smooth: then $c_Y = c_X \circ m$, if $m : Y \rightarrow X$ is bimeromorphic.*

When X is smooth, we denote by $\text{ess}(X)$ the dimension of $C(X)$, and call it the **essential dimension** of X . Thus $\text{ess}(X) = 0$ iff X is special, and $\text{ess}(X) = d = \dim(X)$ iff X is of general type, by Theorem 5.5 below. In the first case, the core is the constant map; in the second case, it is the identity map.

Theorem 3.5. *Let $X \in \mathcal{C}$ be normal. Let $c_X : X \dashrightarrow C(X)$ be its core, and let $a_X : X \dashrightarrow A(X)$ be its algebraic reduction. There exists a factorisation $b_X : A(X) \dashrightarrow C(X)$ of c_X such that $c_X = b_X \circ a_X$.*

In particular: $C(X)$ is always Moishezon.

Proof: By Theorem 2.3, the fibres of a_X (which is a bimeromorphic invariant of X) are special, hence contained in the fibres of c_X $\square$

We shall prove Theorem 3.3 below. Before we start with some easy observations.

Proposition 3.6. *Let $c_X : X \rightarrow C(X)$ be the core of X , smooth. Let $h : Z \dashrightarrow X$ be any meromorphic map. Assume $h(Z)$ meets some c_X -general fibre of c_X . There exists then a natural meromorphic map $c_h : C(Z) \dashrightarrow C(X)$ such that $c_h \circ c_Z = c_X \circ h$.*

Proof. : By the assumption, if $z \in Z$ is general, its image $x := h(z)$ in X belongs to a c_X -general fibre F_z of c_X . The fibre of c_Z through z is special. Thus so is its image V_z by h . Since V_z meets F_z , it is contained in F_z , by property [2.] in Theorem 3.3. Hence the existence of c_h $\square$

Corollary 3.7. *Let $h : Z \dashrightarrow X$ be as in proposition 3.6 above. Then c_h as above exists in the following cases:*

1. $c_X \circ h : Z \rightarrow C(X)$ is surjective. (In particular if h itself is surjective).
2. $Z \subset X$ is the general member of a family $(Z_t)_{t \in T}$ of subvarieties of X such that the varieties $c_X(Z_t)$ cover $C(X)$.
3. $Z = X_y$ is a general fibre of $(\psi \circ c_X)$, where $\psi : C(X) \rightarrow Y$ is any fibration. In this case, c_Z is simply the restriction of c_X to Z .

Let us give some easy examples in which c_X can be described:

Proposition 3.8. *Let $X \in \mathcal{C}$ be a manifold, and assume that $f : X \dashrightarrow Y$ is a special fibration of general type. Then: $f = c_X$*

In particular: there is at most one fibration both special and of general type on X .

Proof: Because f is special, there is a factorisation $g : Y \dashrightarrow C(X)$ such that $g \circ f = c_X$. Indeed: the general fibre F of f is special and meets the general fibre C of c_X . Thus $F \subset C$.

But f is of general type, and so by Theorem 2.11, there exists a factorisation $h : C(X) \dashrightarrow Y$ such that $f = h \circ c_X$. Thus $f = c_X$, as claimed. $\square$

Remark 3.9. *We shall later conjecture (and prove in some cases (see section 5.8)) that c_X is a fibration of general type. So that c_X should always be characterised as the unique fibration of domain X both special and of general type.*

Corollary 3.10. *Let X be a manifold of general type. Then: $c_X = id_X$ (the identity map of X), and so: $ess(X) = dim(X)$.*

Proof: Indeed: id_X is then special and of general type. Apply then 3.8 $\square$

Remark 3.11. *We shall see later in Theorem 5.5 that the converse also holds true: if $ess(X) = dim(X) > 0$, then X is of general type.*

Corollary 3.12. *Let $X \in \mathcal{C}$ be a manifold, with $n := dim(X)$. Then $ess(X) = (n - 1)$ in the following two cases:*

- a. $\kappa(X) = (n - 1)$, and J_X is a fibration of general type. Then: $c_X = J_X$.
- b. $R(X)$ (the rational quotient of X) is of dimension $(n - 1)$ and of general type.

Proof: Indeed, in case (a.) (resp. (b.)), the generic fibre of J_X (resp. r_X) is an elliptic (resp. rational) curve. The fibration J_X (resp. r_X) is thus special. The other conditions imply that it is also of general type. It is thus the core of X . In particular, $ess(X) = (n - 1)$ $\square$

Remark 3.13. *We shall see later in Theorem 5.7, as a consequence of orbifold additivity theorems, that the converse also holds true: if $ess(X) = dim(X) - 1 > 0$, then X is of the type (a) or (b). For the case $ess(X) = (n - 2)$ see below 3.35.*

Finally, in the same section, we shall also give a conjectural description of the core as a canonical and functorial composition of orbifold rational quotients and Iitaka fibrations.

Corollary 3.14. *Let $f : X \dashrightarrow C$ be a special fibration, with $X \in \mathcal{C}$ and C a curve. Then: either f is of general type, and $f = c_X$, or f is not of general type, and X is special.*

Proof: In the first case, the claim follows from Proposition 3.8. In the second, from the fact that if $g : X \dashrightarrow Z$ is a fibration of general type, there exists by 2.11 a factorisation $h : C \dashrightarrow Z$ such that $g = h \circ f$. But C is curve, and f is not of general type. Thus Z is a point, and g is not of general type. Contradiction. $\square$

The core can be constructed in a relative setting, as well, by a simple application of Theorem 12.27:

Theorem 3.15. *Let $f : X \rightarrow Y$, with $X \in \mathcal{C}$. There exists a unique factorisation $f = g_f \circ c_f$ by two fibrations $c_f : X \rightarrow C(f)$ and $g_f : C(f) \rightarrow Y$ such that, for $y \in Y$ general, the restriction $c_f : X_y \rightarrow C(f)_y$ is the core of X_y .*

*We call the factorisation $f = g_f \circ c_f$ the **core of f** .*

We now prove Theorem 3.3:

3.2. Construction of the Core as the Lowest Special Fibration. Proof of Theorem 3.3:

For this, we shall simply apply theorem 12.31 to the family $A(X)$ of special subvarieties of X , supposed to be smooth and to be in $\mathcal{C}$.

We know that $A(X)$ is $\mathbb{Z}$ -regular (see section 12.4 for this notion). It is thus sufficient to show that $A(X)$ is also stable (see 12.5 for this notion). The property [stab2] is obtained by applying corollary 2.5.

The property [stab1] is the content of the next Theorem, which thus establishes at the same time Theorem 3.3:

Theorem 3.16. *Let $T \subset T(A(X)) \subset \mathcal{C}(X)$ be a special family as above, with X smooth in $\mathcal{C}$. Assume each irreducible component of T is X -covering. Then:*

1. *X is special if X is T -connected (ie: q_T is the constant map).*
2. *The general fibre of q_T is special.*

Proof: The second assertion is a consequence of the first, because the if X_y is a general fibre of q_T , then X_y is smooth since q_T is almost holomorphic, and the family T_y consisting of $t \in T$ such that $V_t \subset X_y$ is a finite union of covering families of X_y with general member special, and such that X_y is T_y -connected.

We thus only need to show the first statement: assume there existed a meromorphic fibration $f : X \dashrightarrow Y$ of general type. By 2.21, f is almost holomorphic, since X is supposed to be smooth. By Theorem 2.11 applied to each irreducible component V_i of V , we have a meromorphic factorisation $\phi : T \dashrightarrow Y$ such that $\phi \circ g = f \circ \psi : V \dashrightarrow Y$.

Assume first that f is holomorphic. Thus f is constant on every V_t , and so on every T -chain. Because X is T -connected, f takes the same value on two arbitrary points of X . Thus f is constant and Y is a single point, in contradiction with the fact that it is of general type. Thus X is special, as claimed.

If f is only almost holomorphic, the same argument applies, provided we choose an f -regular point $y \in Y$: for every $t \in T$, if V_t meets X_y , then V_t is contained in X_y (because of the factorisation property $\phi \circ g = f \circ \psi$, and the usual rigidity lemma. More precisely: approximate V_t in $\mathcal{C}(X)$ by a sequence V_{t_n} , such that $V_{t_n} \subset X_{y_n} \subset f^{-1}(U)$, for some Stein-or affine in the algebraic category-neighborhood U of y in Y , and n large. This is possible because the generic $V_{t'}$ is contained in a fibre of f . Then $V_t \subset X_y$, by an easy argument, based on the fact that f is holomorphic on $f^{-1}(U)$).

So we get that the generic member, hence every member of the family T is contained in some fibre of f (here the notion of fibre of f is the usual Chow-scheme theoretic one, defined in section 1.1.2).

Thus every T -chain meeting X_y is contained in X_y . Because X is T -connected, X is contained in X_y and f is constant, and so not of general type, as assumed. Contradiction. $\square$

We can now establish the following often useful characterisations of the core:

Theorem 3.17. *Let $f : X \dashrightarrow Y$ be a special fibration, with $X \in \mathcal{C}$, such that for any special fibration $g : X \dashrightarrow Z$, there exists a factorisation $\phi : Z \dashrightarrow Y$ of $f := \phi \circ g$. Then $f = c_X$ is the core of X .*

In particular, if $f : X \dashrightarrow Y$ is a special fibration of general type, it is the core of X .

The relevant diagram is:

$$\begin{array}{ccc} X & \xrightarrow{\quad f \quad} & Y \\ g \downarrow & \nearrow \phi & \\ Z & & \end{array}$$

Proof: First, because c_X is special, there exists a factorisation $\psi : \mathcal{C}(X) \dashrightarrow Y$. Let F be a c_X -general fibre of c_X . By the existence of ψ , it is contained in some fibre G of f . But G is special, because f is. By the defining property of c_X , we have the reverse inclusion $G \subset F$.

The last assertion follows now from theorem 2.11, because f being of general type, for any special fibration g , theorem 2.11 shows that the factorisation ϕ exists. $\square$

3.3. Rationally connected manifolds. We now come to our first basic example of special manifolds: the rationally connected ones.

We recall first their definition and some properties.

Recall ([C3],[Ko-Mi-Mo]) that an irreducible compact complex space X is said to be **rationally connected** if any two generic points of X are contained in a rational chain of X (ie: a connected projective curve of X , all irreducible components of which are rational (possibly singular) curves). Examples of rationally connected manifolds include unirational and Fano Manifolds, twistor spaces. This property is bimeromorphically stable among manifolds, but not among varieties (the cone over a projective manifold which is not rationally connected will again provide such an example).

(Of course, the above definition can be given for algebraic varieties defined over arbitrary fields).

We refer to [Ko-Mi-Mo] for some of the fundamental properties of this class of manifolds.

Rational connectedness has a slightly different characterisation, by the following fundamental result ([G-H-S]):

Theorem 3.18. ([G-H-S]): *Any fibration $f : X \rightarrow C$ over a projective curve C with X smooth projective and generic fibre rationally connected has a holomorphic section.*

Definition 3.19. ([C95]) *Let $X \in \mathcal{C}$ be irreducible. We say that X is **rationally generated** if for any surjective meromorphic map $f : X \dashrightarrow Y$, Y is uniruled.*

Any rationally connected $X \in \mathcal{C}$ is thus rationally generated. But, conversely:

Theorem 3.20. *Let $X \in \mathcal{C}$ be rationally generated. Then X is rationally connected.*

Proof: By induction on the dimension: X is uniruled (take $f = id_X$). Let $r_X : X \rightarrow R(X)$ be the rational quotient of X . Then $R(X)$ is also rationally generated (obvious). By induction, it is rationally connected. So it is in particular Moishezon (by [C81], which says in particular that if any two points of $X \in \mathcal{C}$ can be joined by a chain of curves, then X is Moishezon). Thus X too is Moishezon (by [C85], which says among others that $X \in \mathcal{C}$ is Moishezon if there is a fibration $u : X \dashrightarrow Y$ with Y and the generic fibre F of u Moishezon, if F has $q(F) = 0$), because the fibres of r_X are Moishezon, with vanishing irregularity. We can assume X to be projective, by the bimeromorphic invariance of the rational generatedness. But then the conclusion follows easily from [G-H-S], which allows to lift rational curves from $R(X)$ to X $\square$

Theorem 3.21. *Let $X \in \mathcal{C}$ be a rationally connected manifold. Then X is special.*

Notice that the smoothness of X is essential, as shown again by the cone over a projective manifold of general type.

Proof: Let $c_X : X \dashrightarrow C(X)$ be the core. Assume it is not the constant map. Let F be a c_X -general fibre of c_X . Because X is rationally connected, some rational curve in X meets F , but is not contained in F . Contradiction. $\square$

We now extend these results to the relative case.

3.4. The Rational Quotient and The Core. We now turn to the study of the **rational quotient** of X from the point of view of special varieties. The rational quotient of $X \in \mathcal{C}$ was introduced in [C3] as an application of T -quotients. It was also independently constructed in [Ko-Mi-Mo] (under the name of “maximal rationally connected fibration” (M.R.C for short)) by a different method based on their “glueing lemma” for rational curves, in the algebraic context.)

Theorem 3.22. *Let $X \in \mathcal{C}$ be normal. There exists a unique meromorphic fibration $r_X : X \dashrightarrow R(X)$, called the **rational quotient** of X such that:*

1. *The general fibre of r_X is rationally connected.*
2. *The general fibre of r_X contains any rational curve of X that it meets.*

As usual, r_X is almost holomorphic.

The proof is given in section 12; see theorem 12.26. Notice that, by Theorem 12.28, the rational quotient also exists in relative version.

Corollary 3.23. *Let $X \in \mathcal{C}$ be smooth. The rational quotient r_X of X is then a special fibration. There exists a factorisation: $(cr)_X : R(X) \dashrightarrow C(X)$ of $c_X = (cr)_X \circ r_X$.*

Proof: This is simply because X , and so the generic fibre of r_X is special, by Theorem 3.21. This shows the first assertion. Notice the second is obvious, and does not require X to be smooth. $\square$ We have the following easy property (already shown for c_X). The proof for r_X is similar:

Proposition 3.24. *Let $f : X \rightarrow Y$ be a surjective meromorphic map, with $X \in \mathcal{C}$. Then f induces functorial maps (always denoted f_*): $f_* : R(X) \rightarrow R(Y)$ and $f_* : C(X) \rightarrow C(Y)$.*

In the above proposition, taking $f := r_X$, we get in particular a natural map $(r_X)_* : C(X) \dashrightarrow C(R(X))$. For the rational quotient, we have a particular property not valid for arbitrary special fibrations:

Theorem 3.25. *Let X be smooth and Moishezon. Let $r_X : X \rightarrow R(X)$ be the rational quotient of X . Let $c_{R(X)} : R(X) \rightarrow C(R(X))$ be the core of $R(X)$. Then: $(r_X)_* : C(X) \dashrightarrow C(R(X))$ is bimeromorphic. Equivalently: $(c_{R(X)} \circ r_X) : X \rightarrow C(R(X))$ is the core of X .*

In other words: $C(R(X)) = C(X)$.

Remark 3.26. *The hypothesis that X is Moishezon can certainly be weakened to: $X \in \mathcal{C}$. For this, it is sufficient to make the same weakening in the hypothesis for G in 3.28 below.*

Proof: We have natural fibrations $\phi : R(X) \dashrightarrow C(X)$ and $\psi = (r_X)_* : C(X) \dashrightarrow C(R(X))$ defined above. We thus only need to show that ψ is isomorphic, or that the general fibre F of $(c_{R(X)} \circ r_X) : X \dashrightarrow C(R(X))$ is special.

Observe that we have, by restricting r_X to F a map $r_X : F \dashrightarrow G$, where $G = (r_X)(F)$ is the corresponding fibre of $c_{R(X)}$. Thus F is fibered over G , which is special, with fibres which are generically rationally connected. The claim thus follows from the next:

Proposition 3.27. *Let $r : F \rightarrow G$ be a fibration with $F \in \mathcal{C}$ such that G is Moishezon and special, and the generic fibre of f is rationally connected. Then F is special.*

Proof: Let, if any, $g : F \rightarrow H$ be an admissible holomorphic fibration of general type. Since f has special fibres, there exists by Theorem 2.11 a factorisation $\phi : G \rightarrow H$ of $g = \phi \circ f$. But now by Lemmas 3.28 and 3.29 below, we see that $\Delta(g) = \Delta(\phi)$. Thus ϕ is of general type, too. But this contradicts G being special. Such a g thus does not exist, and X is special.

Lemma 3.28. *Let $f : F \rightarrow G$ be a fibration with generic fibres rationally connected and G Moishezon. Then f is multiplicity free (ie: $\Delta(f)$ is empty).*

Proof: We may assume that G is projective. The claim then follows immediately from [G-H-S] (and is actually the most difficult part of the proof), by considering the restriction of f over a very ample curve of G meeting transversally any irreducible component of $\Delta(f)$ $\square$

The proof of the following is immediate from the definition of multiplicities and the computation of the base orbifold divisor of a composed fibration, as in section 1.6:

Lemma 3.29. *Let $f : F \rightarrow G$ and $\phi : G \rightarrow H$ be fibrations. Assume that f is multiplicity free. Then $\Delta(\phi \circ f) = \Delta(\phi)$.*

3.5. Surfaces. We can describe the core of a surface as follows, in terms of its rational quotient or Iitaka-Moishezon fibration. This will be extended to threefolds in the next section. A conjectural description in arbitrary dimension will be given in section 6.4.

Recall that for any compact connected complex manifold X with $\kappa(X) \geq 0$, we denote by $J_X : X \dashrightarrow J(X)$ its Iitaka-Moishezon fibration. Let also $\kappa'(X) := \kappa(J(X), J_X)$. Obviously:
 $\kappa(X) \geq \kappa'(X) \geq -\infty$.

Theorem 3.30. *Let X be a compact Kähler smooth surface. Then its core c_X is described as follows:*

1. $\kappa(X) = 2$: then $c_X = id_X$, and $ess(X) = 2$
2. $\kappa(X) = \kappa'(X) = 1$: then $c_X = J_X$ and $ess(X) = 1$.
3. $\kappa(X) = 1 > \kappa'(X)$: X is special.
4. $\kappa(X) = 0$: X is special.
5. $\kappa(X) = -\infty$ and $q(X) \geq 2$: then $c_X = r_X$, $ess(X) = 1$.
6. $\kappa(X) = -\infty$ and $q(X) \leq 1$: X is special.

Proof: If $\kappa(X) = 2$, the claim is given by 3.10. If $\kappa(X) = 1$, the fibration $J_X : X \dashrightarrow C = J(X)$ is special, and the claim follows from 3.14. If $\kappa(X) = 0$, X is special from the facts just recalled above. If $\kappa(X) = -\infty$, from the classification of surfaces, X is bimeromorphic to $\mathbb{P}^1 \times C$, C a curve with $g(C) = q(X)$, and r_X is the projection to C if $q > 0$, and the constant map if $q = 0$. The claims are then obvious $\square$

Corollary 3.31. *Let X be a Kähler surface. Either X is special (and c_X is the constant map); or $\kappa(X) \geq 1$ and $c_X = J_X$ (the Iitaka fibration); or $\kappa(X) = -\infty$, and $c_X = r_X$ (the rational quotient of X).*

One can compute $ess(X)$ as follows:

1. $ess(X) = 2$ iff $\kappa(X) = 2$;
2. $ess(X) = 1$ iff $\kappa(X) \in \{1, -\infty\}$ and $\pi_1(X)$ is not virtually abelian;

3. $\text{ess}(X) = 0$ iff $\kappa(X) \leq 1$ and $\pi_1(X)$ is virtually abelian.

(Recall that a group G is said to be almost (or virtually) abelian if it has a finite index subgroup which is abelian)

Proof: All claims are deduced immediately from 3.30, except for the ones concerning the fundamental group, when $\kappa(X) \leq 1$.

If $\kappa(X) = -\infty$, then $\pi_1(X) \cong \pi_1(C)$, with the notations of the proof of 3.30. The assertion is obvious. If $\kappa(X) = 0$, we know that $\pi_1(X)$ is almost abelian from classification theory. If $\kappa(X) = 1$, the assertion follows from Lemma 3.33, applied to J_X $\square$

Corollary 3.32. *A compact Kähler surface X is special if and only if it has a finite étale cover which is bimeromorphic to one of the following surfaces:*

1. $\mathbb{P}_2(\mathbb{C})$
2. $\mathbb{P}_1(\mathbb{C}) \times E$, with E elliptic.
3. $K3$, or Abelian.
4. Elliptic over a curve C with m multiple fibres, C either rational (then $m \leq 2$) or elliptic (then $m = 0$).

Proof: The surfaces listed above are special, by 3.30 above. Thus so are their undercovers. Conversely, if X is special, it has a finite étale cover in the preceding list. This is clear if $\kappa(X) \leq 0$, by classification, and from the next Lemma 3.33 if $\kappa(X) = 1$ $\square$

Lemma 3.33. *Let $f : X \rightarrow C$ be a relatively minimal elliptic fibration on the compact Kähler surface X .*

1. Let $f^*(c) := \sum_{j \in J} m_j.D_j$ be any (scheme-theoretic) fibre X_c of f . Then: its multiplicity $m(c, f) := \inf\{m_j\}$ is also equal to: $m^+(c, f) := \gcd\{m_j\}$.
2. There exists a finite étale cover $u : X' \rightarrow X$ such that if $(v \circ f') = (f \circ u)$ is the Stein factorisation of $(f \circ u)$, with: $f' : X' \rightarrow C'$ connected and $v : C' \rightarrow C$ finite, then: f' has no multiple fibre if $g(C') \geq 1$, and at most 2 multiple fibres of coprime multiplicities if C' is rational.
3. Moreover: $g(C') = \kappa(C', f') = \kappa(C, f)$ in the preceding situation.
4. X is special if and only if $\pi_1(X)$ is almost abelian.

Proof: (1) follows from Kodaira's classification of singular fibres of elliptic fibrations ([B-P-V; Chap.V.7]). This equality actually also follows from an elementary argument in the more general case of fibrations with generic fibre a complex torus.

Assertion (2) follows from [C98'] and [N]: [N] shows that if a curve C with points $a_1, \dots, a_m$, affected with multiplicities $n_1, \dots, n_m$ is given, there exists a cover C' of C ramified above the a_i 's only, each point above any a_i having ramification exactly n_i . The only exception is when $C = \mathbb{P}^1$, $m = 1, 2$, and when $n_1 \neq n_2$ if $m = 2$. In [C98'], it is shown (it is a simple computation) that the base change over C' leads to the sought after étale cover $u : X' \rightarrow X$.

Then property (3) follows from Theorem 1.7 for the second equality, and from the fact that $m = 0$ if $g(C') \geq 1$.

We show (4): if X is special then $\kappa(C, f) \leq 0$, so that C' is rational or elliptic. We apply [C98'], which shows that the natural sequence of maps:

$$\pi_1(F') \rightarrow \pi_1(X') \rightarrow \pi_1(C') \rightarrow 1$$

is exact, with F' a generic fibre of f' (so that F is an elliptic curve, and $\pi_1(F') \cong \mathbb{Z}^{\oplus 2}$). Thus $\pi_1(X')$ is almost abelian if X' is special, which is true if so is X (because f' is special and $\kappa(C', f') = \kappa(C, f)$).

Conversely, assume that $\pi_1(X)$ is almost abelian. Then so is $\pi_1(C')$, and C' is either rational or elliptic. Thus $\kappa(C, f) \leq 0$, and X is special by 3.30 $\square$

3.6. Higher Kodaira Dimensions. We shall define higher Kodaira dimensions of any connected manifold $X \in \mathcal{C}$ as follows (this works for compact connected manifolds as well, actually): The first Kodaira dimension of X is the usual one: $\kappa(X)$.

If $\kappa(X) = -\infty$, the second Kodaira dimension of X is not defined.

Otherwise: $\kappa(X) \geq 0$, and $J_X : X \rightarrow J(X)$, the Iitaka-Moishezon fibration of X is defined.

Let then: $\kappa'(X) := \kappa(J(X), J_X)$.

We of course have: $\kappa(X) := \dim(J(X)) \geq \kappa'(X) \geq \kappa(J(X)) \geq -\infty$.

If $\kappa'(X) = -\infty$, the next Kodaira dimension ($\kappa''(X)$) is not defined.

Otherwise: $\kappa'(X) \geq 0$.

Let then $J_X^{(0,1)} : J(X) \rightarrow J'(X)$ be the Iitaka fibration defined on $J(X)$ by the $\mathbb{Q}$ -divisor: $(K_{J(X)} + \Delta(J_X))$ (for any admissible model of the fibration J_X).

Define: $J'_X := J_X^{(0,1)} \circ J_X : X \rightarrow J'(X)$.

Define next $\kappa''(X) := \kappa(J'(X), J'_X) \geq \kappa(J'(X)) \geq -\infty$.

Of course, we have: $\kappa(X) \geq \kappa'(X) \geq \kappa''(X) \geq -\infty$, for the invariants so defined.

Continuing inductively, we can define a decreasing sequence of invariants:

$\kappa(X) \geq \kappa'(X) \geq \dots \geq \kappa^{(r)}(X) \geq -\infty$, and iterated orbifold Iitaka fibrations

$J_X^{(r)} : X \dashrightarrow J^{(r)}(X)$. If the sequence is defined til $J_X^{(r)}$, define $\kappa^{(r+1)}(X) := \kappa(J^{(r)}(X), J_X^{(r)})$, and if this is nonnegative, define $J_X^{(r,r+1)}$ as being the Iitaka fibration defined by the $\mathbb{Q}$ -divisor $(K_{J^{(r)}(X)} + \Delta(J_X^{(r)}))$ (for any admissible model of the fibration $J_X^{(r)}$).

We thus have fibrations $J_X^{(r-1,r)} : J^{(r-1)}(X) \dashrightarrow J^{(r)}(X)$ such that $J_X^{(r)} = J_X^{(r-1,r)} \circ J_X^{(r-1)}$.

Observe also that $\kappa(F_r, J_X^{(r)}) = 0$, if F_r is a general fibre of $J_X^{(r-1,r)}$, by the standard property of Iitaka fibrations.

This sequence stops at the first term (if any) equal to $-\infty$, and is stationary if any two terms $\kappa^{(r)}(X) = \kappa^{(r+1)}(X)$ are equal (and nonnegative, necessarily). This happens clearly if and only if the corresponding map $J_X^{(r)}$ is of general type.

The following is easily shown by induction on r :

Proposition 3.34. *The sequence of higher Kodaira dimensions is invariant under bimeromorphic maps and finite étale covers.*

We shall later extend these notions, and even conjecture that these higher Kodaira dimensions are invariant under deformation (for X Kähler).

As an illustration for the introduction of these invariants, we show the:

Proposition 3.35. *Let $X \in \mathcal{C}$ be smooth of dimension $n \geq 2$. Then $\text{ess}(X) = n - 2$ in each of the following cases (a-e). Moreover, the core c_X and its generic fibre F (a special surface) are described as follows:*

- a. $\kappa(X) = (n-1)$, $\kappa'(X) = \kappa''(X) = n-2$. Then $c_X = J'_X$, and $\kappa(F) = 1$, $\kappa'(F) = 0$.
- b. $\kappa(X) = (n-1)$, $\kappa'(X) = -\infty$, there exists $r : J(X) \dashrightarrow Z$ with $\dim(Z) = (n-1)$ such that $f \circ J_X$ is of general type. Then $\kappa(F) = 1$, and $\kappa'(F) = -\infty$.
- c. $\kappa(X) = \kappa'(X) = n-2$. Then $c_X = J_X$, and $\kappa(F) = 0$.
- d. $R(X)$ (the rational quotient of X) is of general type, of dimension $(n-2)$. Then F is a rational surface.
- e. $R(X)$ has dimension $(n-1)$, and $\kappa(R(X)) = \kappa'(R(X)) = (n-2)$. Then F is birationally elliptic ruled.

Proof: Case (a): Indeed, the fibre of J'_X is special, because F has an elliptic fibration $J : F \rightarrow C$ with $\kappa(C, J) = 0$, so the assertion follows from 3.30. By assumption, J'_X is of general type, because $\kappa'(X) = \kappa''(X)$. To show that $\kappa(F) = 1$, use the easy addition theorem, applied to X and J'_X : it says that $(n-1) = \kappa(X) \leq \kappa(F) + \dim(J'(X)) = \kappa(F) + (n-2) \leq \dim(F) + (n-2) = (n-1)$. We shall skip the proofs of the other cases, which are easier or similar $\square$

Remark 3.36. We shall also nearly show the converse, as a consequence of additivity theorems, in section 6.4. Actually, the converse holds under the general additivity conjecture. The formulation of case (b) is unnatural. A natural formulation would require the conjectural notion of rational quotient for orbifolds. See 6.4.

3.7. Threefolds. We shall describe the core of a compact Kähler threefold. For this we shall need Theorem 5.1 shown later in section 5, which says that X is special if $\kappa(X) = 0$, in all dimensions.

Theorem 3.37. Let $X \in \mathcal{C}$ be a nonspecial threefold. The core c_X of X is a fibration of general type. Moreover, one can describe c_X , its generic fibre F , and $\text{ess}(X)$ as follows:

1. $\text{ess}(X) = 3$ iff $\kappa(X) = 3$. Then $c_X = \text{id}_X$.
2. $\text{ess}(X) = 2$ in the following two cases:
 - a. $\kappa(X) = \kappa'(X) = 2$. Then $c_X = J_X$ is an elliptic fibration of general type.
 - b. $\kappa(X) = -\infty$, $R(X)$ is a surface of general type, and $c_X = r_X$ is a $\mathbb{P}^1$ -fibration over $R(X)$.
3. $\text{ess}(X) = 1$ iff one of the following cases occurs:
 - a. $\kappa(X) = 2$, $\kappa'(X) = \kappa''(X) = 1$. Then $c_X = J'_X$ is a fibration of general type onto a curve, with F a special surface with $\kappa(F) = 1$ and $\kappa'(F) = 0$.
 - b. $\kappa(X) = 2$, $\kappa'(X) = -\infty$, X nonspecial. Then c_X is a fibration of general type onto a curve with $\kappa(F) = 1$ and $\kappa'(X) = -\infty$.
 - c. $\kappa(X) = \kappa'(X) = 1$. Then $c_X = J_X$ is a fibration of general type onto a curve with $\kappa(F) = 0$.
 - d. $c_X = r_X$ is a fibration onto a curve of general type, with F a rational surface.
 - e. $c_X = J_{R(X)} \circ r_X$ is a fibration of general type over a curve with F a birationally ruled elliptic surface.

We now give a very rough list of the special threefolds:

Theorem 3.38. Any special threefold $X \in \mathcal{C}$ is one of the following:

1. $\kappa(X) = 2$.

- a. $\kappa'(X) = 1 > \kappa''(X)$: J'_X is a non-general type fibration over a curve with fibre F a special surface with $\kappa(F) = 1 > \kappa'(F) \in \{0, -\infty\}$.
 - b. $\kappa'(X) = 0$: J_X is an elliptic fibration with a klt orbifold base a normal surface with torsion canonical bundle $(K_S + \Delta)$. (The log-Enriques case).
 - c. $\kappa'(X) = -\infty$: X has either a non-general type fibration over a curve with generic fibre a special surface F with $\kappa(F) = 1, \kappa'(F) = -\infty$, or an elliptic fibration with base orbifold a klt normal surface with Picard number one, and log-Del Pezzo (ie: $-(K_S + \Delta)$ is ample).
2. $\kappa(X) = 1$: then J_X is a non-general type fibration over a curve with generic fibre F a surface with $\kappa(F) = 0$.
 3. $\kappa(X) = 0$
 4. $\kappa(X) = -\infty$: X is either rationally connected, or a fibration over an elliptic curve with generic fibre a rational surface, or a $\mathbb{P}^1$ -fibration over a special surface S with $\kappa(S) \geq 0$. (The case where X is simple non-Kummer conjecturally does not exist, but strictly speaking additionally belongs to the last part (4) of the above list, because $\kappa(X) \leq 0$, then).

Proof: We shall prove both results 3.37 and 3.38 at the same time.

The case where $\kappa(X) = -\infty$ is clear from the section 12.26 above, because X is then uniruled if X is projective (by [Miy]), and by [C-P] otherwise if X is not simple.

So we proceed case-by-case, assuming $\kappa(X) \geq 0$.

If $\kappa(X) \geq 0$, we are done by Theorem 5.1. If X is non-projective, we could also have applied [C-P], theorem (8.1), which says that if X has a nonzero holomorphic 2-form, it is covered by either a torus, or by the product of an elliptic curve and a $K3$ -surface. Thus X is special in this case, too. (The existence of a nonzero 2-form when X is non-projective is a famous result of K.Kodaira).

We now classify the cases occuring according to the pairs $(\kappa(X), \kappa'(X))$, $\kappa(X) \geq 1$.

When the two terms are equal and positive, we conclude from 3.8 that J_X is the core, and of general type.

If $\kappa(X) = 1$, and $\kappa'(X) \leq 0$, we conclude from 3.14 that X is special.

We are thus left with the cases where $\kappa(X) = 2$, $\kappa'(X) \leq 1$.

We thus now assume that $\kappa(X) = 2$.

Assume first that $\kappa'(X) = 1$. Consider the map $J'_X : X \rightarrow J'(X)$ (using the notations of 3.6): it has general special fibres, because if F is its general fibre, and $G := J_X(F)$, then the restriction to F of J_X defines $J' : F \rightarrow G$, which has generic fibres elliptic curves, while G is a curve and, by the definition of $J_X^{0,1}$, we have: $\kappa(G, J') = 0$. We conclude from 3.14 that F is a special surface. The easy addition theorem shows that $\kappa(F) = 1$.

If $\kappa''(X) = 1$, J'_X is a fibration both special and of general type. So we conclude from 3.8 that it is the core of X .

Otherwise (if $\kappa'''(X) \leq 0$), X is special. We are thus done with this case ($ess(X) = 0$, or 1).

We are now left with the more difficult case when $\kappa'(X) \leq 0$. Assume that X is not special. Let $f : X \rightarrow Y$ be a fibration of general type. The fibration J_X being special, we get from (1.12) a factorisation $\phi : J(X) \rightarrow Y$ of $f = \phi \circ J_X$. Thus Y has to be a curve.

To conclude the proof we thus just need to show that this does not happen if $\kappa'(X) = 0$, because of the Minimal Model Program applied to a klt surface orbifold, as described below. The assertion we need follows from Proposition 3.40 below. But we need first some definitions for its statement.

3.8. Orbifold Surfaces. A **surface orbifold** will be a klt pair (S, Δ) , with S a normal projective surface, together with an orbifold divisor Δ on it.

If $g : S \rightarrow C$ is a holomorphic fibration, we define the **orbifold base** of $g : (S/\Delta) \rightarrow C$ as the pair $(C, \Delta_g := \Delta(g, \Delta))$, where Δ_g is the Weil $\mathbb{Q}$ -divisor on C defined as follows:

$\Delta_f := \sum_{c \in C} (1 - 1/m(c, g, \Delta)) \cdot c$, with $m(c, g, \Delta) := \inf_{j \in J} \{m_j \cdot n_j\}$, and $g^*(c) := \sum_{j \in J} m_j \cdot D_j$. Here n_j is simply $(1 - d_j)^{-1}$, if $d_j = (1 - 1/n_j)$ is the multiplicity of D_j in Δ .

Remark 3.39. *As in 1.6, we see that if $h : X \rightarrow S$ is a fibration with $\Delta = \Delta(h)$, then $\Delta_f = \Delta(g \circ h)$ (on a suitable model of h).*

We define then as usual the canonical bundle and Kodaira dimension of (C/Δ_f) and (S/Δ) (by definition, $(K_S + \Delta)$ is supposed to be $\mathbb{Q}$ -Cartier).

We then say that g is of **general type** if $\kappa(C/\Delta_f) = 1$, and that (S/Δ) is **special** if $\kappa(S/\Delta) < 2$, and if there is no holomorphic fibration of general type $g : (S/\Delta) \rightarrow C$ onto a curve.

Proposition 3.40. *Let (S/Δ) be a surface orbifold. If $\kappa(S/\Delta) = 0$, then (S/Δ) is special.*

Proof: Assume there exists a general type fibration g on (S/Δ) . We shall show that $\kappa(S/\Delta) \neq 0$. We thus apply the MMP to our initial pair (S/Δ) . This produces a sequence of elementary contractions of the form: $k : (S/\Delta) \rightarrow (S'/\Delta')$, with (S'/Δ') still a klt pair, such that after at most $\rho(S) - 1$ steps, one gets for the final pair (denoted also (S'/Δ')) one of the three basic cases:

- (1) $K_{(S'/\Delta')}$ is nef.
- (2) There is a fibration $g' : S' \rightarrow C'$ onto a curve such that $-K_{(S'/\Delta')}$ is g' -ample, and $\rho(S') = 2$.
- (3) $-K_{(S'/\Delta')}$ is ample, and $\rho(S') = 1$. (“log-Del Pezzo” case).

Notice that at each step, the Kodaira dimension of the pair (S/Δ) is preserved, and that the curve being contracted is rational smooth (because S itself is klt).

We refer to [K-M] and [F-M] for the existence and usual properties of these reduction steps.

In cases (2,3), we have $\kappa(X'/\Delta') = \kappa(X/\Delta) = -\infty$. So these cases do not occur, because we assumed that $\kappa(X/\Delta) = 0$.

Thus: $K_{(S'/\Delta')}$ is nef.

Claim: $D' := (K_{S'} + \Delta') \equiv 0$. Indeed, we have : $(D')^2 = 0$, otherwise we had: $\kappa(X/\Delta) = 2$. From [F-MK],(11.3), we get the claim (their theorem asserts that $\kappa := \kappa(X'/\Delta')$ is equal to the numerical Kodaira dimension of $\kappa(X'/\Delta')$).

The rest of the proof rests on the following two lemmas 3.41 and 3.42. The first one will be proved at the end of this section.

Lemma 3.41. *Let $g : (S/\Delta) \rightarrow C$ be a fibration of general type, with (S/Δ) klt. Let E be a rational curve mapped surjectively onto C by g . Then: $\Delta \cdot E > 2$.*

Corollary 3.42. *Let $k : (S/\Delta) \rightarrow (S'/\Delta')$ is the contraction of an irreducible smooth rational curve E , with: S, S' klt surfaces, $(K_S + \Delta) \cdot E \leq 0$, and $\Delta' := k_*(\Delta)$.*

If $g : (S/\Delta) \rightarrow C$ is a fibration of general type, then there exists $g' : (S'/\Delta') \rightarrow C$ such that $g = g' \circ k$. Moreover, g' is still of general type, and $(K_S + \Delta) = k^*(K_{S'} + \Delta')$ if $(K_S + \Delta).E = 0$.

Proof (of 3.42): The second assertion is clear, if the first one is.

This is because $m(c, g', \Delta) \leq m(c, g', \Delta')$, since in the definition of the left hand side of the inequality, the infimum is taken over a smaller subset.

The first assertion is clear also if $g(C) \geq 1$, because the rational curve E contracted by k cannot be mapped surjectively to C by g .

We shall show that this also cannot happen when C is rational, because g is of general type (actually, as the proof shows, the condition $\kappa(C, g) \geq 0$ is sufficient, even).

We use the following numerical conditions:

- (a) $E^2 \leq 0$ (E exceptional)
- (b) $(K + E).E = -2$ (E rational, smooth)
- (c) $(K + \Delta).E \leq 0$

So we assume by contradiction that E is mapped onto C by g .

We then get: $\Delta.E \leq -K.E = 2 + E^2 \leq 2$, which contradicts 3.41 $\square$

We now complete the proof of 3.40:

We then apply the Minimal model program to S' , but relative to $K_{S'}$. At each step again only smooth rational K -negative curves are contracted. Let (S'', Δ'') be the resulting pair. It has, by 3.42, all properties of (S', Δ') . In particular, $D'' := (K_{S''} + \Delta'') \equiv 0$, and $g'' : (X''/\Delta'') \rightarrow C$ is of general type. Put $K := K_{S''}$.

Assume first that $\kappa(X) \geq 0$.

In addition to the above properties, $K_{S''}$ is nef. By our assumptions, the generic fibre of g'' is elliptic, and Δ'' is “vertical” (ie: contained in fibres of g'').

Thus: $K^2 = 0$, $D''.\Delta'' = 0 = (D'')^2$. Thus: $(D'')^2 = 0$. And so D'' is a union of complete fibres of g'' , by Zariski’s Lemma.

There exists thus an orbifold divisor δ'' on C such that $\Delta'' = (g'')^*(\delta'')$.

Because $\kappa(X'') \geq 0$, and $\kappa(C/\delta'') = 1$, we easily get that $\kappa(X''/\Delta'') \geq 1$ (see Lemma 4.10 below, for example). A contradiction.

We now treat the remaining case in which $\kappa(X) = -\infty$.

Now: $-D''$ is g'' -ample, and every fibre of g'' has an irreducible reduction. (As above, $\Delta'' = (g'')^*(\delta'')$, for an orbifold structure δ'' of general type on C).

We thus have: $\rho(S'') = 2$, and the arguments of [F-MK], theorem 11.2.3, show that:

$K_{S''} + (\Delta'')^{hor} \equiv \lambda F''$, $(\Delta'')^{hor}$ is the horizontal part of Δ'' , defined as usual, and F'' is any fibre of g'' . Here $\lambda \in \mathbb{Q}$ is such that $\lambda \geq (2q - 2)$, with q the genus of C .

Thus $D'' \equiv (\lambda + \deg(\delta'')).F''$ has Kodaira dimension at least 1, because $\deg(\delta'') > (2 - 2q)$.

Contradiction $\square$

We still have to show Lemma 3.41.

Proof of 3.41:

From Hurwitz’s formula, we get: $-2 = -2d + \sum_{e \in E} (r_e - 1)$, where d is the degree of the restriction $h : E \rightarrow C$ of g to E , and for each $e \in E$, r_e is the ramification order of h at e .

Fix $c \in C$. Write:

$$g^*(c) = \sum_{j \in J} m_j \cdot D_j;$$

$$m := \inf\{m_j.n_j, j \in J\},$$

$$n_j := (1 - d_j)^{-1}, d_j := \text{multiplicity of } D_j \text{ in } \Delta.$$

Let $E_c := E \cap S_c$, where $S_c := g^{-1}(c)$ is the fibre of g over c .

Claim: $\sum_{e \in E_c} (r_e - 1) \geq (1 - 1/m).d - \Delta_c.E$, where Δ_c is the union of components of Δ contained in S_c , with their corresponding multiplicities.

Then Lemma 3.41 is an easy consequence of the claim. Indeed:

$$\begin{aligned} -2 &= -2d + \sum_{e \in E} (r_e - 1) \\ &\geq -2d + \sum_{c \in C} \sum_{e \in E_c} (r_e - 1) \\ &\geq -2d + g^*(\delta).E - \Delta.E, \text{ if } \delta := \Delta(g, \Delta) \\ &= (\deg(\delta) - 2).d - \Delta.E \end{aligned}$$

But $\deg(\delta) - 2 > 0$, because g is of general type. Hence the conclusion of 3.41.

To complete the proof, we establish the preceding **claim**.

Proof of the claim:

$$\begin{aligned} (1 - 1/m).g^*(c).E &= (1 - 1/m). \sum_{j \in J} m_j.(D_j.E) = \sum_{j \in J} (m_j - m_j/m).(D_j.E) \\ &\leq \sum_{j \in J} ((m_j - 1) + (1 - 1/n_j)).(D_j.E) \text{ (because } m_j.n_j \geq m, \forall j \in J) \\ &= (\sum_{j \in J} (m_j - 1).(D_j.E)) + \Delta.E. \end{aligned}$$

We are thus reduced to show that:

$$\sum_{j \in J} (m_j - 1).(D_j.E) \leq \sum_{e \in E_c} (r_e - 1).$$

Because $\sum_{j \in J} m_j.(D_j.E) = \sum_{e \in E_c} r_e = d$, we just need to establish that:

$$\sum_{j \in J} D_j.E \geq \sum_{e \in E_c} 1, \text{ which itself follows from the inequality:}$$

$\sum_{j \in J} (D_j.E)_e \geq 1, \forall e \in E_c$, where the intersection number $(D_j.E)_e$ is the local intersection number near e . (Recall that S is only assumed to be normal).

We now show this last inequality.

Indeed: we have $\sum_{j \in J} (m_j.D_j)_e = r_e$, and the conclusion follows from the inequality:

$$m_j \leq r_e, \forall j \in J.$$

To show this inequality, we make a base change over $h : E \rightarrow C$. Let X' be the normalisation of the fibre product $X \times_C E$, and let $k : X' \rightarrow X$, $g' : X' \rightarrow E$ such that $k \circ g = h \circ g$ be the natural maps induced from this base change.

Let $E' \subset X'$ be the lift of E to X' : it is a section of g' . Let e' be the point of E' lying above e .

The components of $(g')^*(c')$ which contain e' are thus reduced. This easily implies the conclusion, by looking at a generic point of D_j , near which the projection g is locally given by the equation $g(t, z) = z^{m_j}$. The fibre product is thus locally given by an equation of the form: $\zeta^{r_e} = z^{m_j}$.

Dividing by $d := \gcd(r_e, m_j)$, we can assume that $d = 1$, since we normalised the fibre product.

Thus we have on X' local coordinates (t, s) with $\zeta = s^{m_j}$, and $z = s^{r_e}$.

Locally, the projection g' is given in these coordinates by $g'(t, s) = \zeta$. From which we deduce that $m_j = 1$ Thus m_j divides r_e . This in particular proves the claimed inequality $\square$

Remark 3.43. *What we actually proved in this section is the orbifold Additivity for Kodaira dimensions in dimension 2. The general case is stated in the next section at 4.4.*

4. Orbifold Additivity

4.1. Orbifold Conjecture $C_{n,m}$. We use the notations and notions introduced in section 1.6.

So, if $g : (Y/H) \rightarrow Z$ is a holomorphic fibration from the manifold Y equipped with the orbifold divisor H , we defined the base orbifold $\Delta = \Delta(g, H)$ on Z .

The fundamental property of this definition is that, when $H = \Delta(f)$, for some fibration $f : X \rightarrow Y$, then $\Delta = \Delta(g \circ f)$ (for suitable models of f, g which can be chosen so that $f, g, g \circ f$ are admissible).

Definition 4.1. *We shall say that $g : (Y/H) \rightarrow Z$ is **high** (resp. **very high**) if there exists a modification $u_0 : Y \rightarrow Y_0$ with Y_0 smooth such that (a) and (b) (resp. (a) and (b')) below are satisfied:*

(a) *Every g -exceptional divisor of Y is u_0 -exceptional.*

(b) $\kappa(Y/H) = \kappa(Y_0/H_0)$, with $H_0 := (u_0)_*(H)$.

(b') $K_Y + H \geq (u_0)^*(K_{Y_0} + H_0)$.

The following properties are immediate (using the arguments of 1.11 and 1.28, for (3)):

Proposition 4.2. (1) *If g is very high, it is high.*

(2) *If g is high, then $\kappa(Y, K_Y + H + B) = \kappa(Y, K_Y + H)$, for any effective g -exceptional $\mathbb{Q}$ -divisor B on Y .*

(3) *If $H = \Delta(f)$, then one can choose models of f, g in such a way that g is high, prepared, that $\Delta(g \circ f) = \Delta(g, H)$, and $f, g, g \circ f$ are admissible.*

Now we can state the orbifold additivity conjecture $C_{n,m}^{orb}$:

Conjecture 4.3. ($C_{n,m}^{orb}$) *Let $g : (Y/H) \rightarrow Z$ be a holomorphic fibration between manifolds, as above, with $Y \in \mathcal{C}$. Assume g is prepared and high.*

Then: $\kappa(Y/H) \geq \kappa((Y/H)_z) + \kappa(Z/\Delta(g, H))$, where $z \in Z$ is general and $(Y/H)_z := (Y_z/H_z)$.

Of fundamental importance for the considerations of the present paper is the following special case, shown by suitably adapting the classical methods of proof (T.Fujita, Y. Kawamata, E. Viehweg):

Theorem 4.4. (C_{gt}^{orb}) *Let $g : (Y/H) \rightarrow Z$ be a holomorphic fibration between manifolds, as above, with $Y \in \mathcal{C}$, and Z projective. Assume g is prepared, high and of general type (ie: $\kappa(Z/\Delta(g, H)) = \dim(Z)$).*

Then: $\kappa(Y/H) = \kappa((Y/H)_z) + \dim(Z)$, where $z \in Z$ is general and $(Y/H)_z := (Y_z/H_z)$.

Of course, the above $C_{n,m}^{orb}$ is a simple generalisation and refinement of the classical conjecture of S.Iitaka, dictated by the constructions made in the previous chapters.

Let us list some of its corollaries or special cases of the above conjecture:

Proposition 4.5. *Assume $C_{n,m}^{orb}$ holds. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be fibrations, with $X \in \mathcal{C}$. Then: $\kappa(Y, f) \geq \kappa(Y_z, f_z) + \kappa(Z, g \circ f)$, where $z \in Z$ is general and $f_z : X_z \rightarrow Y_z$ is the restriction of f .*

Proof: By 1.6 and 1.25, we can choose models of g and f in such a way that $f, g, g \circ f$ are admissible prepared, with g high, $\Delta(g \circ f) = \Delta(g, \Delta(f))$, and such moreover that f_z is admissible.

We conclude then from $C_{n,m}^{orb}$ and the following equalities:

$\kappa(Y/\Delta(f)) = \kappa(Y, f)$, $\kappa(Y_z/\Delta(f)_z) = \kappa(Y_z, f_z)$, and:

$$\kappa(Z, g \circ f) = \kappa(Z/\Delta(g \circ f)) = \kappa(Z/\Delta(g, \Delta(f))) \quad \square$$

In the special case where $X = Y$, we get:

Proposition 4.6. *Assume $C_{n,m}^{orb}$ holds. Let $g : Y \rightarrow Z$ be a fibration, with $Y \in \mathcal{C}$. Then:*
 $\kappa(Y) \geq \kappa(Y_z) + \kappa(Z, g) \geq \kappa(Y_z) + \kappa(Z)$.

The extreme inequality is of course the classical Iitaka conjecture.

Let us first list some immediate consequences of theorem 4.4:

Corollary 4.7. *Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be fibrations, with $X \in \mathcal{C}$. Assume $(g \circ f)$ is of general type, then: $\kappa(Y, f) = \kappa(Y_z, f_z) + \dim(Z)$.*

Corollary 4.8. *Let $g : Y \rightarrow Z$ be a fibration, with $Y \in \mathcal{C}$. Assume g is of general type, then: $\kappa(Y) = \kappa(Y_z) + \dim(Z)$.*

We shall now give the proof of Theorem 4.4. It is classically done in two steps: first an easy reduction to weak-positivity statements for direct images by g of twisted pluricanonical forms; second the proof of semipositivity. The first step is entirely similar to the known cases, so we shall be brief on it. The second step is simply obtained by introducing the orbifold divisors at appropriate places in the classical proofs of Y.Kawamata and E.Viehweg. See also the initial work [Fuj78].

4.2. Reduction to Weak Positivity. We start by briefly recalling the notion of weak-positivity introduced in [Vi 83] (see also the survey [E]).

A torsionfree coherent sheaf $\mathcal{F}$ on Z , projective, is said to be **weakly-positive** (written (w.p) for short) if for any ample line bundle A on Z , and every $a > 0$, integer, there exists an integer $b > 0$ such that $S^{ab}(\mathcal{F}) \otimes A^b$ is generated over some (nonempty) open subset U of Z by its global sections (defined over Z).

Here $S^{ab}(\mathcal{F})$ denotes the extension to Z of the sheaf denoted by the same symbol, naturally defined over the open subset where $\mathcal{F}$ is locally free.

Remark 4.9. *The following properties are shown in [Vi 82]:*

- (1) *If $\mathcal{F}$ is locally free and nef, it is w.p.*
- (2) *Let $v : Z' \rightarrow Z$ be bimeromorphic, and $\mathcal{F} \subset \mathcal{G}$ a inclusion of torsionfree coherent sheaves of the same rank on Z' . If $\mathcal{F}$ is w.p, then so is $v_*(\mathcal{G})$.*
- (3) *If $v : Z' \rightarrow Z$ is a ramified flat covering, with Z, Z' smooth, and if $\mathcal{F}$ is torsionfree coherent on Z , then $\mathcal{F}$ is w.p if so is $v^*(\mathcal{F})$.*

We now state without proofs two lemmas, shown but not separately stated in [E], and in various more or less implicit forms in [Vi 82] and [Kw 81]. Together with the weak-positivity result shown in the next section, they imply immediately Theorem 4.4.

Lemma 4.10. *Let $g : Y \rightarrow Z$ be a fibration with Z projective. Let E and L be $\mathbb{Q}$ -divisors on Y and Z respectively, such that: L is big and $\kappa(Y, E) \geq 0$. Then:*
 $\kappa(Y, E + g^*(L)) = \dim(Z) + \kappa(Y_z + E|_{Y_z})$, for z general in Z .

The crucial place where weak-positivity enters is to check that $\kappa(Y, E) \geq 0$:

Lemma 4.11. *Let $g : Y \rightarrow Z$ be a fibration, E a line bundle on Y and L a $\mathbb{Q}$ -divisor on Z . Assume L is big and $g_*(E)$ is weakly positive and nonzero. Then: $\kappa(Y, E + g^*(L)) \geq 0$.*

In the next section, we shall show:

Theorem 4.12. *In the situation and hypothesis of Theorem 4.4, for any sufficiently divisible integer $m > 0$, there exists an effective g -exceptional divisor B on Y such that the sheaf: $g_*(m(K_{Y/(Z/\Delta(g,H))}) + H) + B$ is weakly positive on Z . If g is very high, we can choose $B = 0$.*

Let us now explain how to deduce Theorem 4.4 from the preceding lemmas and Theorem 4.12: apply first 4.11 to $E := m.(K_{Y/(Z/\Delta(g,H))}) + H$, with $m > 0$ an integer sufficiently divisible, so chosen that $g_*(E)$ is nonzero (and w.p), and:

$L := (m/2).K_{(Z/\Delta(g,H))}$, which is big by hypothesis.

We conclude that $\kappa(Y, E + g^*(L)) \geq 0$.

Thus $E' = E + g^*(L) + H$ is such that $\kappa(Y, E') \geq 0$.

Next apply 4.10 to E' and L to conclude that $\kappa(Y, K_Y + H + B/m)$ satisfies the inequality stated in 4.4. Use finally the fact that g is high to conclude the proof of 4.4, because $\kappa(Y, K_Y + H + B/m) = \kappa(Y, K_Y + H)$ $\square$

4.3. Orbifold Weak-Positivity. Our objective in the next two sections is to establish Theorem 4.12 stated and used above .

Notation 4.13.

We consider thus a prepared holomorphic fibration $g : Y \rightarrow Z$, with Y and Z smooth and Z projective.

That g is prepared means that its non-smooth locus is contained in a simple normal crossing divisor of Z , and that the inverse image by g of this non-smooth locus is also a divisor of simple normal crossings on Y . We let $\Delta' := \Delta(g)$, and $\Delta := \Delta(g, H)$ be the orbifold divisors for g , and (g, H) respectively, the second one is defined in 1.6 to which we refer for details.

Thus $\Delta' := \Delta(g) = (\sum_{i \in I} (1 - 1/m_i) \cdot \Delta_i)$ also is of simple normal crossings. We define also $K_{Z/\Delta'} = K_Z + \Delta'$, and $K_{Y/(Z/\Delta')} := K_Y \otimes g^*(K_{Z/\Delta'})^{-1}$.

We apply similar notations for Δ . Notice that $\Delta \geq \Delta'$.

Remark 4.14. *Notice that $\Delta(g, H) = \Delta(g, H^{vert})$, where H^{vert} is the g -vertical part of H , equal to $\sum_{k \in K'} (1 - 1/n_k) \cdot H_k$, where $K' \subset K$ consists of the components of $|H|$ which are not mapped onto Z by g .*

We shall also denote by H^{hor} the g -horizontal part of H , defined such that $H = H^{vert} + H^{hor}$.

Our objective in this section is to establish the following orbifold generalisation of famous results of Y.Kawamata and E.Viehweg, initiated by T.Fujita in the case where Z is a curve:

We shall actually essentially just reduce our case to the cases they treated.

Theorem 4.15. *Let $g : Y \rightarrow Z$ be as above, prepared. Let H be an orbifold structure on Y , and $\Delta := \Delta(g, H)$ as above. Assume Z is projective and Y is Kähler compact. Let $m > 0$ be an integer such that all $\mathbb{Q}$ -divisors involved are integral.*

Then: $g_*(m.(K_{(Y/(Z/\Delta))} + H) + B)$ is weakly positive, for B a suitable effective g -exceptional divisor on Y .

We shall obtain this result as the consequence of three intermediate steps.

The first step is a generalisation of the standard weak positivity results for direct image sheaves of pluricanonical forms in Kähler geometry:

Theorem 4.16. *Let $g : Y \rightarrow Z$ be a fibration as in 4.15 above, with the same assumptions. Let $D = \sum_{j \in J} d_j D_j$ be a divisor with integer positive coefficients on Y , the D_j being (as always) pairwise distinct. Assume also that the support of D is a divisor of simple normal crossings, and that D is horizontal, which means that each D_j is mapped surjectively onto Z by g . Let m be a positive integer such that $m \geq d_j, \forall j \in J$. Then: $g_*(mK_{Y/Z} + D)$ is weakly positive.*

This result will be proved in the next section. Observe that the restriction that D is horizontal is obviously unnecessary, because in general, one can remove the vertical part of D , if any, and then get an injection from the direct image sheaf relative to the horizontal part of d into the same sheaf, relative to all of D . Notice that there is then no restriction needed on the multiplicities of the vertical components of D , besides nonnegativity.

The second and third steps are given by the following lemma 4.17 and proposition 4.18:

Lemma 4.17. *Let $g : Y \rightarrow Z$, D and m satisfy the same assumptions as the preceding 4.16. If $v : Z' \rightarrow Z$ is a flat finite map with Z' smooth, and if $g' : Y' \rightarrow Z'$ is deduced from g by smoothing the base change $\hat{Y} := Y \times_Z Z'$ of Y by v , there is a natural injection of sheaves:*

$g'_*(mK_{Y'/Z'} + D') \rightarrow v^*(g_*(mK_{Y/Z} + D))$, where $D' := u^*(D)$, if $u : Y' \rightarrow Y$ is the natural map obtained by composing the desingularisation $d' : Y' \rightarrow \hat{Y}$ with the base change map $\hat{u} : \hat{Y} \rightarrow Y$.

Proposition 4.18. *Let g, D, m be as in 4.17 above. Let, in addition, $H = H^{vert}$ be an orbifold vertical divisor on Y , which means that no component of H is mapped onto Z by g .*

There exists a finite flat map $v : Z' \rightarrow Z$, with Z' smooth, such that if $g' : Y' \rightarrow Z'$ is constructed as in 4.17 above from v , the above injection of sheaves extends to an injection:

$g'_*(mK_{Y'/Z'} + D') \rightarrow v^*(g_*(mK_{Y/(Z/\Delta(g,H))} + H) + D + B)$, for some effective g -exceptional divisor B on Y .

We shall give below the proofs of the preceding lemma and proposition. Let us first show that they imply, together with 4.16, the Theorem 4.15: write $mH = D + mH^{vert}$, with $D := mH^{hor}$. This is an integral divisor, if m is sufficiently divisible. Moreover, $0 < d_j := (1 - 1/m_j).m < m$, for each j . So that 4.16 applies to g' and D' , if g' is deduced from g by any base change $v : Z' \rightarrow Z$ as in Lemma 4.17 above, and if D' is the strict transform by $u : Y' \rightarrow Y$ of D , so that it is the horizontal part (for g') of $D' := u^*(D)$. Actually, by the obvious injection of sheaves, the same result applies also if $D' := u^*(D)$, because then only effective vertical divisors are added.

If we now apply the Proposition 4.18, we see that the conclusion of 4.15 holds.

We shall now prove 4.17 and 4.18.

Proof of 4.17: We use the notations of 4.17. It is proved in [Vi83], Lemma 3.3, pp. 335-336 to which we refer, that in our situation, we have: $d_*(mK_{Y'/Z'})$ naturally injects into $\hat{u}^*(mK_{Y/Z})$. This statement implies that $d_*(mK_{Y'/Z'} + D')$ injects into $\hat{u}^*(mK_{Y/Z} + D)$, since $D' = u^*(D)$. We then just need to apply $\hat{g}_*$ to both sides, noticing that $\hat{g}_*(\hat{u}^*) = v^*(g_*)$, by flatness of v .

Here $\hat{g} : \hat{Y} \rightarrow Z'$ is, of course, deduced from g by the base change v $\square$

Proof of 4.18: We start with the construction of v :

Definition 4.19. *Let g as above. A finite covering $v : Z' \rightarrow Z$ is said to be $\widetilde{\Delta}'$ -nice (in similarity to $[Kw]$) if :*

(1) *It is flat, and Z' is smooth.*

(2) *$v^*(\Delta_i) = \widetilde{m}_i \Delta_i''$, for some reduced divisor $\Delta_i'' \subset Z'$, this for any $i \in I$.*

Here $\widetilde{m}_i$ is any integer divisible by $\text{lcm}(m_{ij}$'s), these being the same as above, used to define the multiplicity of g along Δ_i' . Observe in particular that $v^(\Delta')$ is Cartier on Z' .*

(3) *$v^{-1}(|\Delta'|)$ is a divisor of normal crossings on Z' .*

By [Kw] and [Vi82], such coverings exist.

Construction: Let $v : Z' \rightarrow Z$ be a $\widetilde{\Delta}'$ -nice covering. Let $u : Y' \rightarrow Y$ be the composition $u = u'' \circ n \circ d$, in which: $u'' = \hat{u} : Y'' := \hat{Y} = (\widehat{Y \times_Z Z'}) \rightarrow Y$ is the base change by v , $n : Y'' \rightarrow Y''$ is the normalisation, and: $d : Y' \rightarrow Y^n$ is a desingularisation isomorphic above the smooth locus of Y^n . One thus also has, with the previous notations of 4.17: $d' = n \circ d$.

We also denote by $\hat{g} = g'' : Y'' \rightarrow Z'$ and $g' := g \circ d : Y' \rightarrow Z'$ the maps deduced from g .

Then 4.18 is an immediate consequence of 4.9 and the following:

Proposition 4.20. *: In the preceding situation, there exists on Z' a natural injection of sheaves: $u_* : (g')_*(m \cdot (K_{Y'/Z'} + D' + (g')^*((v)^*(m\Delta))) \rightarrow v^*(g_*(m \cdot (K_{Y/Z} + H^{\text{vert}}) + D + B))$, this for any integer $m > 0$ sufficiently divisible, and some effective g -exceptional divisor B on Y .*

Proof: The main point is that we just need to check this injection on the complement of a codimension two subvariety A of Z' , since by 4.17, the result holds with $(g')^*(v^*(m\Delta))$ deleted from the left-hand side, and H^{vert} deleted from the right-hand side, on all of Z' . This provided $\mathcal{O}_Y(B)$ is defined by the poles of maximal order acquired by an arbitrary extension across A of a section of $(g')_*(m \cdot (K_{Y'/Z'} + (g')^*((v)^*(m\Delta)))$ defined outside of A .

Observe that we are working here on the fibre product $Y \times_Z Z'$ which is Cohen-Mac Cauley by the normal crossing hypothesis on Δ' , so that the poles of the sections considered actually occur in codimension one. (This observation is due to S.Druel).

We shall thus check the above injection only above the generic point z of some Δ_i . Let U be a sufficiently small analytic open neighborhood of z in Z .

Let $Y_U := g^{-1}(U)$, and let $W \subset Y$ be a small analytic neighborhood of y , a generic point of D_{ij} , notations being as in (1.1.2).

Thus m_{ij} divides $\widetilde{m}_{ij} := m_{ij} \cdot q$, for some integer $q = q_{i,j}$, by the definition of a $\widetilde{\Delta}'$ -nice covering. Factorise $v := v' \circ v^*$ over U , with $v' : Z' \rightarrow Z^*$, and $v^* : Z^* \rightarrow Z$, in such a way that v^* (resp. v') ramifies at order exactly m_{ij} (resp. q) above Δ_i (resp. $\Delta_i^* := (v^*)^{-1}(\Delta_i)$).

Construct Y^* from Y over Z by taking base change by v^* , followed by normalisation and then smoothing. We have also a natural fibration $g^* : Y^* \rightarrow Z^*$.

Possibly modifying Y' , we thus get a factorisation $u = u^* \circ u'$. Moreover, from the usual commutation properties, we have: $v' \circ g' = g^* \circ u'$, and $v^* \circ g^* = g \circ u^*$.

The crucial property in this construction is that u^* is étale over W , if sufficiently small. This is an easy standard local computation which we already used several times before (see [C2], for example). Thus u^* is étale over the generic point of D_{ij} .

From the following lemma 4.21 below, we deduce that, over the generic point of D_{ij} , we have a natural injection of sheaves: $m.(K_{Y^*/Z^*}) \subset (u^*)^*(m.(K_{Y/(Z/\Delta)} + H^{vert}))$, for any integer $m > 0$ sufficiently divisible.

From this injection, we can deduce the following, by tensorising with $\mathcal{O}_Y(D)$ and its lift $\mathcal{O}_{Y^*}(D^*)$ to Y^* by (u^*) : $m.K_{(Y^*/Z^*)} + D^* \subset (u^*)^*(m.(K_{Y/(Z/\Delta)} + H^{vert}) + D)$

The very same argument as in the proof of 4.17 above shows the existence, for any j (on which the preceding factorisations depend), of a natural injection of sheaves:

$$g'_*(m.(K_{Y'/Z'}) + D') \subset (v')^*((g^*)_*(m.(K_{Y^*/Z^*}) + D^*)).$$

By composing the above injections, and restricting over W , we see that the sections of:

$$(g^*)_*(m.(K_{Y^*/Z^*}) + D') \text{ actually belong to } (v^*)^*(g_*(m.(K_{Y/(Z/\Delta)} + H^{vert}) + D)).$$

(We identify local sections of $g'_*(m.(K_{Y'/Z'}) + D')$ over Z' and sections of $(m.(K_{Y'/Z'}) + D')$ over corresponding open subsets of Y').

The conclusion now follows from Hartog's extension theorem, applied over Y_U , to extend the sections thus obtained across the intersection of two or more such $D_{i,j}$'s $\square$

We used the following:

Lemma 4.21. *With the above notations, over the generic point of D_{ij} , we have a natural injection of sheaves: $m.(K_{Y^*/Z^*}) \subset (u^*)^*(m.(K_{Y/(Z/\Delta)} + H^{vert}))$, for any integer $m > 0$ sufficiently divisible.*

Proof: We shall argue using, instead of sheaf injections, rather inequalities between $\mathbb{Q}$ -divisors; the inequality $A \geq B$ meaning as usual that $(A - B)$ is effective.

From the equalities:

$$K_{Y^*} = (u^*)^*(K_Y) \text{ (because } u^* \text{ is étale), and:}$$

$$K_{Z^*} = (v^*)^*(K_Z + (1 - 1/m_{i,j}).\Delta_i) \text{ (ramification formula), we deduce that:}$$

$$K_{Y^*/Z^*} = (u^*)^*(K_{Y/Z} - (m_{i,j} - 1).D_{i,j}), \text{ because } g^*(\Delta_i) = m_{i,j}.D_{i,j}, \text{ and } g \circ u^* = v^* \circ g^*.$$

On the other hand:

$$\begin{aligned} K_{Y/(Z/\Delta)} + H^{vert} &= K_{Y/Z} - g^*((1 - 1/m'_i).\Delta_i) + (1 - 1/n_{i,j}).D_{i,j} \\ &= K_{Y/Z} + (-m_{i,j}.(1 - 1/m'_i) + (1 - 1/n_{i,j})).D_{i,j} \\ &= K_{Y/Z} + (-m_{i,j} + 1 + (m_{i,j}/m'_i) - (1/n_{i,j})).D_{i,j} \geq K_{Y/Z} - (m_{i,j} - 1).D_{i,j}, \text{ since: } m_{i,j}.n_{i,j} \geq m'_i, \\ &\text{by the very definition of } m'_i. \end{aligned}$$

This concludes the proof, by applying $(u^*)^*$ $\square$

4.4. Twisted Weak Positivity. The aim of this section is to prove Theorem 4.16. For this, we shall refer to the proof given in [Vi83] of the classical case where D is empty, and simply indicate the changes needed. I would like to heartily thank E.Viehweg, who gave me a decisive hint for the proof of the following lemma.

The single change needed lies in the corollary 5.2 of [Vi83]. We restate this corollary in the form we need.

Lemma 4.22. *Let $g : Y \rightarrow Z$ be as above. Let A^* be ample on Z , let $A = g^*(A^*)$.*

Assume that $S^N(g_*(mK_{Y/Z} + D + mA))$ is generated by its global sections on some nonempty open subset of Z .

Then: $g_*(mK_{Y/Z} + D + (m-1)A)$ is weakly positive on Z .

(One can thus withdraw one A in the above situation. This is sufficient to conclude to the desired weak positivity. See [Vi83] for the ingenious details.)

Proof: We shall closely follow the proof of the same corollary 5.2 (where D is empty) in [Vi83]. The proof in fact reduces to add D everywhere in an appropriate way.

Observe also that the proof given there uses only the fact that $g_*(K_{Y/Z})$ is weakly positive when Y is projective. The case when Y is Kähler can be obtained from the different proof of this fact sketched in [Vi86], based on [Ko86] (see the references in [Vi86] for more details), because it uses only the Hodge-theoretic result due to P.Deligne that holomorphic forms with logarithmic poles on a compact Kähler manifold are d -closed.

Take now $L := K + D + A$, with $K := K_{Y/Z}$. Define next;

$$M := \text{Image } (g^*(g_*(mK + D + mA)) \rightarrow (m(K + D + A)))$$

Assume that the base locus of $(NmL - (m-1).ND)$ does not contain any component of D . We can of course always easily reduce to this case, by diminishing the relevant d_j 's, without modifying the conclusion.

We now reproduce the proof of [Vi83, 5.2]. Blowing up Y if needed, we can assume that M is a line bundle, and that $NmL = NmM + E + (m-1)ND$, for E an effective divisor on Y , not containing any component of D , such that $E + D$ has a support of normal crossings. By hypothesis, $Nm.M$ is generated by global sections over a nonempty Zariski open subset of Z .

We first treat the case in which $m > d_j, \forall j \in J$.

The corollary 5.1 of [Vi83] then applies without any change and shows that the subsheaf:

$$g_*(K + L^{(m-1)}) \subset g_*(K + (m-1)L)$$

is weakly positive, in the notations $L^{(m-1)}$ of [Vi82, (2.2)].

Here $L^{(m-1)} := (m-1).L - [((m-1)/mN).(E + (m-1)ND)]$, the integral part of a $\mathbb{Q}$ -divisor being computed just by taking the integral part of the coefficients, componentwise.

From the constructions made, we see that $g_*(K + L^{(m-1)}) \subset g_*(K + (m-1)L - [((m-1)^2/mN).D])$, and that they coincide over the generic point of Z .

The sheaf on the right is thus weakly positive, too.

Computing, we get:

$$\begin{aligned} K + (m-1)L - [((m-1)^2/mN).D] &= K + (m-1)L - \sum_{j \in J} [(m-1)^2 m_j / m].D_j \\ &= mK + \sum_{j \in J} ((m-1)m_j - (m-2)m_j + [m_j/m]).D_j = mK + D, \text{ as desired.} \end{aligned}$$

This ends the proof of the special case (all $d_j < m$) considered.

When $d_j = m$ for certain components Δ_j of D , so that $D = D^- + m\Delta$, each component D_j of D^- having $d_j < m$, one just needs to replace K by $K + \Delta$ in the above proof, using the fact that $g_*(K + \Delta)$ is weakly positive on Z , by [Kw, Thm 32] (which can be proved also using the above result of Deligne in the Kähler case) $\square$

Once this Lemma 4.22 is obtained, the rest of the proof of the weak positivity of $g_*(mK + D)$ is the same as the one given in [Vi83] when D is empty.

5. Geometric Consequences of Additivity

5.1. Varieties with $\kappa=0$. The second fundamental example of special manifold is given by the:

Theorem 5.1. *Assume $X \in \mathcal{C}$ has: $\kappa(X) = 0$. Then X is special.*

Proof: This is an immediate application of Theorem 4.4: let indeed $f : X \rightarrow Y$ be a fibration of general type. Then the results apply and give: $\kappa(X) \geq \kappa(X_y) + \dim(Z)$. But $\dim(Z) > 0$, by hypothesis, and $\kappa(X_y) \geq 0$. Contradiction $\square$

Remark 5.2. *This theorem together with Theorem 2.26 shows that X has no Bogomolov sheaf if $\kappa(X) = 0$. Actually, more is expected: $\kappa^+(X) = 0$ if $\kappa(X) = 0$ (see [C95] for details and definitions). This equality is shown for X projective (or Kähler, using Yau's results) if $c_1(X) = 0$.*

5.2. The Albanese Map. The following result uses indirectly only the easiest part of the Additivity theorem 4.4. It nevertheless seems to belong to this section.

Proposition 5.3. *Let $X \in \mathcal{C}$ be special. Let $\alpha_X : X \rightarrow \text{Alb}(X)$ be the Albanese map of X . Then: α is surjective, connected, and has no multiple fibres in codimension one (ie: $\Delta(\alpha)$ is empty). If X is only w-special (see section 6.6 below for this notion), then α is surjective and connected.*

Proof: Assume $\alpha := \alpha_X$ is not onto.

Let $Z \subset \text{Alb}(X)$ be its image. After ([U],(10.9)) , there exists a fibration $g : Z \rightarrow W$ with W of general type (thus $\dim(Y) > 0$). Let $\psi : X \rightarrow W'$ and $\sigma : W' \rightarrow W$ be the Stein factorisation of $\phi \circ \alpha = \sigma \circ \psi$. Then W' is of general type, too, since σ is finite. This contradicts the fact that X is special. Thus α is onto, and $Z = \text{Alb}(X)$.

Let now $\alpha = \beta \circ \alpha'$ be the Stein factorisation of α , with $\alpha' : X \rightarrow A'$ connected and $\beta : A' \rightarrow \text{Alb}(X)$ finite. A slight variation of the arguments of [K-V] shows that if β is ramified, there exists a fibration $\phi : A' \rightarrow W'$, with W' of general type. Considering $\phi \circ \alpha'$, we get as above a contradiction to the fact that X is special. Thus β is unramified, hence isomorphic, by the universal property of α , which is thus connected and onto.

Assume now that the fibration $\alpha : X \rightarrow \text{Alb}(X)$ has $\Delta := \Delta(\alpha)$ nonempty. Let Δ' be any component of Δ .

There exists a connected submersive quotient map $q : \text{Alb}(X) \rightarrow A := \text{Alb}(X)/B$, for some subtorus B of $\text{Alb}(X)$, such that $\Delta' = q^{-1}(D')$, for some big $\mathbb{Q}$ -divisor $D' \subset A$.

Consider the fibration $f := q \circ \alpha : X \rightarrow A$: the support of $\Delta(f) \subset A$ obviously contains D' . We conclude from 1.12 that f is of general type (since K_A is trivial and D' is big on A). Again this contradicts the assumption of X being special. And concludes the proof if X is special.

If X is only w-special, the first two steps of the proof apply without any change to give the last assertion $\square$

Question 5.4. *Let $X \in \mathcal{C}$ be a special manifold. Are then the generic fibres of α_X special ?*

One can easily show that this question has a positive answer if so does the conjecture $C_{n,m}^{\text{orb}}$.

5.3. Varieties of General Type. Although the results of this section are easy consequences of the main result 5.8 below, we give a direct proof.

Theorem 5.5. *For any manifold $X \in \mathcal{C}$, we have: $\text{ess}(X) = \dim(X)$ if and only if X is of general type.*

In particular, X is generically covered by a nontrivial family of special submanifolds (the fibres of c_X) if X is not of general type. The case when $\kappa(X) \geq 0$ is clear, by theorem 5.1 applied to the fibres of the Iitaka-Moishezon fibration of X . So the result applies nontrivially only when $\kappa(X) = -\infty$, and its proof gives the following:

Theorem 5.6. *Let $X \in \mathcal{C}$ be such that: $\kappa(X) = -\infty$ and $\dim(X) > 0$. Then $c_X : X \rightarrow C(X)$ has a general fibre X_y which is special of positive dimension, with: $\kappa(X_y) = -\infty$.*

The proofs are easy applications of Theorem 4.4:

Proof of 5.5 Assume first that X is of general type. So the identity map id_X of X is a special fibration of general type. It is thus the core by 3.8.

Conversely, assume that $ess(X) = \dim(X)$, or what is the same, that: $c_X = id_X$, and that $n := \dim(X) > 0$, so that X is not special.

There exists then a fibration of general type $f : X \rightarrow Y$, with $m := \dim(Y) > 0$. We proceed by induction on $n > 0$. So the assertion holds for manifolds of dimension smaller than n .

If F is a general fibre of f , then F is of general type.

Indeed: otherwise, we could construct a relative core for f , by theorem 12.27, and deduce a nontrivial fibration on X with general fibre special, contradicting our initial hypothesis.

Thus F is of general type. But now f is a fibration of general type with general fibre itself of general type.

By theorem 4.4, we get that X is of general type, as claimed $\square$

Proof of 5.6: We get first from the preceding argument that $\dim(X_y) > 0$, otherwise $\kappa(X) = n$. Assume that $\kappa(X_y) \geq 0$. From Theorem 4.4 again, we learn that $\kappa(X) = -\infty \geq \dim(Z) + \kappa(X_y) \geq \dim(Z) \geq 0$. Contradiction $\square$

In a similar way, we can get a very simple description of the next step: $ess(X) = \dim(X) - 1$.

Theorem 5.7. *Let $X \in \mathcal{C}$ be a manifold of dimension $n > 0$. Then $ess(X) = (n - 1)$ if and only if one of the following two cases occurs:*

- (a) $\kappa(X) = (n - 1)$ and J_X is a fibration of general type.
- (b) $R(X)$ (the rational quotient of X) is of dimension $(n - 1)$ and of general type.

Proof: The fact that cases (a), (b) imply that $ess(X) = (n - 1)$ was already shown in 3.12. So assume $c_X : X \dashrightarrow C(X)$ has $(n - 1)$ -dimensional image. Let F be the generic fibre of c_X : it is a special curve, hence rational or elliptic.

If F is rational then $C(X) = R(X)$, by the fact that r_X dominates c_X (see 3.23). Thus $R(X)$ is $(n - 1)$ -dimensional. Moreover, we get from 3.5 that $R(X)$ is Moishezon, and so also X is Moishezon, by [C85].

Because $C(R(X)) = C(X)$, by Theorem 3.25 (which applies because X is Moishezon), we conclude that $c_{R(X)} = id_{R(X)}$, and so by the preceding Theorem 5.5, we infer that $R(X)$ is of general type, as claimed.

Next, if F is elliptic, then $c_X = J_X$, because J_X dominates c_X , and is obviously dominated by c_X if $\kappa(X) \geq 0$. We show this by induction on $n \geq 2$ (the case of curves is obvious).

Thus X is non-special, by assumption. Let $f : X \dashrightarrow Y$ be a fibration of general type, and let F be its general fibre. By 2.11, we have a factorisation $f = c_X \circ g$, for a certain fibration $g : C(X) \dashrightarrow Y$. Moreover, because of the functoriality properties of the core, the restriction of c_X to the general fibre X_y of f is the core of X_y . The induction hypothesis thus applies to X_y , and we get that

$\kappa(X_y) = (d - 1)$ if $d := \dim(X_y)$, and also that J_{X_y} (which is the restriction of c_X to X_y) is a fibration of general type.

We are thus in position of applying Theorem 4.4, since f is of general type. We get first that $\kappa(X) = \kappa(X_y) + \dim(Y) = (n - 1)$.

Then, because the general fibres X_y of f have nonnegative Kodaira dimension, we get also from Theorem 4.4 that J_X is of general type, since so is its restriction to the general fibre of f $\square$

The next step, classification of cases when $\text{ess}(X) = (n - 2)$ is similar (but some cases would require the general case of C_{gt}^{orb}).

5.4. The Decomposition Theorem. This is the following assertion, which motivates most of the present paper:

Theorem 5.8. *Let X be non special. Then c_X is a fibration of general type.*

Roughly speaking, this means that X “decomposes” into its “special part” (the fibres of c_X) and its “core” or “essential part”, the orbifold $(C(X)/\Delta(c_X))$, which is either a point, or of general type. Hence the name.

The Decomposition theorem can also be restated in the following form:

Any $X \in \mathcal{C}$ has a fibration both special and of general type. This fibration is unique, and it is the core of X . (Apply 3.8 to see the last two assertions).

Remark 5.9. *We already showed some cases of the preceding theorem 5.8:*

- (1) Up to dimension 3 as a byproduct of the description of the core given in low dimensions.
- (2) When X is of general type, and when $\text{ess}(X) = \dim(X)$: in this last case, showing that X is of general type is actually the same as showing the Decomposition theorem in this case. The crucial step was actually applying C_{gt}^{orb} .
- This is not an accident: we shall show right below that the Decomposition Conjecture is in fact a consequence of C_{gt}^{orb} .
- (3) Exactly the same remarks apply to the classification of the cases in which $\text{ess}(X) = (n - 1)$ or $(n - 2)$.
- (4) We shall show below two consequences of the Decomposition theorem: $C(X)$ is Moishezon, and meromorphic multisections of c_X are of general type.

Theorem 5.10. *Let $X \in \mathcal{C}$, and let $a_X : X \rightarrow \text{Alg}(X)$ be its algebraic reduction.*

There exists a factorisation $\phi : \text{Alg}(X) \rightarrow C(X)$ of $c_X = \phi \circ a_X$. Alternatively: $C(X)$ is Moishezon.

Proof: By Theorem 2.3, the general fibre of a_X is special $\square$

Let us now establish the following weak version of the Decomposition theorem (2.7 shows that it is a consequence of this result):

Proposition 5.11. *Let $j : Z \dashrightarrow X$ be a meromorphic map such that $c_X \circ j : Z \dashrightarrow C(X)$ is surjective and generically finite, $c_X : X \rightarrow C(X)$ being, as usual, the core of $X \in \mathcal{C}$.*

Then Z is a variety of general type.

Proof: By the functoriality of the core, and the surjectivity of $c_X \circ j$, we get the existence of a factorisation $c_j : C(Z) \dashrightarrow C(X)$. Because $c_X \circ j = c_j \circ c_Z$ is generically finite, so is c_Z . The claim thus follows from 5.5 $\square$

Proof of Theorem 5.8: Proceed by induction on $d := \text{ess}(X)$. When $d = 0$, the result trivially holds.

So assume it does when $\text{ess}(F) < d$. Assume that $\text{ess}(X) = d$, and that $\kappa(C(X)) \geq 0$.

By assumption ($d > 0$), X is not special. So let $f : X \dashrightarrow Y$ be of general type. We have, by Theorem 2.11 a factorisation $\psi : C(X) \dashrightarrow Y$ of $f = \psi \circ c_X$.

By the characteristic property of the core, the restriction of c_X to the general fibre X_y of f is the core of X_y (Otherwise X would contain special subvarieties strictly larger than the generic fibre of c_X , a contradiction).

By the induction hypothesis, we conclude that the restriction of c_X to the general fibre X_y of f is a fibration of general type.

We conclude from Theorem 4.4 that c_X itself is a fibration of general type, as claimed $\square$

5.5. Finite étale Covers. Let us indicate that the Decomposition theorem implies the invariance of $\text{ess}(X)$ under finite étale covers:

Theorem 5.12. *Let $u : X' \rightarrow X$ be a finite étale cover. Let $c_u : C(X') \rightarrow C(X)$ be the induced map. Then c_u is generically finite. In particular: $\text{ess}(X)$ is invariant under finite étale covers; and a finite étale cover of a special manifold is special.*

Remark that the proof of this seemingly easy statement requires here fairly deep tools. It were interesting to know if there is an easy proof of it.

Proof: We can assume that u is Galois, of group G . Due to its uniqueness, the map $c_{X'}$ is G -equivariant for an appropriate action of G on $C(X')$. Let $h : C(X') \rightarrow C'(X)$ be the G -quotient. We have natural maps $c'_X : X \rightarrow C'(X)$ (by G -invariance), and $v : C'(X) \rightarrow C(X)$ (since the fibres of c'_X , as images of those of $c_{X'}$ by u are special).

Moreover: $c_X = v \circ c'_X$, by the general properties of c_X .

By the Decomposition theorem, $c_{X'}$ is a fibration of general type. And so c'_X is also of general type, by 1.7. From 2.11, we infer the existence of a factorisation $w : C(X) \rightarrow C'(X)$ such that $c'_X = w \circ c_X$. Thus $C'(X) = C(X)$, and $h = c_u$ is a finite map $\square$

5.6. Essential and Bogomolov Dimensions.

Definition 5.13. *For $X \in \mathcal{C}$, let $B(X) := \max \{p > 0, \text{ such that there exists a Bogomolov sheaf } F \subset \Omega_X^p\}$; if there is no Bogomolov sheaf on X , define $B(X) := 0$.*

From 2.26, we see that $B(X) = 0$ if and only if $\text{ess}(X) = 0$. In general, we have:

Corollary 5.14. *For any $X \in \mathcal{C}$, we have: $\text{ess}(X) = B(X)$.*

Proof: Let F be any Bogomolov sheaf of dimension $p > 0$ on X . The associated fibration is thus of general type, and so dominated by c_X . Thus $\text{ess}(X) \geq p$

Conversely, because c_X is a fibration of general type, the Bogomolov sheaf associated to it has $p = \text{ess}(X)$. Hence the equality $\square$

Let us say that if F, G are Bogomolov sheaves on X that F **dominates** G if the fibration f (of general type) associated to F dominates the fibration g (of general type) associated to G .

Remark 5.15. *The Decomposition theorem may then be restated in the following form:*

For any $X \in \mathcal{C}$, $\text{ess}(X) = B(X)$, there exists on X a unique maximum Bogomolov sheaf, and it is defined by the core of X .

5.7. Fibre Products. We answer here the question 2.33

Theorem 5.16. *Let $X \in \mathcal{C}$, and $f : X \dashrightarrow Y, g : X \dashrightarrow Z$ be two fibrations of general type. Let $h : X \dashrightarrow W$ be the connected part of the Stein factorisation of the product map $f \times g : X \dashrightarrow Y \times Z$. Then h is a fibration of general type.*

Proof: This is an immediate generalisation of the argument in the classical case (when Y, Z are themselves of general type). Indeed: let $g' : W \dashrightarrow Z$ and $f' : W \dashrightarrow Y$ the fibrations such that $f = f' \circ h$ and $g = g' \circ h$.

By the equality (deduced from 4.4): $\kappa(W, h) = \kappa(W_z, g'_z) + \kappa(Z, g = g' \circ h) = \kappa(W_z, g'_z) + \dim(Z)$, we just need to check that the restriction of $g' : X_z \dashrightarrow W_z$ is of general type, for z general in Z . Observe the statement is bimeromorphically invariant. We thus assume that Z is projective (it is certainly Moishezon, since g is of general type). Take $S \subset Z$ be an intersection of generic hyperplane sections such that $(g')^{-1}(S) := T \subset W$ is mapped surjectively and generically finitely onto Y by f' .

Let $V := h^{-1}(T) \subset X$. The restriction $f_V : V \dashrightarrow Y$ of f to V has thus the restriction $h_V : V \dashrightarrow T$ of h to V as Stein factorisation. It is thus a fibration of general type, by 2.6.

If $g' : T \dashrightarrow S$ has positive-dimensional fibres, the conclusion follows from 2.9. If not, this means that g dominates f , in which case the assertion is trivial $\square$

5.8. Construction of the Core as the Highest General Type Fibration. We sketch here a second, shorter, construction of the core. It is more abstract, avoiding the consideration of chains of special varieties. It is something like a reconstruction “a posteriori”, once one already knows from previous more down-to-earth work, the existence of the core.

Theorem 5.17. *Let $X \in \mathcal{C}$. Then X admits a fibration both of general type, and special. This fibration is unique up to equivalence. It is the core of X .*

Proof: Everything is clear from 2.11, once the existence is known. We know from 5.16 above, together with 2.30 that X admits a unique fibration of general type $gt_X : X \dashrightarrow GT(X)$ which dominates all general type fibrations with domain X .

It remains to show that the general fibres of gt_X are special. Assume not. From the construction of the relative gt -reduction in 2.34, we deduce the existence of a factorisation $gt_X = h \circ g$, with $g : X \dashrightarrow G$ and $h : G \dashrightarrow GT(X)$ such that the restriction $g_a : X_a \dashrightarrow G_a$ to the general fibre X_a , $a \in GT(X)$, is the gt -reduction of X_a . Thus g_a is a fibration of general type, since X_a is assumed to be nonspecial.

But then, we get from theorem 5.8 that $g : X \dashrightarrow G$ is a fibration of general type. This contradicts the facts that $\dim(G) > \dim(GT(X))$, by hypothesis, and that gt_X dominates any fibration of general type defined on X $\square$

6. Geometric Conjectures

We shall state in this section some conjectures or questions naturally arising from our previous considerations.

6.1. Orbifold Rational Connectedness. We consider in this section an orbifold (Y/D) with $Y \in \mathcal{C}$ compact smooth and D an orbifold divisor of normal crossings on Y . We define several variants of rational connectedness, directly adapted from the non-orbifold case.

If $g : (Y/D) \rightarrow Z$ is a fibration, we define its orbifold base $(Z/\Delta(g, D))$ as in section 1.6. If g is only meromorphic, we replace (Y/D) by a terminal modification $\mu : (Y'/D') \rightarrow (Y/D)$ such that $g' := g \circ \mu$ is holomorphic, and then define the orbifold base of g to be that of g' . We then define $\kappa(Z, g, D)$ as the minimum of the Kodaira dimensions of the orbifold bases so obtained.

The orbifold general fibre of $g : (Y/D) \rightarrow Z$ is defined as the orbifold (Y_z/D_z) , with Y_z the general fibre of g , and $D_z := i^*(D)$, where $i : Y_z \rightarrow Y$ is the inclusion map.

An **orbifold rational curve** on (Y/D) is an irreducible rational curve C on Y such that $\kappa(\hat{C}, K_{\hat{C}} + n^*(D)) = -\infty$, where $n : \hat{C} \rightarrow C$ is the normalisation map.

Definition 6.1. *We say that (Y/D) is:*

- (1) **uniruled** if it is covered by a family of curves, whose generic member is an orbifold rational curve.
- (2) **rationally connected** if two generic points of Y can be joined by a chain of orbifold rational curves.
- (3) **rationally generated** if, for any fibration $g : (Y/D) \dashrightarrow Z$, the orbifold base of some holomorphic model of g is uniruled.
- (4) **κ -rationally connected** if, for any fibration $g : (Y/D) \dashrightarrow Z$, $\kappa(Z, g, D) = -\infty$.

We note, as usual, for short: R.C (resp. R.G; resp. k-RG) for rationally connected (resp. rationally generated; resp. κ -rationally generated). Notice that R.G was introduced in [C95] in the non-orbifold context, and that the notion of κ -rational generatedness is directly inspired from the definition of κ^+ in the same paper. Actually, one can even define:

Definition 6.2. *For (Y/D) as above, define: $\kappa_+(Y/D) := \max\{\kappa(Z/\Delta(g, D))\}$, where g runs through all fibrations $g : (Y/D) \dashrightarrow Z$ as above.*

Then κ -rational generatedness is defined by the equality: $\kappa_+ = -\infty$.

Remark 6.3. *One can also introduce in the orbifold case many other invariants, based on the vanishing of holomorphic “covariant” tensors, which conjecturally characterise rational connectedness. We do not do this here, because one needs first to define these notions in the orbifold case. So we introduced only the simplest one: κ_+ .*

One could, for example, define $\kappa_+(Y/D) := \max\{\kappa(Y, F)\}$, where F ranges over all rank one coherent subsheaves of $\Omega_Y(\log(D))$. (See section 2.6 for this notion). Another variant of orbifold rational connectedness would be defined by $\kappa_+(Y/D) = -\infty$.

Standard arguments that we do not need to reproduce easily show:

Proposition 6.4. *For any orbifold (Y/D) as above, one has:*

- (1) $\kappa(Y/D) = -\infty$ if (Y/D) is uniruled.
- (2) (Y/D) is R.G if it is R.C and is k-RG if it is R.G.

The reverse implications depend on the answers to the following two questions.

Question 6.5. (1) Assume $\kappa(Y/D) = -\infty$. Is then (Y/D) uniruled?

(2) Let $g : (Y/D) \rightarrow \mathbb{P}^1$ be a fibration with generic orbifold fibre rationally connected, and such that $\kappa(\mathbb{P}^1/\Delta(g, D)) = -\infty$. Does g have a section, that is: an orbifold rational curve mapped isomorphically to $\mathbb{P}^1$ by g ?

Of course, the second question is directly inspired from [G-H-S]. Again, standard arguments show that:

If question (1) has a positive answer, then: R.G and k-R.G are equivalent.

If question (2) has a positive answer, then: R.C and R.G are equivalent properties.

Remark 6.6. Question (1) seems quite delicate (if true): it is open in the non-orbifold case. It is open even in the surface case ($\dim(Y) = 2$), and seems to present great subtlety. It is similar to the case handled in [K-MK], of log-Del Pezzo surfaces with a reduced boundary D . But their methods might apply to treat the above surface case.

Remark 6.7. Another natural question in this context is whether the “glueing Lemma” of [Ko-Mi-Mo] extends to this context: for example, can one add to an orbifold rational curve C' in (Y/D) sufficiently many “free” orbifold rational curves if (Y/D) is uniruled, in such a way that the union deforms to an (irreducible) orbifold rational curve?

We shall now justify the introduction of the notion of κ -generatedness, by showing that it permits to define the notion of rational quotient (even in the orbifold case) without solving question (1) above, this definition however depends on the solution of conjecture $C_{n,m}^{orb}$.

Conjecture 6.8. ($(\kappa = -\infty)$ -Conjecture) Assume that $\kappa(Y/D) = -\infty$.

There exists then a unique fibration $r_{Y/D} : Y \dashrightarrow R(Y/D)$ such that:

(1) the general orbifold fibre $F_r := (Y_r/D_r)$, $r \in R(Y/D)$ of $r_{Y/D}$ has $\kappa_+(Y_r/D_r) = -\infty$.

(2) $\kappa(R(Y/D), r_{Y/D}, D) \geq 0$.

Proposition 6.9. A fibration as in 6.8 is unique if it exists. The existence follows from Conjecture $C_{n,m}^{orb}$.

Proof: Assume there exists another fibration $h : Y \dashrightarrow Z$ with the same properties. Let $u : Y \dashrightarrow W$ be the Stein factorised product of $r_{Y/D}$ and h . Assume h is not equivalent to $r_{Y/D}$. The restriction $h_r : Y_r \dashrightarrow W_r$ then maps the orbifold F_r to a positive-dimensional orbifold of nonnegative Kodaira dimension (by the property (2), assumed to hold for h). This contradicts (1) above for $r_{Y/D}$, and shows unicity.

For the existence: consider the Stein factorised product $p : Y \dashrightarrow V$ of all fibrations $h : Y \dashrightarrow Z$ such that $\kappa(Z/\Delta(h, D)) \geq 0$. By $C_{n,m}^{orb}$, p itself has this property. Assume the general orbifold fibre of p were not κ -RG. We could find (by the result on the relative quotients) fibrations $h : Y \dashrightarrow U$ and $k : U \dashrightarrow V$ such that $p = k \circ h$, and such that the general orbifold fibre of k had nonnegative Kodaira dimension. By $C_{n,m}^{orb}$ again, this would imply that $\kappa(U/\Delta(h, D)) \geq 0$.

But this contradicts the maximality of p among such fibrations, since $\dim(U) > \dim(V)$ $\square$

6.2. Orbifold Iitaka-Moishezon Fibration. The Iitaka-Moishezon fibration of an orbifold (Y/D) as in section 6.1 above can be defined without difficulty by considering the linear system $|m.(K_{Y/D})|$ for m sufficiently big and divisible.

We shall then denote by $J_{Y/D} : (Y/D) \dashrightarrow J(Y/D)$ the resulting Iitaka-Moishezon fibration.

By the usual properties of Iitaka fibrations, one has: $\kappa(Y_s/D_s) = 0$, for its general orbifold fibre.

6.3. Orbifold Rational quotient and Iitaka Fibration of a Special fibration. We describe in this section an alternative construction of the core of an arbitrary X , along the program described in the introduction. This construction actually coincides in dimensions 2 and 3 with the construction of the core given in 3.30 and 3.37.

The idea is quite simple: assume we have a **special** fibration $f : X \dashrightarrow Y$. We try to understand the obstructions to constructing a non-trivial $h : Y \dashrightarrow Z$ such that $g := h \circ f : X \dashrightarrow Z$ is still special, this in terms of the invariant $\kappa(Y, f)$.

This will certainly be impossible if f is of general type, because then f is the core of X , by 3.8. The other cases can actually be reduced (as in the classical Iitaka-Moishezon classification Program) to the cases $\kappa = 0$, or $-\infty$. And we conjecture below that such a non-trivial g then exists.

This leads to a conjectural decomposition of the core in a canonical and functorial sequence of orbifold rational and Iitaka fibrations.

First, we need to give an orbifold generalisation of Iitaka fibrations and rational quotients for special fibrations.

We consider in the sequel a **special** fibration $f : X \dashrightarrow Y$ (thus with $X \in \mathcal{C}$).

Theorem 6.10. *Assume that $\kappa(Y, f) = 0$. Then X is special.*

Proof: This follows from 4.4 and its corollaries. Assume indeed X were not special. By 2.11, there would exist a fibration $g : Y \dashrightarrow Z$ such that $g \circ f : X \dashrightarrow Z$ were of general type.

But then, we would have:

- (1) $\kappa(Y_z, f_z) \geq 0$, because $\kappa(Y, f) = 0$, by the easy addition theorem.
- (2) $0 = \kappa(Y, f) = \kappa(Y_z, f_z) + \dim(Z) \geq \dim(Z) > 0$, because of (1). Contradiction $\square$

From this result, we deduce immediately the notion of **Iitaka fibration** of the special fibration $f : X \dashrightarrow Y$: it is just the fibration $J_f : Y \rightarrow J(f)$ associated to the $\mathbb{Q}$ -divisor $(K_Y + \Delta(f))$, first replacing f by any of its admissible models. Alternatively, it is also the Iitaka-Moishezon fibration of the orbifold base $(Y/\Delta(f))$ of any of its admissible prepared models.

The important property is that the general orbifold fibre of J_f has Kodaira dimension zero (by the remarks made in section 6.2 above).

From this follows immediately, using 6.10, that the composite fibration $J(f) \circ f : X \dashrightarrow J(f)$ is special.

Next we need to investigate the case when $\kappa(Y, f) = -\infty$. Using the preceding conjecture 6.8, we get:

Conjecture 6.11. *Let $f : X \dashrightarrow Y$ be special fibration, with $\kappa(Y, f) = -\infty$. There exists a unique fibration $r_f : (Y/\Delta(f)) \dashrightarrow R(f)$ such that:*

- (1) *The general orbifold fibre of r_f is κ -rationally generated.*
- (2) $\kappa(R(f)/\Delta(r_f, \Delta(f))) = \kappa(R(f), r_f \circ f) \geq 0$.

*We call r_f the **rational quotient** of f .*

Notice that, assuming the existence of r_f , we have: $\dim(R(f)) < \dim(Y)$, and that $f^{-1}(Y_r) := X_r$ is then special, for $r \in R(f)$ general.

By a slight extension of the definition, we also define the rational quotient r_f of f to be the identity map of Y if $\kappa(Y, f) \geq 0$.

Caution: The Iitaka base fibration (resp. the rational base quotient) differ from the (relative) Iitaka fibration (resp. rational quotient) of f , which are defined for any fibration f , and provide a factorisation of f , inducing on the general fibre of f the corresponding fibration of that fibre. Of course, if the questions 6.5 have a positive answer, the general orbifold fibre of r_f is rationally connected.

6.4. The Core as an Iterated Orbifold Rational Quotients and Iitaka Fibrations. We assume in this section the conjecture 6.8.

Definition 6.12. Special Reduction of a Special Fibration *Let $f : X \rightarrow Y$ be a special fibration. Assume that the above conjecture 6.8 hold. Define $s_f : J_{r_f} \circ r_f : Y \dashrightarrow J(r_f \circ f) := S(f)$, and call it the first special reduction of f .*

In vague terms: s_f is obtained by first applying r_f to Y , and then $J(r_f \circ f)$ to $(R(f)/\Delta(r_f \circ f))$. Notice that $s_f : X \rightarrow S(f)$ has still its general fibre special. So the construction can be iterated, and we get a sequence of special fibrations: $s_f^0 := id_Y$, with $Y := S^0(f)$, and then inductively: $s_f^{k+1} := s_{s_f^k} : X \rightarrow S^{k+1}(f)$. We call s_f^k the k^{th} -special reduction of f .

Notice that $s_f = f$ exactly when $\kappa(Y, f) = \dim(Y) \geq 0$.

In this case, by 3.8, f is the core of X . And it is the constant map iff $\kappa(Y, f) = \dim(Y) = 0$.

From this we immediately deduce the following, by applying the above iteration starting with $f := id_X$, the identity map of X , thus getting the sequence of k^{th} -special reductions of X :

Theorem 6.13. *Assume that the conjecture 6.8 holds. Let $X \in \mathcal{C}$, with $n := \dim(X)$. Define $s_X^k := s_{id_X}^k : X \rightarrow S^k(X) := S^k(id_X), \forall k \geq 0$. Then $s_X^n = c_X$, the core of X , and c_X is thus either constant, or a fibration of general type.*

Of course, in general, the sequence of fibrations s_X^k will be stationary before its n^{th} -term is reached.

Definition 6.14. *For $X \in \mathcal{C}$ and $k \geq 0$, define: $s^k(X) := \dim(S^k(X))$, and $rs^k(X) := \dim(R(s_X^k))$, where $r_{s_X^k} : S^k(X) \rightarrow R(s_X^k)$ is the rational quotient of $s_X^k : X \rightarrow S^k(X)$, the k^{th} -special reduction of X .*

Under the said conjecture, these are new bimeromorphic invariants of X . We call $s^k(X)$ the k^{th} -special dimension of X , and $rs^k(X)$ the reduced k^{th} -special dimension of X .

Standard easy arguments give:

Proposition 6.15. *Assume that the conjecture 6.8 holds. The invariants $s^k(X)$ and $rs^k(X)$ are invariant under finite étale covers $u : X' \rightarrow X$ of X . The same holds for the associated fibrations: $s_{X'}^k$ (resp. $rs_{X'}^k$) is the Stein factorisation of $s_X^k \circ u$ (resp. $rs_X^k \circ u$).*

One has the following “strictness” property for the core decomposition, which we quote without proof:

Theorem 6.16. *Assume that conjecture 6.8 holds. Let $X \in \mathcal{C}$, and let $k \geq j \geq 0$ be integers. Let S (resp. R) be the general fibre of s_X^k (resp. rs_X^k). The restrictions of s_X^j and rs_X^j to S and R coincide respectively with s_S^j , rs_S^j , s_R^j and rs_R^j .*

Remark 6.17. *The sequence (s) of $(2n + 2)$ integers $\dim(X) = n = s^0 \geq rs^0 \geq s^1 \geq rs^1 \geq \dots s^k \geq rs^k \geq \dots s^n \geq rs^n$ partitions the class of n -dimensional manifolds in $\mathcal{C}$ into a certain (finite) number $c(n)$ of classes, according to the number of steps and the dimensions of the steps needed to decompose the core in orbifold rational quotients and Iitaka fibrations.*

One has for example: $c(0) = 1$, $c(1) = 3$, $c(2) = 8$, $c(3) = 21$, by direct listing.

One can easily compute the numbers $c(n)$ as follows. Say a sequence (s) is special if its last term is zero.

Notice the sequence is stationary precisely if two consecutive terms $rs^k = s^{k+1}$ coincide, or if one is zero.

We say the sequence (s) **ends** at J (resp. r) if $s^k = rs^k < rs^{k-1}$ (resp. $s^{k+1} = rs^k < s^k$).

let $s(n)$ be the number of special sequences of n -dimensional manifolds.

Let $s'(n)$ (resp. $s''(n)$) be the number of such special sequences which end at r (resp. J).

Thus: $s(n) = s'(n) + s''(n)$.

We quote without proof:

Lemma 6.18. *One has, for any $n \geq 0$:*

- (1) $c(n) := \sum_{m=0,1,\dots,n} s(m)$,
- (2) $s'(n) = 1 + \sum_{h=1,\dots,(n-1)} s''(h)$,
- (3) $s''(n) = 1 + \sum_{h=1,2,\dots,(n-1)} s(h)$.

Because $s(0) = 1, s'(1) = s''(1) = 1, s(1) = 2, c(1) = 3$, we get inductively: $s'(2) = 2, s''(2) = 3, s(2) = 5, c(2) = 8$, for example. We already met these classes, actually, when listing the possibilities for the core of a surface.

Corollary 6.19. *For all $n \geq 0$, one has:*

- (1) $s'(n+1) = s(n)$, $s''(n+1) = 2s(n) - s(n-1)$,
- (2) $s(n+1) = 3s(n) - s(n-1)$, $c(n+1) = c(n) + s(n+1)$.

And, finally:

Corollary 6.20. *If $\alpha := (3 + \sqrt{5})/2$, and $\beta := (3 - \sqrt{5})/2$, one has, for all $n \geq 0$:*

- (1) $s(n) = ((1 + \sqrt{5})/(2\sqrt{5})) \cdot \alpha^n + ((\sqrt{5} - 1)/(2\sqrt{5})) \cdot \beta^n$.
- (2) $c(n) = (\alpha^{n+1} - \beta^{n+1})/\sqrt{5}$.

6.5. Deformation invariance. We next state the following, for $X \in \mathcal{C}$ smooth:

Conjecture 6.21. *(Deformation Conjecture) $\text{ess}(X)$ is a deformation invariant of X . In particular: the class of special manifolds is stable under deformation.*

One may ask more precise versions: does the core of X vary holomorphically with X for suitable models?

Even weaker versions are difficult: is $\text{ess}(X)$ upper or lower semi-continuous under smooth deformations?

Is it lower semi-continuous under degenerations?

The following is important and maybe accessible: is a degeneration of special manifolds still special (in the sense that all of its irreducible components are special)?

A more refined version of the above deformation conjecture is the following, which extends the classical conjecture concerning the deformation invariance of the Kodaira dimension for compact Kähler manifolds:

Question 6.22. *Assume that the $(\kappa = -\infty)$ -Conjecture hold. Let $X \in \mathcal{C}$. Are the invariants $s^k(X)$ and $rs^k(X)$ deformation invariants of X ? More precisely, do the associated fibrations vary holomorphically (for suitable models) when deforming X ?*

Notice that this is the case for $k = 0$, and in the projective case (and deformation among projective manifolds) for s_X^1 , by the invariance of plurigena, due to Y.T.Siu ([Siu']).

A slightly less general version, but which deals with invariants known to exist, is the following, concerning the higher Kodaira dimensions, defined in 3.6:

Question 6.23. *Let X be compact Kähler. Assume the invariants:*

$\kappa(X) \geq \kappa'(X) \geq \dots \geq \kappa^r(X) \geq -\infty$ are defined.

Are then these invariants defined also and constant for small deformations of X ?

Of course, one could still extend the preceding questions to the category of orbifolds, appropriately defined.

6.6. Weakly-Special Manifolds.

Definition 6.24. *$X \in \mathcal{C}$ is said to be **weakly-special** (or *w-special* for short), if there is no pair (u, f') in which $u : X' \rightarrow X$ is a finite étale covering, and $f' : X' \dashrightarrow Y'$ is a surjective meromorphic map onto a manifold Y' of general type (in the usual sense) and positive dimension.*

From 1.7, we get:

Proposition 6.25. *If X is special, it is w-special.*

Proof: It is the same as in 2.33. Assume by contradiction that a triple (u, X', f') as in the preceding definition exists. We can assume the cover u to be Galois, of group G . If f' is G -equivariant, then f' descends to $f : X \dashrightarrow Y := (Y'/G)$, and f is of general type by 1.7. Otherwise, just replace f' by the least upper bound f'' of the (finite) family $(f' \circ g), g \in G$. Now f'' is G -equivariant, and maps X' to Y'' , which is of general type, by 2.29 $\square$

Example 6.26. *All the examples listed in 2.2 are thus w-special.*

Notice that, contrary to the case of special manifolds, it is obvious that any finite étale cover of a w-special manifold is again w-special. In particular, any finite étale cover of a special manifold is w-special.

Conversely, it follows from 3.32 that special surfaces and are weakly special.

The same assertion for threefolds is expected to have a negative answer. More generally:

Question 6.27. *Does there exist manifolds which are w-special, but not special?*

Although it seems reasonable to expect a positive answer, the construction of such examples is far from being immediate.

The following has been shown in 5.3:

Proposition 6.28. *Let X be w -special. The Albanese map of X is then surjective and connected. In particular: if X is special, and X' is a finite étale cover of X , the Albanese map of X' is surjective and connected.*

Observe that, in contrast to the case X special, this result does not assert that the Albanese map of X is multiplicity free if X is w -special.

Thus if X is w -special, but its Albanese map has multiple fibres in codimension one, X cannot be special.

The core as a weak analogue, described below.

6.7. The Weak Version of the Core. It is possible to construct an analogue of the core for weakly-special manifolds and fibrations strongly of general type. We briefly indicate how to do this. The proofs are much easier than in the case of the actual core, and will be omitted, because of being straightforward applications of standard results concerning the classical (not orbifold) Kodaira dimension. These notions will not be used, anyway, but it seems worth mentioning them, for sake of completeness.

Definition 6.29. *We say that the fibration $f : X \dashrightarrow Y$, with $Y \in \mathcal{C}$, is of **strong general type** if there exists a finite étale cover $u : X' \rightarrow X$ such that Y' is of general type, where $f \circ u = v \circ f'$ is the Stein factorisation of $f \circ u$. Here $f' : X' \dashrightarrow Y'$ is connected and $v : Y' \rightarrow Y$ is finite. Thus X is weakly-special iff it has no fibration $f : X \rightarrow Y$ of strong general type.*

Notice that, by 1.7, a fibration of strong general type is of general type.

Proposition 6.30. *Let $f : X \rightarrow Y$ and $g : X \rightarrow Z$ be two fibrations of strong general type. Let $h : X \rightarrow V$ be the Stein factorisation of the product map $f \times g : X \rightarrow W \subset Y \times Z$. Then h is of strong general type.*

Corollary 6.31. *For $X \in \mathcal{C}$, there exists a unique fibration $gt_X^+ : X \rightarrow GT^+(X)$ which is of strong general type and dominates (in the obvious sense) any fibration of strong general type $f : X \rightarrow Y$. We call it the strong general type reduction of X . It is thus the constant map iff X is weakly-special.*

Corollary 6.32. *For $X \in \mathcal{C}$, there is a unique $w_X : C(X) \rightarrow GT^+(X)$ such that $gt_X^+ = w_X \circ c_X$, if $c_X : X \rightarrow C(X)$ is the core of X .*

Proof: By 2.11, if $f : X \rightarrow Y$ is a fibration of general type, there is a unique factorisation $\theta : C(X) \rightarrow Y$ such that $f = \theta \circ c_X$. Apply this to $f = gt_X^+$, which is of general type, since of strong general type.

Remark 6.33.

- (1) The general type reduction $gt_X : X \rightarrow GT(X)$ of X (see 2.30) dominates gt_X^+ (and is dominated by c_X).
- (2) For any fibration $f : X \rightarrow Y$, it is possible to construct a relative strong general type reduction of f , as in the general type case.
- (3) Applying the above reduction to $f := gt_X^+$, we would get that the general fibre of gt_X^+ is weakly-special, provided one could show the following : if $g \circ f : X \rightarrow Z$ is of strong general type,

and if so is the restriction $f_z : X_z \rightarrow Y_z$ of $f : X \rightarrow Y$ to the general fibre X_z of $g \circ f$, then $f : X \rightarrow Y$ is of strong general type.

This statement seems however very unpalatable (it would be much better to have counterexamples!), which is why we were led to consider the more complicated notion of special varieties and general type fibrations.

(4) As D.Abramovitch informed me, in [Ab] a reduction map analogous to the preceding gt_X^+ was constructed, but with fibrations onto general type varieties in place of strong general type fibrations (ie: ignoring étale covers), inspired by “Harris conjecture”. It enjoys nice properties, but is very unstable with respect to finite étale covers (which however preserve Kobayashi pseudo-metric and the arithmetic behaviour).

6.8. Classical gcd -Multiplicities. We can still introduce a third notion of special manifolds, which interpolates between the preceding two ones. It is based on the classical definition of multiple fibres, using (gcd) instead of (inf) . We shall be brief on this.

Definition 6.34. *Let $f : X \rightarrow Y$ be a fibration, and let $\Delta \subset Y$ be an irreducible reduced divisor. Let $f^*(\Delta) = (\sum_{j \in J} m_j \cdot D_j) + R$, where $f(D_j) = \Delta, \forall j$, while $f(R)$ has codimension two or more in Y . Define multiplicities as follows:*

$m^-(\Delta, f) := gcd(m_j, j \in J)$, and $m(\Delta, f) := min(m_j, j \in J)$. Obviously, m^- divides m .

Now using this definition of multiplicity, one can define, exactly as we did, with the same proofs, the notion of base orbifold and Kodaira dimension of a fibration. This leads to the $(gcd$ -versions) of fibration of general type, and special manifold. The gcd -core can be constructed also with the same properties.

One property only is (possibly) lost: gcd -fibrations of general type are no longer naturally in bijective correspondance with Bogomolov sheaves.

On the other hand, the additivity result 4.4 and its proof (even becoming slightly simpler at a point) remain. So it may happen that the gcd -core is still a fibration of gcd -general type.

At this point, one may wonder:

- (1) Are the two theories actually different?
- (2) If yes, which one is the “right” one?

Presently, there is no counterexample to distinguish them. But it seems plausible that the two theories differ. We shall make some general easy comparisons between them right below.

Concerning the second question (b) above, assuming a positive answer to (a), it seems that the version given here (with inf , not gcd) is the right one. This is supported by the correspondance with Bogomolov sheaves, and the direct link between inf fibrations of general type and geometric positivity properties of cotangent sheaves.

This link between geometry and positivity is the main theme of Classification theory.

Other impressions support this choice of inf versus gcd : the inf -notion is better suited for the study of Kobayashi pseudo-metric. Also, one may expect on the arithmetic side the fact that gcd -special manifolds are not always potentially dense.

The only feature in favour of the gcd -theory is that it is better suited for the computation of the fundamental group. This property cannot however compensate the other drawbacks.

Let us end up with some obvious remarks concerning the comparisons between the three notions of specialness introduced:

A special manifold is *gcd*-special (because a fibration of general type is of *gcd*-general type).

A *gcd*-special manifold is *w*-special (by the *gcd*-version of 1.7).

Although no counterexamples are known, it seems plausible that the three notions are distinct.

The easiest counterexample would be a *w*-special, but not special threefold.

The following question is related to the comparison of special and *gcd*-special manifolds. A positive answer (not necessarily expected) would show the coincidence of the two notions):

Question 6.35. *Assume the general fibre F of f is special. Is it then true that $m(\Delta, f) = m^-(\Delta, f)$, for any pair $(\Delta \subset Y)$?*

Remark 6.36.

It is sufficient to answer the above question when Y is a curve and $m^-(\Delta, f) = 1$: one would like to know if one of the m'_j 's = 1, then. The question above is motivated by the fact that it has a positive answer when F is rationally connected (by [G-H-S], the above assertion being actually the main difficulty in the proof), and when F is an abelian variety (or a complex torus), by an easy argument (the addition of local multisections). The next interesting case to consider is that of F a *K3*-surface or a special surface with $\kappa = 1$. The answer does not seem to be known in these cases.

6.9. Irreducible Special Manifolds.

Definition 6.37. *Let $X \in \mathcal{C}$ be a special connected manifold. We say X is **irreducible special** (resp. **strongly irreducible special**) if it is special, and if every fibration $f : X \rightarrow Y$ (resp. covering family $((V_t)_{t \in T})$ of X) with general fibre special (resp. special general member V_t) is trivial (ie: has fibre or base dimension zero); (resp. if V_t has dimension or codimension zero in X).*

Examples: If X is rationally connected of dimension 2 or more, it is not strongly irreducible special.

But it is possible that some rationally connected threefolds (necessarily bimeromorphic to some terminal $\mathbb{Q}$ -Fano threefolds with Picard number one) may be irreducible special. The answer does not seem to be known.

The general quintic threefold is strongly irreducible (because not generically covered by images of elliptic curves or abelian surfaces).

Remark 6.38. *If X is irreducible special, then $\kappa(X) = 0$, or $-\infty$. (By 3.10, 5.1, and the Iitaka-Moishezon fibration).*

Recall the standard:

Uniruledness Conjecture: If $\kappa(X) = -\infty$, then X is uniruled.

Observe that it is obviously equivalent to:

Irreducibility Conjecture: If $\kappa(X) = -\infty$, and if X is special, of dimension $n \geq 2$, then X is not strongly irreducible special.

6.10. Manifolds with nef anticanonical bundle. Question: Let X be a Kähler manifold with $-K_X$ nef (see [D-P-S] for this notion). Is then X special? A positive answer to this question would imply most of the conjectures concerning these manifolds.

Certain important consequences of a positive answer to this question have been established: Q.Zhang has shown in [Zh] the surjectivity of the Albanese map in the projective case.

And M.Paun has shown (in the difficult Kähler case, see [Pa]) the virtual nilpotency of the fundamental group. Combined with Q.Zhang's result and [C4], it shows that if X is projective and has nef anticanonical bundle, then it has virtually abelian fundamental group.

7. The Fundamental group

7.1. The Abelianity conjecture. Our considerations here are guided by the following:

Conjecture 7.1. *Let X be special. Then $\pi_1(X)$ is virtually abelian.*

Recall that a group is said to be **virtually abelian** (one also says: almost abelian) if it has a subgroup of finite index which is abelian.

This conjecture is motivated by the fact that it should be true for klt orbifolds either rationally connected or with $\kappa = 0$, and the fact that a special manifold should be (after section 6.4) a tower of orbifold fibrations with orbifold fibres in the preceding two classes. See section 9.4 for the motivation of a similar conjecture, concerning the Kobayashi pseudo-metric.

Examples 7.2.

(1) Curves: A curve is special iff its fundamental group is virtually abelian.

(2) Rationally connected manifolds: They have trivial fundamental group ([C3],[C5],[Ko-Mi-Mo]), and are special.

More generally, if r_X is the rational quotient of X (see 12.26), then it induces an isomorphism between the fundamental groups of X and $R(X)$ by [Ko]. So that X and $R(X)$ are simultaneously special and have simultaneously virtually abelian fundamental groups. The above conjecture should thus essentially reduce to the case where $\kappa(X) \geq 0$ if the uniruledness conjecture is true.

(3) Manifolds with $a(X) = 0$ or with $\kappa(X) = 0$ are special, and standard conjectures say they should have a virtually abelian fundamental group. These conjectures are thus special cases of conjecture I above.

Notice that Conjecture 7.1 is established for manifolds with $c_1(X) = 0$ (which are indeed special, and have almost abelian fundamental group, by the Calabi-Yau theorem).

This provides strong support for (and motivates) the case where $\kappa = 0$.

Let us give another example of result supporting Conjecture 7.1 in the case $a(X) = 0$:

Theorem 7.3. *([C95]) Let $X \in \mathcal{C}$ be such that: $a(X) = 0$ and $\chi(X, \mathcal{O}_X) \neq 0$. Then: $\pi_1(X)$ is finite (in particular: conjecture 7.1 holds for X).*

More generally, if F is the generic fibre of the algebraic reduction map of any $X \in \mathcal{C}$, with $\chi(X, \mathcal{O}_X) \neq 0$, the image in $\pi_1(X)$ of $\pi_1(F)$ is finite. (Recall that F is special, by 2.3).

(4) Surfaces and Threefolds: By 3.32 and 3.38, Conjecture 7.1 holds for Kähler surfaces and projective threefolds with $\kappa \neq 2$ (and more precisely, for all special threefolds, except maybe for the ones with $\kappa = 2$, and having a model of the Iitaka-Moishezon fibration with orbifold base a log-Enriques or log-Fano normal surface. In this case, the conjecture is open because of an insufficient knowledge of these orbifolds).

This case shows clearly that the solution of Conjecture 7.1 above should be extended to the orbifold situation. Recall that the fundamental group of (Y/Δ) is defined as the quotient of the fundamental group of the complement of Δ in Y , modulo the normal subgroup generated by elements of the form $\lambda_j^{m_j}$, where λ_j is a small loop winding once around Δ_j , if:

$$\Delta := \sum_{j \in J} (1 - 1/m_j) \cdot \Delta_j.$$

Question 7.4. *Let (Y, Δ) be a klt-orbifold, with Y normal. Assume that $-(K_Y + \Delta)$ is ample or torsion. Is then the fundamental group of (Y/Δ) trivial in the first case, and almost abelian in the second case?*

More generally is the fundamental group of (Y/Δ) almost abelian if $\kappa(Y/\Delta) = 0$, or if (Y/Δ) is κ -rationally generated (see section 6.1 for this notion)?

Notice that a positive answer for surfaces solves the remaining cases of the abelianity conjecture in dimension three.

(5) Manifolds with $-K_X$ nef are conjectured to have a virtually abelian fundamental group. This conjecture is thus also a consequence of conjecture 7.1 under a positive answer to the question asked above of whether such manifolds are special. The main results concerning this conjecture are in [D-P-S], [D-P-S'], [Pa] and [Zh].

(6) Kähler manifolds X covered by $\mathbb{C}^d$ are special (by the results of the next section 8.2, and S. Itaka conjectured them to be covered by complex tori. Easy arguments (given below) show that this happens precisely if $\pi_1(X)$ is virtually abelian. So this conjecture is also a special case of the abelianity conjecture above.

We are not presently able to solve substantial examples of the abelianity conjecture. But, at least, the solvable and linear quotients of the fundamental group can be dealt with.

7.2. Linear and Solvable Quotients. Recall first (5.3):

Theorem 7.5. *Let X be any finite étale cover of a special manifold.*

Let $\alpha_X : X \rightarrow \text{Alb}(X)$ be the Albanese map of X . Then it is surjective, connected. Moreover, α_X has no multiple fibres in codimension one if X is special.

Remark 7.6. *Due to 5.1, this extends and sharpens slightly a result of [Kw].*

Remark 7.7. *Assume that conjecture 7.1 holds (or alternatively, that X is special and has almost abelian fundamental group). Easy standard arguments then show, using the surjectivity of the Albanese map and its absence of multiple fibres in codimension one, that the universal cover of X is holomorphically convex (in accordance with the conjecture of Shafarevitch), and that its Remmert reduction makes it proper over an affine complex space, which is the universal cover of its Albanese variety.*

Now using [C4] (see also [A-N], [B], [Sim]), one gets:

Corollary 7.8. *Let $X \in \mathcal{C}$ be special, and let $c : X' \rightarrow X$ to be an étale finite cover. Then:*

- (1) *The derived group $D\pi_1(X)$ of X is finitely generated.*
- (2) *The Green-Lazarsfeld set $\Sigma^1(X')$ consists of finitely many torsion characters of $\pi_1(X)$.*

Again using [C4], corollary 4.2' (based on Deligne's semi-simplicity and invariant cycle theorems), and 7.5, we get:

Theorem 7.9. *If X is w -special and if $\mu : \pi_1(X) \rightarrow G$ is a surjective morphism of groups, with G solvable and torsionfree, then G is virtually abelian. (We say that torsionfree solvable quotients of $\pi_1(X)$ are almost abelian).*

Proof: This is an easy consequence of 7.5 (which says that the Albanese map of X is surjective and connected) and of corollary 4.2' of [C4] (which says that if the Albanese map of X' is surjective for any finite étale cover X' of X , then the conclusion of 7.9 is true for X) $\square$

Remark this applies in particular when $a(X) = 0$ (it is shown by a different method in [C4]), and when $\kappa(X) = 0$.

Now, using a result of [Z], itself based on results of N.Mok [Mk] (see also [Siu]), and 7.9, we get:

Theorem 7.10. *Let X be w -special, and let $\rho : \pi_1(X) \rightarrow \text{Gl}(N, \mathbb{C})$ be a linear representation. Then $G := \text{Image}(\rho)$ is virtually abelian. (We say that linear quotients of $\pi_1(X)$ are almost abelian).*

So, counter-examples to conjecture 7.1, if any, can't be detected by linear representations of their fundamental groups.

Proof: Let G' be the Zariski closure of G . Replacing X by a suitable étale cover, we assume that G' is connected. We have an exact sequence of groups:

$$1 \rightarrow R \rightarrow G' \rightarrow S \rightarrow 1,$$

with S semi-simple and R solvable. Consider the representation $\rho' = \sigma \circ \rho : \pi_1(X) \rightarrow S$, where $\sigma : G' \rightarrow S$ is the above quotient. By [Z], thm.5,p.105, we get that S is trivial (ie: reduced to the unit element). Thus $G \subset R$ is solvable, and can be assumed to be torsionfree, passing to a suitable finite étale cover, by A.Selberg's theorem. The conclusion now follows from 7.9 $\square$

Corollary 7.11. *Conjecture 7.1 holds when $\pi_1(X)$ is linear (ie: has a faithful representation ρ .)*

7.3. Manifolds Covered by $\mathbb{C}^n$. An easy consequence of 7.10 is:

Theorem 7.12. *Let $X \in \mathcal{C}$ be covered by $\mathbb{C}^d$, so $d := \dim(X)$. Assume that $\pi_1(X)$ is either solvable, or linear (ie: admits a faithful linear representation in some $\text{Gl}(N, \mathbb{C})$). Then X is covered by a complex torus. (ie: S.Itaka's Conjecture then holds for X).*

Proof: We know that X is special by 8.2 below (one can also get directly from [K-O] that X is w -special, which is sufficient for our purpose). Thus $\pi_1(X)$ is virtually abelian by 7.9 or 7.10 above.

Replacing X by a suitable unramified cover, the conclusion thus follows from the next Lemma 7.13 $\square$

Lemma 7.13. *Let $X \in \mathcal{C}$ be covered by $\mathbb{C}^d$, with $d := \dim(X)$. Assume that $\pi_1(X)$ is abelian. Then X is a complex torus.*

Proof: Let $\alpha : X \rightarrow A$ be the Albanese map of X . By 7.5, it is surjective and connected. Let F be a generic fibre of α .

It is sufficient to show that $\dim(F) = 0$, because X is covered by $\mathbb{C}^d$ (indeed, α is then birational. And so X contains a rational curve if α is not isomorphic. But this rational curve would lift after normalisation to the universal cover of X . A contradiction).

Assume $\dim(F) > 0$; then $(\text{image}(\pi_1(F) \rightarrow \pi_1(Y)))$ would be infinite, again because $\mathbb{C}^d$ does not contain positive dimensional subvarieties. Because $\pi_1(X)$ is abelian, this implies by dualising that the restriction map $H^0(X, \Omega_X^1) \rightarrow H^0(F, \Omega_F^1)$ is nonzero. A contradiction to the fact that F is a fibre of α $\square$

We can slightly extend 7.10 to the relative case, which shows that interesting linear representations of $\pi_1(X)$, $X \in \mathcal{C}$, come from the core:

Theorem 7.14. *Let $X \in \mathcal{C}$, let $c_X : X \rightarrow C(X)$ be its core with F its generic fibre, and let $\rho : \pi_1 \rightarrow \text{Gl}(N, \mathbb{C})$ be a linear representation.*

Let G (resp. H) be the image of $\pi_1(X)$ by ρ (resp. the intersection with G of the Zariski closure of the image H of $\pi_1(F)$ by ρ). Then H is normal in G , almost abelian, and there is a natural quotient linear representation $\rho' : \pi_1(C'(X)) \rightarrow (G/H) \subset \text{Gl}(N', \mathbb{C})$, where $C'(X)$ is a suitable finite étale cover of $C(X)$.

Proof: This is an immediate consequence of 7.10, using the well-known fact ([C2], for example) that in the situation of 7.14, if $K := \text{Ker}(\pi_1(X) \rightarrow \pi_1(C(X)))$, then $\pi_1(F)_X \subset K$, and the quotient group is generated by torsion elements. Thus, if $K' := \rho(K)$, since H is almost abelian by 7.10, and the quotient group (G/H) is linear by standard linear algebraic group theory, we have an embedding $(G/H) \subset \text{Gl}(N', \mathbb{C})$, for some N' , and we see that (K'/H) is the trivial group by A.Selberg's theorem, provided we replaced X by some suitable finite étale cover.

We thus get then an isomorphic morphism of groups: $\mu : (G/K') \rightarrow (G/H)$, which gives the asserted representation ρ' $\square$

8. An Orbifold generalisation of Kobayashi-Ochiai's Extension theorem

8.1. Statements. We give here an orbifold version of a famous extension theorem, due to Kobayashi-Ochiai ([K-O]).

The setting considered in all of this section 8 is the following:

Notation 8.1. *V be a connected complex manifold, D a reduced divisor on V , and $U := V - D$.*

Let $\phi : U \dashrightarrow X$ be a meromorphic map, with $X \in \mathcal{C}$.

We always assume $\psi := f \circ \phi : U \dashrightarrow Y$ is nondegenerate (ie: has maximal rank $p = \dim(Y)$ somewhere)

Let $f : X \dashrightarrow Y$ be a fibration, with $p := \dim(Y)$.

Our main result here is:

Theorem 8.2. *Let V, U, X, Y, ϕ, f be as above.*

assume f is a fibration of general type. Then:

(a) *ψ extends meromorphically to $\psi' : V \dashrightarrow Y$.*

(b) *For any $m > 0$ sufficiently divisible and $s \in H^0(Y, m(K_Y + \Delta(f)))$, $\psi^*(s)$ extends to:*

$(\psi')^(s) \in H^0(V, (\Omega_V^p)^{\otimes m}((m-1).D))$.*

Remark 8.3. *The result of Kobayashi-Ochiai is the special case where $X = Y$, so that X is of general type.*

The proof will be given in the next sections. It is an orbifold modification of the original proof of [K-O].

We shall first give some applications of this result, which are criteria of transcendental nature for certain varieties to be special.

Definition 8.4. *Let $U = V - D$ be the complement of a reduced divisor D in the connected complex manifold V . Say that the pair (V, D) is **log-special** if: for $p > 0$, there is no rank one coherent subsheaf $L \subset (\Omega_V^p)$ and no $m > 0$ such that the complete linear system defined by $H^0(V, mL + (m - 1)D)$ is of general type (ie: defines a meromorphic map of maximal rank p).*

As a first application, we immediately get from 8.2:

Corollary 8.5. *Let $U = V - D$ be the complement of the divisor D in the connected complex manifold V . Let $\phi : U \dashrightarrow X$ be nondegenerate.*

Assume that the pair (V, D) is log-special. Then: X is special.

We now consider the case when $V = X$ and D is empty. So the extension part of 8.2 is certainly trivial. Observe that the pair $(V, D) := (X, 0)$ is log-special precisely if there is no Bogomolov sheaf on X . We however recover one direction of 2.26:

Corollary 8.6. *Assume that X has no Bogomolov sheaf. Then X is special.*

To get some concrete applications, we take more restrictive conditions than log-specialness.

Definition 8.7. *Let $U = V - D$ be the complement of a reduced divisor D in the connected complex manifold V . Say that the pair (V, D) is **log-RC** (for log-rationally connected) if: $H^0(V, (\Omega_V^p)^{\otimes m}((m - 1)D))$ vanishes for any $m > 0$.*

Of course, if (V, D) is log-RC, it is log-special, if V is projective rationally connected and D is empty, the pair (V, D) is log-RC.

One easily checks the following (shown in a generalised form below):

Example 8.8. *The pair $(\mathbb{P}^n, \mathbb{P}^{n-1})$ is log-RC.*

From 8.2, we get:

Corollary 8.9. *Assume that the pair (V, D) is log-RC. Then: X is special.*

Specialising from Example 8.8, we get the:

Definition 8.10. *Say that X is $\mathbb{C}^d$ -dominable (resp. **covered by $\mathbb{C}^d$**) if there exists $\phi : \mathbb{C}^d \dashrightarrow X$, nondegenerate (resp. if the universal cover of X is isomorphic to $\mathbb{C}^d$).*

The terminology $\mathbb{C}^d$ -dominable is from [B-L]

From 8.9, we then deduce:

Corollary 8.11. *Let $X \in \mathcal{C}$ be $\mathbb{C}^d$ -dominable. Then X is special.*

In particular: if X is covered by $\mathbb{C}^d$, then X is special.

Remark that manifolds $X \in \mathcal{C}$ covered by $\mathbb{C}^d$ are conjectured by S. Itaka to be covered by a torus (ie: to have a finite étale cover which is a torus). This is shown to be true for surfaces in [I2], and for threefolds in [C-Z]. For X projective of arbitrary dimension, this follows (see [C-Z]) from the standard conjectures of the Minimal Model Program. See 7.12 for another case where this conjecture is known to hold true.

We now give a geometric criterion for the pair (V, D) to be log-RC. This criterion may explain the name log-RC, and shows that being log-RC may be seen as a transcendental analogue of rational connectedness.

Definition 8.12. *We say the pair (V, D) is H' -special if there exists a meromorphic map $\phi : U \dashrightarrow X$, where $U = W \times \mathbb{C}$, with W quasi-projective, and moreover such that ϕ is holomorphic and constant at the generic point of $W \times \{0\}$ (ie: $\phi(w, 0) = a$, for some fixed $a \in X$ and any $w \in W$).*

Example 8.13. *The pair $(\mathbb{P}^n, \mathbb{P}^{n-1})$ is H' -special. (Blow-up a point outside of $\mathbb{P}^{n-1}$ to see this).*

Proposition 8.14. *The pair (V, D) is log-RC if it is H' -special.*

The easy but lengthy proof is deferred to section 8.8 below.

Although the following clearly results from the preceding observations, we state it separately:

Corollary 8.15. *Let $X \in \mathcal{C}$ be H' -special. Then X is special.*

Remark 8.16. *Due to 8.2, and in this situation, it is tempting to ask if X is special when V itself is special.*

This is not true, because transcendental holomorphic maps do not preserve multiplicities along D . The following example shows this.

Example 8.17. *Let B, E be elliptic curves, and $a_i \in B$ be distinct points, $i = 1, \dots, 4$. Let B^* be the complement of these points in B . Let $V := E \times B$ and $U := E \times B^* \subset V$. Let now $m \geq 2$ be an integer. Let now X be obtained from V by applying to V the four logarithmic transformations of multiplicity m respectively to its fibres $E \times \{a_i\}$ over B .*

Then X is projective, and contains U as a Zariski open subset. However, V is special, while X is not.

So the specialness of V does not imply that of X , even when ϕ is the embedding of a Zariski open subset of both V and X . (Except when X is a curve, as observed by M. Zaidenberg (this is an immediate consequence of [K-O])).

8.2. Sketch of Proof of the Kobayashi-Ochiai's Extension Theorem. We shall very briefly first sketch the proof of the classical result of Kobayashi-Ochiai, and then indicate in section 8.6 the changes needed to obtain the orbifold version.

We consider here only the extension part of the statement (ie: the fact that ψ extends meromorphically to V). The second part (the control of the order of poles of $(\psi')^*(s)$) is shown in section 8.7 by a lemma of independent interest.

Recall we want to show that if $\psi : U := (V - D) \dashrightarrow Y$ is nondegenerate, then ψ extends meromorphically to V , provided Y is of general type. The idea is thus that maps to general type varieties do not have essential singularities. This is analogous to the fact that a bounded function on the

unit disc minus the origin extends across the origin. Now general type manifolds behave like manifolds with universal cover a bounded domain, in the sense that they admit pseudo-volume forms with negative Ricci form.

The proof (as explained in, [Kob], Chap.7) goes in several steps:

(1) It is sufficient to deal with the special case $V = \mathbb{D}^p$, $D = \{0\} \times \mathbb{D}^{p-1}$, so that:

$$U = \mathbb{D}^p - \{0\} \times \mathbb{D}^{p-1} := \mathbb{D}^{(p)}.$$

This is simply because meromorphic maps to a projective variety (or even a Kähler manifold) extend meromorphically across codimension 2 or more analytic subsets.

The case where $\dim(V) > p$ can be reduced to the equidimensional case (see [Kob], p.374-75).

Notations: For future reference, we denote by $\mathbb{D}(r)$ the disk of radius r centered at 0 in $\mathbb{C}$, by $\mathbb{D} := \mathbb{D}(1)$ the unit disk; $\mathbb{D}^p(r)$ and $\mathbb{D}^{(p)}(r)$ are constructed as $\mathbb{D}^p$ and $\mathbb{D}^{(p)}$ above, replacing $\mathbb{D}$ by $\mathbb{D}(r)$.

(2) Because Y is of general type, it has a pseudo-volume form w of Ricci curvature negative, bounded from above by a constant $-1/C < 0$.

This form is constructed out of a basis of a linear subsystem L of $m.K_Y$ giving an embedding of Y into some projective space (once its fixed components have been removed).

Another property used in the proof is that if w_j is the pseudo-volume form associated to a single element of the basis of L above, the quotient w_j/w is a smooth function on Y . This function is thus bounded.

These notions and constructions are explained in more detail in section 8.4 below.

(3) An elementary lemma (see 8.22 below), together with the argument exposed in (5) below, show that the special case (1) above is true, if one shows that w being the preceding pseudo-volume form on Y , then the integral of $\psi^*(w)$ converges on $\mathbb{D}^{(p)}(1/2)$.

(4) By the celebrated Ahlfors-Schwarz lemma (see 8.21 below), we have the estimate: $w \leq C.\beta$, if β is the homogeneous volume form on $\mathbb{D}^{(p)}$, with constant Ricci curvature -1 , and C is the positive constant of (2) above.

One concludes from the elementary fact that β has finite volume on $\mathbb{D}^{(p)}(1/2)$.

(5) Let us now conclude the proof in the equidimensional case: the forms $w_j := \psi^*(s_j)$ of the linear system $\psi^*(m.K_Y)$ extend meromorphically to $\mathbb{D}^p$, by the steps (2,3,4,5). But the map ψ is defined by the N -tuple of maps: $(w_0 : w_1 : \dots : w_N)$ with values in $\mathbb{P}^N$. The map ψ itself thus extends meromorphically to $\mathbb{D}^p$, as claimed.

We shall refer to [Kob] for the details not given here. For the main points (2,3,4), we shall introduce the basic definitions and properties in the sections 8.4 and 8.3 below. Finally, the proof of the orbifold version will be given in section 8.6.

The proof of point (b) in the statement of Theorem 8.2 will be given in the separate section 8.7 below. (Although it might possibly be deduced from [Kob], which contains a similar result in implicit form, we prefer to give an independent, more general statement).

8.3. Pseudo-Volume and Ricci Forms. We need to recall some classical notions. See, for example, [Kob], Chapters 1 and 7 for more details.

Let M be a complex manifold. We say that v is a **pseudo-volume form** (with holomorphic degeneracies) on M if it is a form of type (p, p) on M , with $p := \dim(M)$, such that locally, in any coordinate system $(x_1, \dots, x_p) = (x)$, one has: $v = |a|^2 . T.vol_{(x)}$, with: $vol_{(x)} := (i^{p^2} . d(x) \wedge \overline{d(x)})$, a

is a holomorphic nonzero function, and T a smooth (ie: $\mathcal{C}^\infty$) everywhere positive function locally defined on M .

For such a form v , its **Ricci form** is the real $(1, 1)$ -form: $Ricci(v) := -dd^c(\log T) = -i\partial\bar{\partial}(\log T)$, and its **Ricci function** is: $K_v := (-Ricci(v))^m/v$.

Both are well-defined independently of local coordinates, because changing charts multiplies T by the square of the modulus of the Jacobian, the $\log$ of which has vanishing dd^c .

From the definitions, one gets also immediately the functoriality of these notions: if $f : N \rightarrow M$ is a dominant holomorphic map, with $p = \dim(N)$, and if $v' := f^*(v)$, then the definitions apply to v' , and one has both: $Ricci(v') = f^*(Ricci(v))$ and $K_{v'} = f^*(K_v)$.

Remark 8.18. *The Ricci form is thus the special case for $-K_M$ of the notion of curvature form of a singular hermitian metric on a line bundle, neglecting the singular part (cohomologically significant, however) of the curvature current.*

Assume that v has holomorphic degeneracies. Then we say that v has **negatively bounded Ricci curvature** if $(-Ricci(v))$ is an everywhere positive or zero $(1, 1)$ -form, and if $-K_v \geq C$ for some constant $C > 0$, everywhere on M .

Observe that K_v is then nonpositive from the condition on $Ricci(v)$, and that K_v may take the value $(-\infty)$ at points where a vanishes. Note that if M is compact and $-Ricci(v)$ everywhere positive, then v has negatively bounded Ricci curvature.

Observe too that v' has negatively bounded Ricci curvature if and only if so has v , for $v' = f^*(v)$ as above.

Example 8.19. *We give the examples used in the proof. As said above, these are bounded domains.*

(1) The unit disc. Take (in the linear coordinates) $v := idx \wedge \overline{dx}/(1 - |x|^2)^2$.

Then $Ricci(v) = -v$, and $K_v = -1$.

(2) The unit disc with origin removed $\mathbb{D}^*$. Take (in the linear coordinates):

$v := idx \wedge \overline{dx}/(|x|^2 \cdot (\log(|x|^2))^2)$. One again has: $Ricci(v) = -v$, and $K_v = -1$. Actually the forms in the preceding two examples correspond under the universal covering map from $\mathbb{D}$ to $\mathbb{D}^*$. This can be checked explicitly using the Poincaré upper half plane as an intermediate step.

An important property used in the proof is that $\int_{\mathbb{D}^*(r)} v < +\infty$ if $r < 1$. (Of course, the domain over which the integral is taken is the disc of radius r with origin deleted);

(3) $M = \mathbb{D}^{(p)}$. Take (in the linear coordinates):

$v := i^{p^2} dx \wedge \overline{dx}/(|x_1|^2 \cdot (\log(|x_1|^2))^2 \cdot \prod_{2 \leq j \leq p} (1 - |x_j|^2)^2)$.

One again has: $Ricci(v) = -v$, and $K_v = -1$.

Again, one has: $\int_{\mathbb{D}^{(p)}(r)} v < +\infty$.

The above volume forms have thus negatively bounded Ricci function.

We give in the next section the example used in the proof in an essential way.

8.4. Pseudo-Volume forms on Varieties of General Type. Let Y be a p -dimensional manifold of general type, that is: with K_Y big.

From Kodaira's lemma, for m large enough, we can write: $mK_Y = H + A$, with H effective and A very ample.

Let L be a free linear subsystem of the complete linear system $|A|$, such that the associated regular map $\Lambda : Y \rightarrow \mathbb{P}^N$ ($N = \dim L$) is an embedding.

Let $(s) := (s_0, s_1, \dots, s_N)$ be a (complex) basis of $H + L$, and h a section of $\mathcal{O}_Y(H)$ vanishing exactly on H .

Define $w := (\sum_{j \in \{0, \dots, N\}} i^{m \cdot p^2} s_j \wedge \overline{s_j})^{1/m}$.

Then w is a pseudo-volume form with holomorphic degeneracies along H , since it is written in local coordinates as: $w = |a|^2 \cdot w_1$, with $w_1 = (\sum_{j \in \{0, \dots, N\}} |t_j|^2)^{1/m} \cdot \text{vol}_{(x)}$, if a is a local equation for H , and $s_j = t_j \cdot (\text{vol}_{(x)})^{\otimes m}$, $\forall j$ (using the notation introduced in section 8.2 above).

Notice that L being free, w_1 is a smooth volume form on Y . We now compute its Ricci form.

By its very definition, we have: $\text{Ricci}(w) = -\Lambda^*(\Theta)$, where Θ is (up to a positive constant) the curvature form of the Fubini-Study metric on the tautological line bundle $\mathcal{O}_{\mathbb{P}^N}(1)$.

(This is in accordance with remark 8.18: the Ricci form of w is just $-\Lambda^*(\Theta)$, since the singular part of the metric is given by h).

The Ricci form of w is thus negative everywhere. As a consequence: $K_w := (\text{Ricci}(w)^p / v)$ is negative everywhere (and $-\infty$ on H). By the compactness of Y , we conclude that this function is negatively bounded by some constant $-1/C$, for some $C > 0$.

Remark 8.20. For $j \in \{0, \dots, N\}$, define: $w_j := (i^{m \cdot p^2} s_j \wedge \overline{s_j})^{1/m}$. This is still a pseudo-volume form with holomorphic degeneracies on Y . The function w_j/w is obviously smooth on Y , and thus bounded. There exists $B > 0$ such that $w_j \leq B \cdot w$, $\forall j \in \{0, \dots, N\}$.

8.5. The Lemma of Ahlfors-Schwarz. Despite its simplicity, it is a very powerful tool to obtain bounds on the growth of pseudo-volume forms. We give the simplest version, sufficient for our applications.

Lemma 8.21. Let v be a pseudo-volume form with holomorphic degeneracies on $\mathbb{D}^{(p)}$.

Assume that $K_v \leq -1/C < 0$ everywhere on $\mathbb{D}^{(p)}$. Then $v \leq \beta$, where β is the volume form on $\mathbb{D}^{(p)}$ with constant Ricci function -1 defined in example 8.19 (3) above.

Sketch of proof: See [Kob], (2.4.14) for details. One can reduce (by introducing the volume forms relative to polyradii $r < 1$) to the case when the quotient (smooth) function $k := v/\beta$ has a maximum at an interior point b of (the closure of $\mathbb{D}^{(p)}$). Obviously, v does not vanish at b .

The real $(1,1)$ -form $dd^c(\log k) = i\partial\bar{\partial}(\log k)$ is then nonnegative at b . (By functoriality, it is sufficient to check this on a complex parametrised curve. But then: $i\partial\bar{\partial}(\log k) = (\partial^2(\log k)/\partial s^2 + \partial^2(\log k)/\partial t^2) \cdot ds \wedge dt$ in real coordinates $x = s + it$, and the claim follows).

Rescaling v , we assume that $C = 1$. Now: $i\partial\bar{\partial}(\log k) = i\partial\bar{\partial}(\log V) - i\partial\bar{\partial}(\log B) = \text{Ricci}(\beta) - \text{Ricci}(v) \leq 0$, if V, B are the functions such that $v = V \cdot \text{vol}_{(x)}$ and $\beta = B \cdot \text{vol}_{(x)}$.

We get: $\text{Ricci}(\beta) \leq \text{Ricci}(v)$. Hence also: $-v \geq K_v \cdot v = \text{Ricci}(\beta)^{\wedge p} \leq \text{Ricci}(v)^{\wedge p} = K_\beta \cdot \beta = -\beta$, and the conclusion follows $\square$

We now state an extension criterion which is used crucially in the proof.

Proposition 8.22. Let $s \in H^0(\mathbb{D}^{(p)}, m \cdot K_{\mathbb{D}^{(p)}})$ be such that:

$\int_{\mathbb{D}^{(p)}(r)} (i^{m \cdot p^2} s \wedge \overline{s})^{1/m} < +\infty$, for some $r < 1$.

Then s extends to a meromorphic section of $mK_{\mathbb{D}^p}$. Moreover, the poles of this extension are of order at most $(m-1)$ along $\{0\} \times \mathbb{D}^{p-1}$.

The proof is given in [Kob], (7.5.7-8). The proof reduces to the case when $p = 1$ and the elementary fact that a holomorphic function on the punctured unit disc which is $L_{(2/m)}$ on some $\mathbb{D}(r), r < 1$ extends meromorphically with a pole of order at most $(m - 1)$.

8.6. Proof of the Orbifold Version. .

Setting: We consider in this section the following data: $f : X \dashrightarrow Y$ is a fibration of general type, $\phi : U \dashrightarrow X$ is a meromorphic map which is dominant (ie: submersive at some point), X, V are connected complex manifolds, with X compact, and $j : U \subset V$ is a Zariski dense open subset, complement of some divisor D of V .

Let $\psi := f \circ \phi : U \dashrightarrow Y$. Blowing-up X and Y if needed, we shall assume that f is holomorphic, that Y is projective, and that f is high (see 4.1).

We let $p > 0$ be the dimension of Y , and $\Delta := \Delta(f)$ be the multiplicity divisor for f , as defined in 1.1.4

Our purpose here is to establish that ψ extends meromorphically to V .

Proposition 8.23. *Let $m > 0$ be an integer such that $m.\Delta$ is Cartier. Then:*

- (1) *f defines an injection of sheaves $f^* : \mathcal{O}_Y(m.(K_Y + \Delta)) \rightarrow (\Omega_X^p)^{\otimes m}$, after suitable modifications of X and Y .*
- (2) *In particular, let $s \in H^0(Y, m.(K_Y + \Delta))$. Then $f^*(s) \in H^0(X, (\Omega_X^p)^{\otimes m})$. (Recall Y has dimension p).*

Proof: Just as in the proof of 4.20, because we have an injection of sheaves (also denoted f^*): $f^* : \mathcal{O}_Y(m.(K_Y)) \rightarrow (\Omega_X^p)^{\otimes m}$, it is sufficient to check the above injection outside a codimension two subvariety of X , provided f is “high”.

So we first assume that we are near (in the analytic topology) the generic point of Δ_i .

But the assertion follows here from an easy local computation: indeed m_i divides $m = (m'.m_i)$ by assumption, and we can choose local coordinates:

$(x) = (x_1, \dots, x_d)$ near the generic point of the divisor $D_i \subset X$, a component of $f^{-1}(\Delta_i)$ mapped surjectively to Δ_i by f in such a way that in suitable local coordinates $(y) := (y_1, \dots, y_p)$, near the corresponding generic point of Δ_i , one has:

$f(x) = (y) = (y_1, \dots, y_p)$, with $y_1 = (x_1)^{m_i+m''}$, and $y_j = x_j$ for $j > 1$, with m'' some appropriate nonnegative integer.

(Local equations for D_i and Δ_i being respectively $x_1=0$ and $y_1=0$).

Thus we get a local generator of $f^*(\mathcal{O}_Y(m.(K_Y + \Delta)))$ in the form: $f^*(d(y)/y_1^{(1-1/m_i)})^{\otimes m}$, and this expression is equal to:

$$f^*(d(y)^{\otimes m}/y_1^{(m-m')}) = x_1^{(m_i+m''-1)m-(m_i+m'').(m-m')} . d(x')^{\otimes m} = (x_1)^{m'.m''} . d(x')^{\otimes m},$$

up to a nonzero constant factor, where $d(y) := dy_1 \wedge \dots \wedge dy_p$ and $d(x') := dx_1 \wedge \dots \wedge dx_p$.

Hence the claim, since the exponent of x_1 in the last expression is obviously nonnegative.

We thus see that f^* is well-defined outside the finite union B of all divisors of X which are mapped to codimension two or more analytic subsets of Y . By a suitable composition of blow-ups $u : X' \rightarrow X$ and $v : Y' \rightarrow Y$, we get $f' : X' \rightarrow Y'$ holomorphic such that $f.u = v.f'$, and moreover such that the strict transform B' of B in X' has all of its irreducible components mapped onto a divisor of Y' by f' (depending on the component, of course).

Then $g := v^{-1} \circ f = f' \circ u^{-1} : X \dashrightarrow Y'$ is holomorphic outside a codimension two or more analytic subset A of X (the indeterminacy locus of u^{-1}).

But then $g^*(\mathcal{O}_{Y'}(m.(K_{Y'} + \Delta(f'))))$ injects into $(\Omega_X^p)^{\otimes m}$ over $(X - A)$. By Hartog's theorem (for example), this injection extends through A . Now composing with $u^* : (\Omega_X^p)^{\otimes m} \rightarrow (\Omega_{X'}^p)^{\otimes m}$, we see that $(f')^*$ injects $(\mathcal{O}_{Y'}(m.(K_{Y'} + \Delta(f'))))$ into $(\Omega_{X'}^p)^{\otimes m}$, as asserted.

The second assertion is an immediate consequence of the first $\square$

The first part of this argument (in simpler form) applies to give:

Proposition 8.24. *Let $u :: Y' \rightarrow Y$ be a Δ -nice covering, in the sense of 4.19. But we choose u such that it ramifies at order exactly m_j above each component Δ_j of $\Delta(f)$. Then:*

- (1) *u defines an injection of sheaves: $u^* : \mathcal{O}_Y(m.(K_Y + \Delta)) \rightarrow \mathcal{O}_{Y'}(m.K_{Y'})$.*
- (2) *In particular, let $s \in H^0(Y, m.(K_Y + \Delta))$. Then $u^*(s) \in H^0(Y', m.K_{Y'})$.*

We now come to the crucial point of our modification of the proof of the extension theorem of Kobayashi-Ochiai. In a first step, we proceed as in section 8.4, by the construction of a pseudo-volume form on Y with meromorphic-not holomorphic- degeneracies.

Notation 8.25.

Let $\phi : \mathbb{D}^{(p)} \dashrightarrow X$ be such that $\psi := f \circ \phi : \mathbb{D}^{(p)} \dashrightarrow Y$ is nondegenerate. We need to extend ψ meromorphically to $\mathbb{D}^p$.

Since $(K_Y + \Delta)$ is big on Y , we can write, by Kodaira's lemma: $m(K + \Delta) = H + A$, with A very ample and H effective on Y , m being choosen sufficiently divisible (by the l.c.m of the m_i 's) and sufficiently big.

Let $h \in H^0(Y, \mathcal{O}_Y(H))$ be an element defining H .

Let $\delta \in H^0(Y, \mathcal{O}_Y(m.\Delta))$ be a section vanishing exactly on $m\Delta$, with the right multiplicities.

For $\sigma \in |A|$, the complete linear system defined by H , we denote by:

$$s := h.\sigma. \in H^0(Y, m.(K_Y + \Delta)).$$

Let now $\sigma_0, \dots, \sigma_N$ a basis of $H^0(Y, A)$; let also $s_j := h.\sigma_j$, for $j \in J := \{0, \dots, N\}$.

Let further $t_j := s_j/\delta, \forall j$. These are meromorphic sections of $m.K_Y$.

Define $v := (\sum_{j=0}^{j=N} (i^{p^2.m} t_j \wedge \bar{t}_j))^{1/m}$. Thus v is a global pseudo-volume form with meromorphic degeneracies on Y .

Notice however that, by Proposition 8.23 above, $f^*(t_j)$ is, for any j , a holomorphic section of $(\Omega_X^p)^{\otimes m}$. This is simply because in local coordinates on Y , we can write:

$t_j = (T_j/d).dy$, for some holomorphic function T_j , where d (resp. dy) is a local generator of the ideal defining $m\Delta$ (resp. K_Y). (I thank M.Paun for this important observation). The assertion thus follows from 8.23 and the fact that dy/d is a local section of $m.(K_Y + \Delta)$.

In particular, $\psi^*(t_j)$ is a holomorphic section of mK_M , with $M = \mathbb{D}^{(p)}$, and so $w := \psi^*(v)$ is naturally a pseudo-volume form with holomorphic degeneracies on M .

For the same reason, $f^*(s_j)$ is, for any j , a holomorphic section of $(\Omega_X^p)^{\otimes m}$.

Exactly in the same way we defined $w := \psi^*(v)$, we can define a pseudo-volume form with holomorphic degeneracies $\bar{w}$ on M by: $\bar{w} := (\sum_{j=0}^{j=N} (i^{p^2.m} \psi^*(s_j) \wedge \overline{\psi^*(s_j)}))^{1/m}$.

Notice that here, however, there is no pseudo-volume form $\bar{v}$ on Y such that $\bar{w} = \psi^*(\bar{v})$.

The crucial point is now:

Lemma 8.26. *Ricci($\bar{w}$) = Ricci(w) = $\psi^*(-\Lambda^*(\Theta))$, if $\Lambda : Y \rightarrow \mathbb{P}^N$ is the regular map defined by the complete linear system $|A|$ on Y , and Θ the curvature form of the Fubini-Study metric on $\mathbb{P}^N$. In particular, Ricci(w) is thus negative everywhere, and there exists $C > 0$ such that $K_{\bar{w}} \leq -1/C$ everywhere on $\mathbb{D}^{(p)}$.*

Proof: Write again $t_j = (T_j/d).dy$, as above. We deduce that: $v = |h/d|^{2/m} \cdot V \cdot \text{vol}_{(y)}$, if h is a local equation for H , and $\text{vol}_{(y)}$ a local volume form on Y . Here $V = (\sum_{j \in \mathbb{N}} |T_j|^2)^{1/m}$.

Now, from 8.23 again, we see that $\psi^*(\text{vol}_{(y)}/|d|^{2/m}) = |G|^2 \cdot \text{vol}_{(M)}$, for a certain holomorphic function G defined on $\psi^{-1}(U)$, for $U \subset Y$, the open subset where the said trivialisations and charts are defined.

Recall that $-i\partial\bar{\partial}V = -\Lambda^*(\Theta)$, by 8.4 above.

A computation of the Ricci curvature then gives the first assertion for w .

The assertion for $\bar{w}$ is deduced from the fact that $\bar{w} = \psi^*(|d|^{2/m}).w$, so that these two pseudo-volume forms have the same Ricci-form.

We then get:

$$K_w = \psi^*(-\Lambda^*(\Theta))^{\wedge p} / \psi^*(v) = \psi^*(-\Lambda^*(\Theta))^{\wedge p} / \psi^*(|h.G^m|^{2/m} \cdot V) \cdot \text{vol}_M$$

$$\text{And so, } K_{\bar{w}} = \psi^*(|d|^{-2/m} \cdot K_w) \leq (-1/C) \cdot (1/\psi^*(|h|^{2/m} \cdot V)),$$

because V is differentiable, and h locally bounded from above everywhere on Y .

To express things differently, one can compute the Ricci function of $\bar{w}$ symbolically, as if it were of the form $\psi^*(\bar{v})$, with $\bar{v} := (\sum_{j=0}^{j=N} (i^{p^2 \cdot m} s_j \wedge \bar{s}_j))^{1/m}$. $\square$

By the argument given in section 8.2, to show 8.2, it is sufficient to show that each of the $\psi^*(s_j)$'s extend meromorphically to $\mathbb{D}^p$, and this is true if the integral of w over $\mathbb{D}^{(p)}(r)$ is convergent, for some $0 < r < 1$.

To show this convergence, we simply need, by the Ahlfors-Schwarz Lemma, to show that w has everywhere negatively bounded Ricci function (by some constant $-1/C < 0$). But this exactly what the preceding Lemma 8.26 claims. This concludes the proof of the first assertion of 8.2.

Remark 8.27. *Observe that the preceding proof shows with minor adaptations (using 8.24), the following orbifold version of Theorem 8.2: If (Y/Δ) is an orbifold of general type (Y smooth and Δ supported on a normal crossings divisor), then any nondegenerate meromorphic map $\phi : U \dashrightarrow (Y/\Delta)$ extends meromorphically to V . Here, to say that ϕ is a meromorphic map to the orbifold (Y/Δ) means that any point $u \in U$ at which ϕ is holomorphic, with $y := \phi(u)$ a smooth point of the support of Δ , lying on Δ_i , then ϕ lifts locally around u to a holomorphic map to the ramified cover of Y near y , ramifying at order exactly m_i above Δ_i .*

8.7. Extension of Pluri-Canonical Meromorphic Forms. .

We now want to control the order of the poles along D of the meromorphic extension to V of $\psi^*(s) \in H^0(U, (\Omega_U^p)^{\otimes m})$, for $s \in H^0(Y, m \cdot (K_Y + \Delta))$. More precisely, notations being as above:

Proposition 8.28. *For any $m > 0$ sufficiently divisible and s any element of $H^0(Y, m(K_Y + \Delta(f)))$, $\psi^*(s)$ extends to $(\psi')^*(s) \in H^0(V, (\Omega_V^p)^{\otimes m}((m-1).D))$.*

Proof: Recall that the support of Δ is of normal crossings. We can assume that ψ' is holomorphic, because the assertion holds for V if it does for any of its blow-ups.

Recall that s can be written in local coordinates as follows, s being a section of $m(K_Y + \Delta(f))$:

$s = (d(y)/(y_1^{(1-1/m_1)} \dots y_p^{(1-1/m_p)}))^{\otimes m}$, if Δ has local equation the denominator of this expression. (Some of the m_i may be one, and so the corresponding terms don't contribute).

Let now $(v) = (v_1, \dots, v_n)$ be local coordinates near a generic point of some component D' of D , such that D' has local equation $w := v_1 = 0$. We can assume that near this point the map ψ' is given by $\psi'(v) = (v_1^{r_1} \cdot Y_1, \dots, v_1^{r_p} \cdot Y_p)$, for integers r_i and nowhere vanishing functions Y_i of (v) .

Computing, we get (letting r be: $r_1 + \dots + r_p$): $(\psi')^*(s) = (w^{r-1} \cdot \Theta / w^{r-s} \cdot Y')^{\otimes m}$, with $w := v_1$, for some holomorphic p -form Θ on V , and Y' a nowhere vanishing function on V (the product of the Y_i 's), and $s := r_1/m_1 + \dots + r_p/m_p$. This shows the proposition, because $s \cdot m$ is a positive integer, since each m_i divides m $\square$

8.8. Proof that H' -Specialness Implies Log-Rational Connectedness. .

We prove here proposition 8.14.

We use the notations and setting of 8.1, 8.12 and 8.7.

Proof: We just need to show that any element s' of $H^0(V, (\Omega_V^p)^{\otimes m}((m-1).D))$ vanishes, provided it is on U of the form $\phi^*(s'')$ for some $s'' \in H^0(X, (\Omega_X^p)^{\otimes m})$. For this, we shall just show that its restriction to any $F := \{w\} \times \mathbb{P}^1(\mathbb{C}) \subset V$ has to vanish for any $w \in W$. The argument is similar to (and motivated by) the one showing that sections of $(\Omega_X^p)^{\otimes m}$ vanish if X is rationally connected (see [C5]).

Let Ω^q denote the restriction of Ω_V^q to F , for $q \geq 0$, and let N be the (trivial) dual of the normal bundle of F in V .

From the exact sequence: $1 \rightarrow N \rightarrow \Omega_F^1 \rightarrow \Omega_F^1 \rightarrow 1$, we get, for $q > 0$:

$$(\textcircled{a}) 1 \rightarrow \wedge^q N \rightarrow \Omega^q \rightarrow \Omega_F^1 \otimes \wedge^{q-1} N \rightarrow 1$$

Notice that all these sequences are split.

We now fix $q = p > 0$, and denote respectively by $B := \wedge^p N$ the kernel, and $A := \Omega_F^1 \otimes \wedge^{p-1} N$ the quotient of the corresponding exact sequence $(\textcircled{a})$.

From this same split exact sequence, we get now for $(\Omega^p)^{\otimes m}$ a decreasing (split) filtration by subbundles:

$$\{0\} := W_{m+1} \subset W_1 \subset \dots \subset W_0 := (\Omega^p)^{\otimes m},$$

with successive quotients: $W_0/W_1 = A^{\otimes m}; \dots; W_k/W_{k+1} = A^{\otimes(m-k)} \otimes B^k; \dots; W_m = B^{\otimes m}$.

Notice that, for $k = 0, \dots, m$, the bundle $A^{\otimes(m-k)} \otimes B^k$ is isomorphic to $\mathcal{O}_F(-2(m-k))$ tensorised by a trivial vector bundle N_k on F .

We now take local coordinates $((w), t)$ on V near $(w, 0)$, where $(w) := (w_1, \dots, w_{n-1})$ are local coordinates on W , and t is a local coordinate near $0 \in \mathbb{P}^1$.

The map ϕ takes the form $\phi((w), t) = (t \cdot \phi_1, \dots, t \cdot \phi_d)$, for holomorphic functions ϕ_i of $((w), t)$, with $d := \dim(X)$, in local coordinates near $a \in X$, since $\phi(W \times \{0\}) = \{a\}$, by our H' -speciality assumption.

Locally, if σ'' is a p -form on X near a , then $\sigma' := \phi^*(\sigma'')$ is written: $\sigma' = dt \cdot \alpha + t \cdot \beta$, for α and β forms in the dw_i 's. If now σ'' is a p -form, then $\sigma' = t^{p-1} \cdot dt \wedge \alpha + t^p \cdot \beta$, with the same conditions.

We now consider the case when $s' = \phi^*(s'')$ for some $s'' \in H^0(X, (\Omega_X^p)^{\otimes m})$: from the preceding remarks, we can write $s'' = s_0'' + \dots s_k'' + \dots + s_m''$, where each of the s_k'' is the tensor product of $(m-k)$ terms of the form: $t^{p-1}.dt \wedge \alpha$, and of k terms of the form: $t^p.\beta$.

So that, for each k , s_k'' is actually the piece of degree k of s'' for the above graduation, and so turns out to be a section of the bundle $A^{m-k} \otimes B^k$ defined above, and vanishing at 0 to order $(m-k).(p-1) + k.p = m.(p-1) + k$.

Now, we assumed that s'' has a pole of order at most $(m-1)$ at the point at infinity of F . Thus s_k'' defines a meromorphic section of $\mathcal{O}_F(-2(m-k)) \otimes N_k$, with N_k a trivial bundle, this section vanishing at 0 to order $(m(p-1) + k)$ at least, and having a pole of order $(m-1)$ at most at infinity.

This section thus has to vanish, because otherwise the number of poles should be the opposite of the degree. But here: $(m-1) - m.(p-1) - k < 2(m-k)$ for any integers $m, p > 0$ and $0 \leq k \leq m$. This concludes the proof of 8.14 $\square$

9. The Core and the Kobayashi Metric

9.1. The Kobayashi Pseudo-Metric. We refer to [Kob] for a systematic treatment of the following notions.

Let X be a complex manifold. Let d_X be the **Kobayashi pseudometric** on X (which is the largest pseudometric d on X such that $h^*(d) \leq p_{\mathbb{D}}$ for any holomorphic map $h : \mathbb{D} \rightarrow X$, where $\mathbb{D}$ is the unit disc in $\mathbb{C}$ and $p_{\mathbb{D}}$ is the Poincaré metric on $\mathbb{D}$).

For example, it is easy to check (using the Ahlfors-Schwarz lemma) that $d_{\mathbb{C}^n} \equiv 0$, and $p_{\mathbb{D}} = d_{\mathbb{D}}$. A fundamental property of d is that holomorphic maps are distance decreasing: if $h : X \rightarrow Y$ is any holomorphic map, then $h^*(d_Y) \leq d_X$. Hence if $h : \mathbb{C} \rightarrow X$ is holomorphic, then d_X vanishes on the (metric) closure of the image of h . Rational and elliptic curves have thus vanishing d .

In particular, if $h : \mathbb{C} \rightarrow X$ is holomorphic, then d_X vanishes on the (metric) closure H of the image of h . Thus d_X will vanish identically if, for example, $H = X$, or if any two points of X can be connected by such a chain of H_i 's.

On the opposite side, X is said to be **hyperbolic** if d_X is a metric on X , defining its usual “coherent” (or “metric”) topology. The unit disc and the curves of genus 2 or more which it covers are hyperbolic.

A fundamental result of R. Brody states that a compact complex manifold X is hyperbolic if and only if every holomorphic map $h : \mathbb{C} \rightarrow X$ is constant.

The study of projective hyperbolic manifolds is guided by the two nearly converse conjectures of S. Kobayashi and S. Lang: Kobayashi's conjecture states that a hyperbolic projective manifold has ample canonical bundle (and is so of general type). Lang's conjecture states that a manifold X is of general type if and only if it is “pseudo”-hyperbolic (in Lang's terminology). Which means that there is a proper algebraic subset $S \subset X$ such that any holomorphic map $h : \mathbb{C} \rightarrow X$ has image contained in S . See [La 1,2] for more information, also on the relations with arithmetic geometry.

Notice that being pseudo-hyperbolic in the above sense is a priori weaker than being hyperbolic modulo S , in Kobayashi's terminology (which means that d_X is a metric outside S). Although it seems plausible that both properties are equivalent.

9.2. Hyperbolically Special Manifolds. Our aim here and in section 9.10 below is to give a conjectural description of d_X in the general case.

Definition 9.1. $X \in \mathcal{C}$ is said to be *H-special* (H for *Hyperbolically*) if $d_X \equiv 0$.

Our investigations are guided by :

Conjecture 9.2. (Conjecture III_H): *Let $X \in \mathcal{C}$. Then X is special if and only if it is H-special.*

We shall give a motivation for conjecture 9.2 in section 9.4 below.

In Section 9.10, we shall complete conjecture III_H by conjecture IV_H, which (hopefully) describes d_X for any $X \in \mathcal{C}$.

Examples 9.3. *We list special manifolds known to be H-special.*

(1) **Curves:** A curve is *H-special* iff special, that is: either rational or elliptic.

(2) **Rationally connected manifolds** are *H-special* (and special). This is obvious. More generally:

Proposition 9.4. *Let $f : X \rightarrow Y$ be a rationally connected fibration (ie: so is its generic fibre), with Y projective. Then: $d_X = f^*(d_Y)$. In particular, X is H-special iff so is $R(X)$.*

Proof: Clearly, there exists a pseudometric d_Y^+ on Y such that $d_X = f^*(d_Y^+)$.

We need to show that $d_Y^+ = d_Y$. The inequality $d_Y^+ \geq d_Y$ is obvious (by the decreasing property of d_X under holomorphic maps). By the algebraic approximation theorem of [D-L-S], it is sufficient to show this when Y is a curve. But then, by [G-H-S], f has a section. This provides us with the opposite inequality $\square$

Remark 9.5. *This is another instance in which it is shown that rationally connected manifolds are in many aspects “negligible”, also in fibrations: indeed we already saw that if $f : X \dashrightarrow Y$ is a rationally connected fibration with X projective, then X and Y have the same core, the same fundamental group, and the same Kobayashi (pseudo-)metric. Only the arithmetic aspects are still unknown. Although the function field case is, by [G-H-S], much better understood.*

Notice that, by the birational invariance of d_X , the above result applies if Y is Moishezon. In fact, as in 3.28, it should still be true when $X \in \mathcal{C}$.

(3) **Manifolds with trivial canonical class** They provide important examples of manifolds with $\kappa = 0$. Then conjecture III_H is obviously true for complex tori, true for K3-surfaces (see below).

Using Bogomolov decomposition theorem, it is sufficient to test the irreducible holomorphic symplectic manifolds and the Calabi-Yau manifolds.

The known symplectic manifolds form a very short list (The two series of Beauville-Fujiki, and the two examples of O’Grady). For all these examples, a direct verification seems quite feasible, by reduction to the case of abelian surfaces and K3-surfaces.

For others symplectic manifolds (if any), the method of [C90], obviously underexploited there, gives a very positive indication.

The first significant example of Calabi-Yau manifolds, that of quintics in $\mathbb{P}_4(\mathbb{C})$, might be accessible also, by a strategy explained to me by C.Voisin.

(4) **Manifolds with $\kappa = 0$:** This is a crucial case, motivated by the special case where K_X is trivial; the conjecture then implies (by 5.1) the vanishing of d_X . This question is standard, and was already raised in [Kob 76].

(5) **Finite étale covers** of an H -special X are H -special, too (by the invariance of Kobayashi pseudometric under étale covers).

(6) **H'' -special manifolds:**

Let us now consider the opposite implication (that H -specialness should imply specialness). Its difficulty lies in the fact that the vanishing of d_X has no known geometric translation in dimension 3 or more. See 9.16 for a simple discussion of this topic. So we start with a stronger hypothesis that gives such an implication.

Definition 9.6. *We say that $X \in \mathcal{C}$ is H'' -special if there exists a holomorphic map with dense image (in the metric topology) $\phi : U \rightarrow X$, where $U = W \times \mathbb{C}$, with W quasi-projective, and moreover such that $\phi(w, 0) = a$, for some fixed $a \in X$ and any $w \in W$.*

Then X is obviously H' - and H -special if it is H'' -special. By 8.12, X is also special.

In particular, of course, manifolds X which are $\mathbb{C}^d$ -dominable or covered by $\mathbb{C}^d$ are H -special. They are special, too by 8.11

9.3. Similarities. Three stability properties: We list here, to stress their analogies, three stability properties which are common to the classes of special and H -special manifolds:

(a) Let $f : X \rightarrow Y$ be a fibration, and $g : V \rightarrow X$ be holomorphic with $f \circ g : V \rightarrow Y$ surjective. Then X is special (resp. H -special) if so are V and the generic fibres of f .

(b) Assume X is S -connected (resp. H -connected). In this case, X is special (resp. H -special). Here being S -connected (resp. H -connected) means that two generic points of X can be joined by a chain of subvarieties of X which are special (resp. H -special).

(c) The Albanese map is surjective if X is special (resp. H -special, provided its Albanese torus is simple). See 5.3 (resp. 9.13 below) for the special (resp. H -special) case).

Notice that Conjecture III $_H$ implies that specialisations of special varieties are special, because this specialisation stability holds for H -special varieties, by the metric continuity of the Kobayashi metric.

9.4. The Origin of Conjecture III $_H$. Conjecture III $_H$ is motivated by the following:

Question 9.7. *Let $f : X \rightarrow Y$ be an admissible fibration. Assume that $d_F \equiv 0$ if F is a general fibre of f . Then $d_X = f^*(d_Y^+)$, for a certain pseudometric d_Y^+ on Y .*

(a) *Is then $d_Y^+ = d_{(Y/\Delta(f))}$ the Kobayashi pseudometric of the orbifold $(Y/\Delta(f))$ (defined in 9.23 below)?*

(b) *Is $d_{(Y/\Delta(f))} \equiv 0$ if either $\kappa(Y/\Delta(f)) = 0$, or if there exists a family of rational curves $(C_t)_{t \in T}$ with $X_t := f^{-1}(C_t)$ special for $t \in T$ general, and such that any two points of Y can be connected by a finite chain of such C_t 's?*

Notice that the above theorem 9.4 shows that the preceding question has a positive answer for rationally connected fibrations $f : X \rightarrow Y$, if Y is projective, because then $\Delta(f)$ is empty.

Let us show briefly how to deduce Conjecture III_H from a positive answer to the above question, assuming moreover Conjectures $\kappa = 0, -\infty$ expressing the core as a composition of orbifold Iitaka fibrations and rational quotients. We keep the notations of 6.4.

We argue by induction on the largest number $k \geq 0$ such that $s_X^k : X \rightarrow S^k(X)$ is not the constant map: if $(s_X^k)_R$ is the constant map, then: there exists a family of rational curves C_t as in the question (b) above. By the positive answer to that question, the orbifold pseudometric on the base vanishes, while the fibres have vanishing pseudometric by the induction hypothesis. Hence the conclusion, by (a). If $g := r \circ s_X^k$ is not constant, then s_X^{k+1} is constant. The preceding argument, applied to JB_g , instead of $(s_X^k)_R$ gives the conclusion.

The following result, due to [B-L] and [L] can be rephrased as giving a second example (abelian fibrations over curves) in which the above question 9.7 has a positive answer:

Theorem 9.8. *Let $f : X \rightarrow C$ be a fibration with C a curve, X projective, and the generic fibres of f being abelian varieties. Then 9.7 above has a positive answer.*

More precisely: $d_X = c_X^(d_C^*)$, where d_C^* vanishes if f is not of general type, and otherwise is a metric on C such that $g^*(d_C^*) = d_{C'}$, where $g : C' \rightarrow C$ is a finite Galois cover with C' of genus 2 at least, the map g having ramification order exactly m_i above each of the points c_i of $|\Delta(f)| \subset C$.*

9.5. Surfaces.

Proposition 9.9. *A compact Kähler surface not of general type is H -special if and only if it is special, and if and only if its fundamental group is virtually abelian.*

Thus conjecture III_H holds for them, except for potential surfaces of general type and vanishing d_X , the existence of which is excluded by Lang's conjectures.

For surfaces, conjecture III_H is thus equivalent to Lang's conjecture.

Proof: This follows easily from the results of [B-L] (which can be rephrased as saying that non- $K3$ special surfaces are $\mathbb{C}^2$ -dominable) and the fact that every $K3$ -surface is H -special, by the metric continuity of Kobayashi pseudometric used in conjunction with the usual argument of approximating an arbitrary $K3$ -surface by Kummer ones. (More generally, being H -special is a property which is closed under deformation in the metric topology). Thus special surfaces are H -special.

The reverse direction is clear, because a non-special surface not of general type has (by 3.30) a finite étale cover which maps surjectively onto a curve of genus two or more, and cannot thus be H -special $\square$

Remark 9.10. *Let $X \in \mathcal{C}$ be H -special with $\kappa(X) = 0, 1$. Then X is special. Indeed, the case $\kappa = 0$ (resp. $\kappa = 1$) follows from 5.1 (resp. 3.14).*

.

9.6. Threefolds. Our results are very partial. The complete solution of Conjecture III_H in this case depends in particular on the solution of several other difficult conjectures: Lang's conjecture in dimension 3, its orbifold generalisation (as stated in section 9.23 below) for orbifold surfaces, and finally the case where $\kappa = 0$.

Proposition 9.11. *Let X be a projective threefold.*

- (1) Assume $\kappa(X) \neq 3$, $\text{ess}(X) \neq 2$, and X is H -special. Then X is special.
- (2) Assume $\kappa(X) \neq 0, 2$, and X is special. Then X is H -special, unless possibly in the following case: $\kappa(X) = 1$ and J_X has generic fibres which are K3-surfaces.

Let us remark that the non-algebraic case presents some specific difficulties which can however be solved at the cost of some lengthy arguments. The case $\kappa(X) = 2$ left open in the above statement seems also accessible by known methods. We hope to come back on this in a later work.

The proof that the special 3-folds listed above are H -special does not present any difficulty, using the description of special 3-folds given in 3.38. The converse direction rests on the fact that if $f : X \dashrightarrow Y$ is surjective with X H -special, then Y is H -special too.

9.7. The Fundamental group. Conjectures 7.1 and III_H imply that if X is H -special, then its fundamental group should be virtually abelian.

It is thus natural to conjecture independantly that being H -special should force the fundamental group to be virtually abelian.

Notice that this would imply for example that being H' -special, or $\mathbb{C}^d$ -dominable, or being covered by $\mathbb{C}^d$ implies the virtual abelianity of the fundamental group. Recall that this last implication, by far the easiest one, is equivalent to S. Iitaka's conjecture, and is widely open.

A typical example of a question suggested by the preceding context is:

Question 9.12. Let $f : \mathbb{C} \rightarrow X$, $X \in \mathcal{C}$, be a holomorphic map with Zariski dense image. Is then $\pi_1(X)$ almost abelian?

9.8. The Albanese map. However, in complete analogy with the special case (see 5.3), we have, using strengthenings of a classical result of Bloch-Ochiai (surprisingly, it does not seem to have ever been stated that way):

Proposition 9.13. Let $X \in \mathcal{C}$ be H -special. Then its Albanese map $\alpha_X : X \rightarrow \text{Alb}(X)$ is surjective if $\text{Alb}(X)$ is a simple torus

Of course, the hypothesis that $\text{Alb}(X)$ is simple is artificial. It could be removed, using another result of Y. Kawamata in [Kw], provided one could positively answer question 9.14 below, which is a natural strengthening of Brody's theorem. (Alternatively, it is possible that a suitable sharpening of the proof of Bloch-Ochiai's theorem permits to remove this artificial hypothesis, without answering question 9.14 in general):

Question 9.14. Let X be a compact complex connected manifold. Let $Z \subset X$ be a proper compact analytic subset. Assume that any holomorphic map $h : \mathbb{C} \rightarrow X$ has image contained in Z . Is then the Kobayashi pseudo-metric d_X a metric on its complement X/Z ?

Notice also that certain recent results of Noguchi-Winkelmann (or at least the methods there) might imply that in 9.13 the Albanese map considered is surjective and has no multiple fibres in codimension one, as expected from Conjecture III_H and Theorem 5.3

Corollary 9.15. If $X \in \mathcal{C}$ which is H -special (and with simple Albanese torus), then:

- (a) any torsionfree solvable quotient of $\pi_1(X)$ is almost abelian.
- (b) any linear quotient of $\pi_1(X)$ is almost abelian.

Proof: The proof is exactly the same as that of 7.10 (where X was special), just observing that any finite étale cover of X satisfy the same hypothesis. Also we need to apply Zuo's result ([Z]) that if $\pi_1(X)$ had a non-trivial semi-simple linear representation, then X were generically Brody hyperbolic (in particular: not H -special) $\square$

9.9. Geometric variants of H -speciality. Brody's theorem says that the obstruction to the hyperbolicity of a compact X is due to the presence of “entire” curves in X , that is images of nonconstant holomorphic maps $h : \mathbb{C} \rightarrow X$. It is thus natural to try to express the fact that X is H -special by the geometry of “entire” curves lying on X . In this sense, being H -special appears as a transcendental analogue of rational connectedness.

Unfortunately, the analogue of a deformation theory for entire curves does not exist (and would certainly present great difficulties). So we introduced above the notions of H' -speciality, of H'' -speciality, and of $\mathbb{C}^d$ -dominability to try to give some examples in which the vanishing of d_X is deduced from the geometry of “entire” curves.

Remark 9.16. *Let $X \in \mathcal{C}$. One has the following implications between the different notions of speciality: $\mathbb{C}^d$ -dominability implies H'' -speciality, which implies both H -speciality and H' -speciality, which implies speciality (by 8.2).*

Of course, there are lots of other geometric variants of H -speciality which one can introduce, replacing for example the metric topology by the Zariski topology, and/or considering jets (see [Gr]), and which can be considered as approximations of H -speciality. One can ask about the relationships between the various notions so defined. In particular, do the notions relative to metric and Zariski topologies always coincide?

We shall just ask one question of whether some strong reverse implications hold between the various notions introduced in 9.16.

Question 9.17. *Assume X is H -special. Is X then also H'' -special?*

This question is motivated by the fact that a positive answer (which does not seem unrealistic, although far beyond reach if true) would imply, using 8.2, that H -special manifolds are special, which is one of the two implications of conjecture III_H.

It is shown in [B-L] that for surfaces not of general type, $\mathbb{C}^d$ -dominability, H' -speciality and H -speciality are equivalent.

This may however be a low dimensional phenomenon, similar to the fact that rational connectedness, unirationality and rationality coincide for complex surfaces.

This suggests the following obvious observations concerning rationally connected manifolds:

Remark 9.18. *Let X be a rationally connected manifold. Then X is H'' -special. If X is unirational, then X is $\mathbb{C}^d$ -dominable.*

This suggests of course also the following (again far beyond reach, since no rationally connected manifold is known which is not unirational, although most of them are suspected not to be):

Question 9.19. *Let X be a rationally connected manifold. If X is $\mathbb{C}^d$ -dominable, is X unirational?*

The answer is not expected to be necessarily positive. It seems quite possible that the class of rationally connected manifolds which are $\mathbb{C}^d$ -dominable is strictly intermediate between the other two ones (unirational and rationally connected), as the following example may suggest:

Example 9.20. *The threefold X constructed in [U,(11.7.1),p.137] is obviously $\mathbb{C}^3$ -dominated, and (less obviously) rationally connected. But it is asked by K.Ueno whether it is unirational or not.*

Recall that X is nothing but a smooth model of the quotient of E^3 , with E the Gauss elliptic curve, by the cyclic group of order 4 generated by complex multiplication by $i := \sqrt{-1}$ on each factor.

In the other direction (discrepancy between rational connectedness and $\mathbb{C}^d$ -dominability), even the $\mathbb{C}^3$ -dominability of the general quartic in $\mathbb{P}^4$ is far from obvious, if true.

9.10. The Core and the Splitting of the Kobayashi Metric. Let $X \in \mathcal{C}$, and let $c_X : X \rightarrow C(X) = Y$ be its core. By conjecture III_H, its general fibres are H -special, since special. Thus d_X should vanish on all of them, by Conjecture III_H and the metric continuity of d_X . There should thus exist a unique pseudometric $d_{C(X)} := d_Y^*$ on $Y = C(X)$ such that $d_X = c_X^*(d_Y^*)$.

We give a first version of:

Conjecture 9.21. Conjecture IV_H:

The pseudometric $d_{C(X)}^$ is a metric on some Zariski nonempty open subset $C(X)^*$ of $C(X)$.*

Remark 9.22.

- (1) Assume X is of general type. Then $c_X = id_X$, and we just recover the usual conjecture of Lang in its strong form.
- (2) Recall that, after Conjecture II, $(C(X)/\Delta(c_X))$ is an orbifold of general type. Thus Conjecture IV_H appears as an extension of Lang's conjecture to the orbifold case.
- (3) The preceding remark can be made more precise, if one introduces as in 9.23 below an orbifold Kobayashi pseudometric on any orbifold Y/Δ . One may then ask whether $d_{C(X)}^* = d_{C(X)/\Delta(c_X)}$. Then Conjecture IV_H appears as the orbifold extension of Lang's conjecture. See conjecture IV'_H below. We hope to come back on this latter.

We shall now propose an orbifold definition of the Kobayashi pseudometric:

Definition 9.23. *Let X be a complex manifold and $\Delta = \sum(1 - 1/m_i) \cdot \Delta_i$ be a $\mathbb{Q}$ -divisor on X , with m_i positive integers and Δ_i irreducible distinct divisors. Let $\mathbb{D}$ be the unit disc in $\mathbb{C}$, and $h : \mathbb{D} \rightarrow X$ a holomorphic map.*

*We say that h is an **orbifold map** to (X/Δ) if, for each $z \in \mathbb{D}$ such that $h(z)$ belongs to the smooth locus of $|\Delta|$, say to Δ_i , for some i , then for any nonnegative integer j which is not divisible by m_i , the j^{th} derivative of h at z , seen as a tangent vector to X at $h(z)$, is tangent to Δ_i at $h(z)$.*

We then define $d_{(X/\Delta)}$ as the largest pseudometric on X such that $h^(d_{(X/\Delta)}) \leq d_{\mathbb{D}}$, for any orbifold map from $\mathbb{D}$ to X .*

*We call $d_{(X/\Delta)}$ the **orbifold Kobayashi pseudometric of (X/Δ) .***

(I thank J.P. Demailly for supporting my impression that such a definition is meaningful, and correcting my first attempt). Remark that by the same arguments as in the classical case, this

pseudometric enjoys the same properties as when Δ is empty (in which case we just recover the usual Kobayashi pseudometric on X).

In particular, we have the following obvious refinement of the distance decreasing property of holomorphic maps:

Proposition 9.24. *Let $f : X \rightarrow Y$ be a fibration. Assume that $\Delta(f) = \Delta^-(f)$. Then:*
 $d_X \geq f^*(d_{(Y/\Delta(f))})$.

Proof: For any holomorphic map $h : \mathbb{D} \rightarrow X$, $f \circ h : \mathbb{D} \rightarrow Y$ is an orbifold map to $(Y/\Delta(f))$. (This trivial argument is in fact the motivation for the definition 9.23 given above).

Notice that for the reverse inequality, (at least when f is a special fibration) one would need to approximate arbitrarily closely holomorphic orbifold maps from $\mathbb{D}$ to Y by liftable (to X) ones. The definition of multiplicities (based on *inf*) shows that there is no local (on Y) obstruction for this.

We can now formulate more precisely Conjecture IV_H :

Conjecture 9.25. Conjecture IV'_H : *Let $c_X : X \rightarrow C(X)$ be the core of X , with c_X admissible. Then:*

- (1) $d_X = (c_X)^*(d_{(X/\Delta(c_X))})$. (In other terms: d_X should be nothing but the lift of the orbifold pseudometric of $(X/\Delta(c_X))$ by c_X).
- (2) Let (X/Δ) be a klt orbifold of general type. Then its Kobayashi pseudometric is a metric outside some proper algebraic subset $S \subset X$.

Remark that Conjecture IV'_H obviously implies Conjecture IV_H .

10. Arithmetic Aspects

10.1. Fields of Definition. We consider here the situation in which X is a projective manifold defined over the field $K \subset \mathbb{C}$, supposed to be finitely generated over $\mathbb{Q}$ (this can always be achieved).

Proposition 10.1. *Let $f : X \rightarrow Y$ be an almost holomorphic fibration. Then:*

- (a) f is defined over some finite extension K' of K .
- (b) If f is either J_X the Iitaka fibration, or r_X the rational quotient, or c_X the core of X , then f is defined over K itself.

Proof: I would like F.Bogomolov for explaining me the proof and indicating me some of the examples below.

(a) The family of fibres of f (with their reduced structure for the generic one) is then an irreducible component of the Hilbert Scheme $Hilb(X)$ of X . Let $\overline{K}$ be the algebraic closure of K . Then $G := Gal(\overline{K}/K)$ acts on the set of irreducible components of $Hilb(X)$ with finite orbits, since this action preserves the Hilbert polynomial of any component. This shows the claim.

(b) When $f = J_X$, the claim is clear, since J_X is defined by the linear system $|m.K_X|$ for some suitable integer $m > 0$, and this system is invariant under the Galois action.

In the same way, because of the maximality properties of the general fibres of r_X and c_X , we see that G preserves the component of $Hilb(X)$ parametrising the generic fibres of either r_X or c_X . More precisely: the conjugate of a rational curve (resp. a special variety) is still rational

(resp. special). (Notice that the same argument in fact applies to any fibration enjoying a similar maximality property) $\square$

10.2. Arithmetically Special Manifolds.

Definition 10.2. *The projective manifold X is said to be **Arithmetically special** (A -special, for short), if its set $X(K')$ of K' -rational points is Zariski dense in X , for some suitable finite extension K'/K . (This notion is usually called “potential density”).*

In analogy with the notion of H -speciality, we formulate the:

Conjecture III_A: X is special if and only if it is A -special.

(In the first version of the present article, I hesitated to include this conjecture; but the independent kind suggestion from B.Totaro to include it decided me. I thank him for this suggestion).

One can strengthen the above conjecture by replacing the Zariski topology by the metric topology. Examples in [C-T-S-SD] show that further field extensions may be needed (ie: $X(K')$ may be Zariski dense without being metrically dense).

Now guided by Lang’s arithmetic conjecture that a variety of general type should be “Mordellic”, and the connections he found between hyperbolicity and “Mordellicity”, we infer from the observations above in 9.1, the following definitions and conjectures:

Conjecture IV_A: Let X be a projective manifold defined over the field $K \subset \mathbb{C}$, assumed to be finitely generated over $\mathbb{Q}$.

Let $c_X : X \rightarrow C(X)$ be its core (which is defined over K , by the above remark). Let K'/K be any finite extension. Then the K' -rational points of X are mapped by c_X to a proper algebraic subset which is finite outside of the proper algebraic subset S of $C(X)$ defined in Conjecture IV’_H.

Known cases that Special implies A -special:

(a) **Curves.** For genus 0 curves, the existence of a K -rational point implies already the density of $X(K)$. For genus 1 curves, this is a special case of [L-N], for example.

(b) **Abelian varieties.** This is shown in [L-N], but is older (going back to A.Weil at least).

(c) **Unirational manifolds.** The conjecture holds, almost by definition. For rational surfaces, see [CT], especially th.5 there, as well as the references in [CT-S-SD] and [CT], for example. (The works devoted to this class of surfaces lie much deeper than what is needed for our purpose here, because the authors usually work with a **fixed** base field K).

(d) **Special elliptic surfaces with two sections.** (due to [B-T]) The base is a rational or elliptic curve. So (finitely) extending the base field, we can assume everything to be defined over K , and the two sections having a dense set of K -points. They induce then, using addition in the fibres of the elliptic fibration a dense set of K -rational points.

This is the principle of construction of [Bo-T] for a $K3$ -surface over a number field, and a dense set of K -rational points. By taking a suitable base change by an elliptic double cover of the base $\mathbb{P}^1(\mathbb{C})$, one can even get a surface X with $\kappa(X) = 1$, and $X(K)$ dense. Of course, X is special.

(Colliot-Thélène informed me that $K3$ examples with this properties are already in [SD], and with $\kappa = 1$ in [Cox]).

(e) **Quartics in $\mathbb{P}^4(\mathbb{C})$.** These are A -special if defined over a number field (by [H-T]).

(f) **Fano Manifolds, Rationally connected Manifolds.** The answer is not known (although the analogous property property is known over function fields (see 11.1 below).

Let us list some properties common to special and A -special manifolds.

First, in complete analogy with 5.3 and 9.13:

Theorem 10.3. *Let X be A -special. Then:*

- (a) *The Albanese map of X is surjective.*
- (b) *Any finite étale cover $u : \tilde{X} \rightarrow X$ of X is A -special (if X is defined over a number field K).*

Proof: (a) If not, we have a surjective map $\beta : X \rightarrow Z$, where Z is a subvariety of general type of an Abelian variety. Everything is defined over the base field K , after a finite extension. After [Fa], $Z(K)$ is however not Zariski dense. This contradicts our assumption that X is A -special.

(b) This is an immediate consequence of the Chevalley-Weil Theorem ([H-S], Ex.C7, p.292), which states that in this situation, $\tilde{X}(K')$ contains $u^{-1}(X(K))$, for a suitable finite extension K'/K . (The restriction that K is a number field can probably be removed, here).

Known cases of conjecture IV_A :

The list here is very short.

- (a) **Curves.** This is known by the solution of Mordell's conjecture in [Fa] for curves of general type and previous papers by the same author.
- (b) **Subvarieties of Abelian varieties.** Such a variety is not A -special if of general type, by [Fa]. The distribution of K -rational points is as predicted by Lang's conjectures.
- (c) **Non-special surfaces not of general type.**

Theorem 10.4. *Let X be a projective surface defined over the number field K . Assume X is not special and not of general type. Then conjecture IV_A holds for X .*

Proof: Because of the hypothesis, either X is birationally ruled over a curve of genus 2 or more, and the conclusion follows from [Fa], or X has an elliptic fibration which is of general type (in the sense of (1.9)). But then X has a finite étale cover $\tilde{X}$ which is elliptic over a curve of general type. The conclusion now follows from combining [Fa] and 10.3 (the Chevalley-Weil theorem). $\square$ (Notice that the preceding argument in fact solves, more generally, conjecture IV_A when $ess'(X) = 1$ over number fields:

Proposition 10.5. *Assume $ess'(X) = 1$, and X is defined over a number field. Then Conjecture IV_A holds for X .*

The hypothesis here means that the core of X is a fibration of gcd -general type (not just of general type). See 6.8 for this notion. It is an interesting question to know if the arguments can be adapted to the case where $ess(X) = 1$.

Remark 10.6.

- (1) The preceding result is known in various versions (see [Da], [Da-G], [CT-S-SD]. I thank D. Abramovitch and J.L. Colliot-Thélène for learning me these references).
- (2) Presumably, the conclusion extends if K is finitely generated over $\mathbb{Q}$.
- (3) Lang's connection between hyperbolicity and arithmetic properties and the reformulation above of Conjecture IV_H as Conjecture IV'_H also suggest strongly that one should be able to attach to any projective orbifold (Y/Δ) defined over K the notion of its K -rational points $(Y/\Delta)(K)$.

Such a definition can actually be given quite naturally, in complete analogy with the hyperbolicity and function field cases considered here, as explained to me by E. Peyre ([Pey]).

An important property is that if $f : X \rightarrow Y$ is a fibration defined over K , then $f(X(K)) \subset (Y/\Delta(f))(K)$, the latter being the set of K -rational points of the base orbifold $(Y/\Delta(f))$ of f .

Then Conjecture IV_A above can be reformulated as the assertion that an orbifold of general type defined over K does not have a Zariski dense set of K -rational points.

Conjecture IV'_A : Let $c_X : X \rightarrow C(X)$ be an admissible representative of the core of X , defined over K , finitely generated over $\mathbb{Q}$.

If K'/K is any finite extension $S \subset C(X)$ is defined as in conjecture IV'_H , then $(C(X)/\Delta(c_X))(K')$ has only finitely many points outside of S (and contains $c_X(X(K'))$).

10.3. A Related Conjecture of Abramovitch and Colliot-Thélène. A conjecture related to conjecture III_A is formulated in [H-T], and attributed to D. Abramovitch and J.L. Colliot-Thélène (I thank J.L. Colliot-Thélène for learning me this reference) : it claims that if X is what we called w -special and defined over a number field, then X is A -special (“potentially dense”, there), and conversely.

This conjecture thus coincides with conjecture III_A if the properties of being special and w -special coincide.

Although no example of a w -special, but not special manifold is known today, it seems plausible that the two properties are distinct (special is anyway more restrictive).

The difference in the two conjectures is that if there exists an X (defined over a number field) which is w -special, but not special, it should be A -special by [H-T], but not by IV_A , which would imply that its K' -rational points should lie in finitely many fibres of c_X , outside of $c_X^{-1}(S)$.

11. The Function Field Case

11.1. The (Complex) Function Field Case. The following extremely interesting function field variant of Conjectures III_A and IV_A above was suggested to me by P. Eyssidieux. They are included here with his kind authorisation.

(One could of course consider other fields than $\mathbb{C}$ as well).

We need first some conventions.

Conventions. Let $f : X \rightarrow C$ be a fibration onto a complex projective curve C , with X complex projective, smooth connected. Let $S^0(f) \subset \text{Chow}(X)$ be the Zariski open subset consisting of holomorphic sections of f , and let $S(f)$ be its closure in $\text{Chow}(X)$, the Chow variety of X .

We also denote by $S^0(X) \subset X$ the union of all images of sections of f , and by $S(X)$ the Zariski closure of $S^0(X)$ in X . We say that an irreducible component Σ of $S^0(f)$ is X -covering if the union of all images of sections in Σ is Zariski dense in X (and so covers a nonempty Zariski open subset of X).

Let finally $b : C' \rightarrow C$ be any finite covering (possibly ramified), with C' a smooth connected projective curve. We then let $f' : X' \rightarrow C'$ be deduced from f by the base change b . We define also $S(f')$, $S(X')$,... as before, with f' in place of f .

The arithmetic interpretation is that if K is the (non algebraically closed) function field of C (ie: its field of meromorphic functions), then X is a model of the K -variety X_K defined over K by

the function field of X . Moreover, the sections of f are exactly the K -rational points $X(K)$ of X_K ; and $X(K)$ is Zariski dense in X_K if and only if so is $S^0(X)$, or equivalently: $S(X) = X$.

The finite extensions K'/K are exactly the ones given by the coverings $b : C' \rightarrow C$ above.

We shall say that $f : X \rightarrow C$ is **split** if there exists a manifold F and a birational map $\sigma : F \times C' \rightarrow X$ over C' , after some suitable base change b . This is (essentially) Lang's terminology in [La 2].

We need to introduce the orbifold version of the preceding notions:

An orbifold (X/Δ) over C will be just a fibration $f : X \rightarrow C$ as above, together with an orbifold structure on X . The notions of Kodaira dimension, canonical bundle, general type,... are defined by restriction to the general fibre.

If $f : X \rightarrow C$ and $g : Y \rightarrow C$ are fibrations, a fibration $h : X \dashrightarrow Y$ over C is the usual h such that $g \circ h = f$.

When h is holomorphic, we can naturally define its base orbifold (an orbifold structure on Y) in the expected way. This leads to the notion of Kodaira dimension of h , denoted $\kappa(Y, f)/C$.

A special fibration $f : X \rightarrow C$ is thus defined in the same way as we already did. By the existence of a relative core, this is the same as having a special general fibre.

In this context, it is maybe useful to define the sections of an orbifold (X/Δ) over C . These are simply the sections of f which meet each component Δ_j of Δ at order at least m_j at each point where the section meets Δ_j , if $\Delta = \sum_{j \in J} (1 - 1/m_j) \cdot \Delta_j$.

We say that (X/Δ) is **split** if it becomes birationally trivial after some base change b as in the non-orbifold version. (Birationally trivial means that (X/Δ) is birational to $(F/D) \times C$ over C , for some "constant" orbifold (F/D)).

Definition 11.1. *We say that the K -variety X_K is **special** (resp. **rationally connected**; resp. **an abelian variety**; resp. **of general type**...) if so is the general fibre of f .*

Let $f = g \circ h$ be factorisation of f by fibrations $h : X \rightarrow Y$ and $g : Y \rightarrow C$. We say that this factorisation is the K -core (resp. the K -rational quotient; resp. the K -Albanese map) if it induces the corresponding reduction on the general fibre of f . These factorisations thus exist (by 12.27) and analogous results for the other cases), are uniquely defined, and moreover compatible with the base changes b .

Because of the split case, the formulation of the function field version of conjectures III_A and IV_A is not exactly the same.

To formulate them we need to introduce the K -core $f = g_f \circ c_f$ of f (see 12.27). We thus have a factorisation of f by $c_f : X \rightarrow C(f)$ and $g_f : C(f) \rightarrow C$.

Conjecture 11.2. (Conjecture III_{FF}) *The K' -rational points of X_K are Zariski dense in X , for some finite extension K'/K if and only if $g_f : (C(f)/\Delta(c(f))) \rightarrow C$ is split. In particular, if X_K is special, $X(K')$ is Zariski dense in X , for a suitable finite extension K'/K . (One might say, in analogy with the terminology introduced in 9.1 and 10.2, that they are FF-special (FF standing for: function field)).*

Geometrically, this just means that $S(X') = X'$ if and only if $C(f)$ is birationally a product over C (after a preliminary covering of C , maybe).

Conjecture 11.3. (Conjecture IV_{FF}) Assume that $g_f : (C(f)/\Delta(c(f))) \rightarrow C$ is not split. There exists a Zariski closed subset $T \subset C(f)$, $T \neq C(f)$ such that for any finite extension K'/K , $c_f(S(X'))$ is contained in T and a finite union of images of sections of g_f .

Remark 11.4. If g_f is split, one can consider also only the sections which are not constant to then recover exactly the same formulations as in the arithmetic case.

Known cases when X_K is special. One wants to know if $X(K')$ becomes Zariski dense for a suitable finite extension K'/K .

(a) **Curves.** For rational curves, this is clear, because such surfaces X are birationally ruled. For elliptic fibrations, it is a special case of the next case (alternatively, by a suitable base change, construct 2 sections; the first one is used as the zero section to define a relative group law; then choose the second section so as to have infinite order. This is an immediate construction).

(b) **Abelian varieties.** This is shown in a more general form in [L-N].

(c) **Rationally connected varieties.**

They are FF -special (for “function-field-special”). This immediately follows from the glueing lemma of [Ko-Mi-Mo], which implies that if the generic fibre of $f : X \rightarrow C$ is rationally connected, and if f has a section, then any finite set of X mapped injectively to C is contained in a section of f .

Notice that [G-H-S] announces that f itself has a section, so that no field extension is needed in that case.

(d) **Surfaces.** The answer is known, except for $K3$ -surfaces as fibres:

Theorem 11.5. Let $f : X \rightarrow C$ be a fibration with its generic fibre a special, but not $K3$, surface. Then $S(X') = X'$ for a suitable finite covering $b : C' \rightarrow C$.

The proof will be given elsewhere. The case of elliptic $K3$ surfaces is true and immediate. It is certainly worth investigating the case of general $K3$'s.

Known cases when X_K is not special.

The list is then much more limited. It is in fact restricted to cases when X_K is of general type, and where conjecture IV_{FF} reduces to the function field case of Lang's conjectures.

(a) **Curves.** This case has been settled by various approaches by H.Grauert [Gr], Y.Manin [Man] and A.N.Parshin [Par].

(b) **Subvarieties of general type of Abelian varieties.** This the case where the general fibre F of f is of general type and mapped birationally into its Albanese variety by its Albanese map. We now turn to another question, quite natural in our context, but for which we don't know any reference.

11.2. Split and “Dominated” Fibrations: the Rational Quotient. We shall make here some easy observations on the **non-orbifold versions**.

It would be interesting to extend them to the orbifold case as well.

Definition 11.6. The fibration $f : X \rightarrow C$ is said to be **dominated** if there exists a projective manifold F' and a dominant rational map $g : F' \times C' \rightarrow X'$ over C' , for some suitable $b : C' \rightarrow C$.

If the above map g can be chosen to be regular (resp. isomorphic above the generic point of C), we say that f is **regularly dominated** (resp. **isotrivial**).

Remark that if f is dominated, the manifold F' can be chosen to be of the same dimension as F , the generic fibre of f , so that g is generically finite. And remark also that a split fibration is dominated.

Our objective now is to introduce Conjecture III_{SD} below, which says that the obstruction to reverse implication lies in the rational quotient of f .

The relevance of notion of “dominatedness” to our context is the following:

Proposition 11.7. *The following properties are equivalent, for a fibration $f : X \rightarrow C$:*

- (a) f is dominated.
- (b) $S^0(X') = X'$, for a suitable $b : C' \rightarrow C$. (See above for the definition of $S^0(X')$).
- (c) There exists an irreducible component T' of $S(f')$ which is X' -covering, for some suitable $b : C' \rightarrow C$. (See 12.)

Proof: Obviously, (a) implies (b). To show that (b) implies (c), use the countability at infinity of the Chow-Scheme of X . To show that (c) implies (a), take $T' := F'$, and use the (surjective) evaluation map of the incidence graph G , say, of the algebraic family of cycles parametrised by T' . The conclusion is implied by the fact that G is bimeromorphic to $C \times T'$, because the generic point of T' parametrises a section of f .

Conjecture 11.8. (Conjecture III_{SD})

(i) Let $r_K : X \rightarrow R(X_K)$ be the rational quotient of the fibration $f : X \rightarrow C$. Then X_K is dominated if and only if $R(X_K)$ is split.

A weaker form (shown below to be equivalent) is:

(ii) Assume X_K is dominated, but not split. Then X_K is uniruled (ie: so is F).

This means in particular that X_K is dominated if rationally connected, and that it is split if it is dominated and not uniruled.

We shall show several cases where the above conjecture holds. First, the “if” part:

Theorem 11.9. *If $R(X_K)$ is split, then X_K is dominated.*

A special case (when $R(X_K) = C$) is:

Corollary 11.10. *Assume X_K is rationally connected. Then it is dominated.*

It would be interesting to have concrete examples when F is a three-dimensional cubic or quartic. For example: the above result asserts that one can uniformly dominate one-dimensional families of such manifolds, although they are birationally distinct.

Proof (of 11.9): Assume $R(X) = C \times R$ (after finite base change, if needed). Let $r \in R$ be generic, and let $r'_K : X_r := r_K^{-1}(C \times \{r\}) \rightarrow (C \times \{r\})$ be the restriction of r_K to X_r . This is a rationally connected fibration, and so by [G-H-S] and [Ko-Mi-Mo], it is dominated. Thus so is f , using characterisation (b) above of dominatedness $\square$

Theorem 11.11. *If the weaker form (ii) of conjecture III_{SD} holds, the stronger form (i) also holds.*

Proof: The proof is actually a direct application of 3.20, or of [G-H-S], which asserts the equivalence of rational connectedness and rational generatedness. (This last notion is defined as follows:

X is rationally generated if, for any fibration $h : X \rightarrow Y$, Y is uniruled). Indeed: since X_K is dominated, so is $R(X_K)$. If $R(X_K)$ is not split, it is uniruled by . But [G-H-S] for details) in particular says that $R(X_K)$ is not uniruled.

If, conversely, $R(X_K)$ is split, we deduce from 11.9 above that X_K is dominated $\square$

We now give some positive answers to the conjecture III_{SD} , (ii).

Proposition 11.12. *Assume that through the generic point of X' there exists a positive-dimensional family of sections of $f' : X' \rightarrow C'$. Then X_K is uniruled.*

This is a simple application of Mori's bend-and-break technique. Indeed:

Proof: This is a simple application of Mori's bend-and-break technique. Indeed: let $x \in X$ be generic with a positive-dimensional family of sections through it. By Mori's techniques (no characteristic $p > 0$ is needed, here), there is a rational curve Γ in X through x . If the base C is not rational, Γ is vertical (ie: contained in the fibre of f through x). We can reduce to this case by a base change $b : C' \rightarrow C$, with C' non-rational, because sections of f lift to sections of f' . $\square$

So the remaining case is when X is dominated, but the set of sections through the general point of any X' is at most countable, not empty.

Proposition 11.13. *Assume X_K is regularly dominated. Then X_K is isotrivial.*

This is essentially (up to the terminology) shown in [Mw].

Corollary 11.14. *Assume F (the generic fibre of f) is an absolutely minimal model (in the sense that any birational rational map $r : F_0 \rightarrow F$, with F_0 smooth, is regular). Then: X_K is isotrivial if dominated.*

Example 11.15. *The above condition is satisfied if F is an abelian variety, or a minimal surface with $\kappa(F) \geq 0$.*

Another important case is given by:

Proposition 11.16. *Assume F is of general type. If X_K is dominated, it is split.*

This is shown in [Mae]. Observe that conjecture III_{SD} , (ii) consists in extending this to the case where F is not uniruled. Already the (conjecturally equivalent) case where $\kappa(F) \geq 0$ would be of interest.

12. Appendix: Geometric quotients

We now turn to the exposition of various techniques of construction of quotients for equivalence relations of geometric origin.

These techniques have been introduced in [C80,81] and have found many subsequent applications ([C92], [Ko-Mi-Mo], [C94], [C01] among others).

They are based on fundamental properness properties of the Chow-Scheme $\mathcal{C}(X)$ of X (see [Ba] if X is not projective) if $X \in \mathcal{C}$, discovered in [Lieb] (see [Fu] for the Hilbert-Douady space case), further properties having been shown in [C80].

When X is projective, these properties reduce to the well-known projectivity of the irreducible components of $\mathcal{C}(X)$ of X

The cycle of X naturally parametrised by $t \in T$ is denoted by V_t . Conversely, if V is a subvariety of X (ie: irreducible reduced), we denote by $[V]$ (or sometimes simply V) the corresponding point $t \in \mathcal{C}(X)$.

We need some preliminary definitions and facts.

12.1. Covering Families. Let X be a compact connected normal analytic space (not necessarily in $\mathcal{C}$).

We shall consider finite or countable unions $T \subset \mathcal{C}(X)$ of compact irreducible subvarieties $T_i \subset \mathcal{C}(X)$, such that:

(a) VV_t is irreducible for t generic in T .

(b) X is the union of all V_t 's, for $t \in T$.

(Equivalently: if $V \subset X \times T$ is the incidence graph of the family $(V_t)_{t \in T}$, and $q : V \rightarrow T$ and $p : V \rightarrow X$ the restrictions to V of the projections, then p is surjective and for any index i , q_i has generic fibres which are irreducible, where V_i , q_i , and p_i denote the incidence graph of the family T_i , and the restrictions of p and q to V_i respectively).

We say then that $T \subset \mathcal{C}(X)$ is a **covering family** of X . Notice that by Baire's Category Theorem, at least one of the families $T_i \subset \mathcal{C}(X)$ has to be a covering family of X .

Because the maps q_i are equidimensional, we have the following important:

Proposition 12.1. *If T is a covering family of X , then:*

- a. V_t is connected, for every $t \in T$.
- b. V_i is irreducible, for every i . More generally:
- c. If V'_i is deduced from $q_i : V_i \rightarrow T_i$ by any surjective base change by $\beta : T'_i \rightarrow T_i$, with T'_i irreducible, then V'_i is irreducible, too.

Sometimes, we shall denote by V not the incidence graph of the family T , but its normalisation, or a smooth model of it, without further notice.

Remark 12.2. : *All results in this appendix hold, together with their proofs without any change if X is arbitrary, irreducible (not necessarily compact), provided the maps q_i, p_i are all assumed to be proper (so the T_i 's are not necessarily compact either). This was observed and used in [C94].*

(14.2) T-Equivalence (see also [C81])

Let $T \subset \mathcal{C}(X)$ be a covering family of X . Let $R(T)$ be the equivalence relation on X which is generated by the symmetric reflexive binary relation $R^1(T)$ on X for which $x, x' \in X$ are related if and only if there exists $t \in T$ such that V_t contains x and x' .

Thus $x, x' \in X$ are $R(T)$ -equivalent iff they can be joined by (ie: are contained in) a **T -chain**. This is by definition a **connected** union of finitely many of the V_t 's.

We say that X is **T -connected** if $R(T)$ has a single equivalence class (ie: any two points of X can be connected by some T -chain).

For these definitions, we do not assume that T is irreducible. We only assume that each irreducible component of T is compact, and that T has at most countably many such components.

Observe that $R^1(T) \subset X \times X$ has the following description: $R^1(T) = (p \times p)(V \times_T V)$, with obvious notations. From this and 12.1, we conclude:

Corollary 12.3. *If T is a covering family of X , then for any i , $R^1(T_i)$ is irreducible.*

The fundamental result on T -equivalence is now the following:

Theorem 12.4. *([C81]) Let $T \subset \mathcal{C}(X)$ be a covering family of X . Assume that X is normal. There exists a fibration $q_T : X \rightarrow X_T$ such that its general fibre is an equivalence class for $R(T)$. Of course, this map is unique, up to equivalence. Moreover, q_T is almost holomorphic. The map q_T is called the **T -quotient** of X .*

Remark 12.5. *If T has only finitely many irreducible components, the word “general” can be replaced by “generic” in 12.4. See below.*

It is not necessary to assume that $X \in \mathcal{C}$ in 12.4.

Remark 12.6.

The above result is shown in [C6] only when T is irreducible. The general case stated here can however be easily reduced to this special case by arguments sketched below in 12.3.

See also [C94] for a more general version (but the proof goes through) in which X is not assumed to be compact, but p_i is assumed to be only proper. This version is used to construct Γ -reductions (also called Shafarevitch maps). These T -quotients appear in many other problems.

Finally, the normality assumption is essential, as shown by examples given in [C81].

Theorem 12.4 is exposed in [Ko1] in the algebraic context, via an approach slightly different from (and no appropriate quotation of) the original one.

A careful, corrected exposition along this line is given in [De].

The proof of theorem 12.4 will be done in two steps: first the proof of the case where T has finitely many components, each covering; second: the easy reduction to this special case.

12.2. Construction of the Quotient for an Irreducible Covering Family. We shall prove here Theorem 12.4 when T is irreducible. The proof is exactly the same as in [C81], except for a slight simplification at one point, which shortens the exposition. This simplification consists in replacing T by T' in step 2. below.

Let us remark that the proof works also if T is not necessarily irreducible, provided all components T_i of T are covering.

We thus start with a diagram:

$$\begin{array}{ccc} V & \xrightarrow{p} & X \\ q \downarrow & & \\ T & & \end{array}$$

in which V, T, X are irreducible compact, p, q surjective, and q is equidimensional with generic fibre irreducible. Also X is normal.

Because T is fixed, we delete T from the notations introduced in 12.1, obtaining R, R^1 instead of $R(T), R^1(T)$.

We define an **n -chain (of T)** as the connected union of n of the V_t 's.

We say that $x, x' \in X$ are **n -equivalent (for T)** if they are connected by (ie: contained in) some n -chain of T .

We denote by R^n (or $R^n(T)$) the symmetric set of $X \times X$ consisting of pairs $x, x' \in X$ which are n -equivalent. When $n = 0$ (resp. $n = \infty$), we define R^n as being Δ_X , the diagonal of X (resp. $R = \cup_{n \geq 0} R^n$).

The sequence R^n is of course increasing for the inclusion.

Lemma 12.7. *For any $n \geq 1$, one has:*

1. $R^{n+1} = (p)'((q')^{-1}((p)')^{-1}(R^n))$, where $(p)'$ and $(q)'$: $V \times_T V \rightarrow X = \Delta_X$ denote the restrictions to $V \times_T V$ of $(p \times p) : (V \times V) \rightarrow (X \times X)$ and $(q \times q)$ respectively.
2. $R^n \subset (X \times X)$ is Zariski closed.
3. R^1 is irreducible. (see 12.1).

Proof: Easy. See [C81] for details $\square$

Definition 12.8. *We shall now say that an irreducible component of R^n is **significant** if it contains Δ_X .*

We note $R_n \subset R^n$ the union of the significant irreducible components of R^n . Thus $R_n \subset X \times X$ is Zariski closed, and R^n has only finitely many significant components, for any n . Notice that $R_1 = R^1$ is irreducible (if T is).

We also note $d_n(T) := d_n := \dim(R_n)$, for every $n \geq 0$. This is an increasing, bounded sequence of integers. So it stabilises at some n_0 .

Definition 12.9. *We shall say that the irreducible covering family T of X is **stationary** if:*

- a. p is a modification
- b. $d_2 = d_1$

Theorem 12.4 is proved in two steps, corresponding to the following two propositions:

Proposition 12.10. *Let T be an irreducible covering family of X , normal. There exists a **stationary** irreducible covering family T' of X such that $R(T) = R(T')$.*

Proposition 12.11 will be proved below.

Proposition 12.11. *Let T be an irreducible stationary covering family of X , normal. Then:*

- a. $h := q \circ (p^{-1}) : X \dashrightarrow T$ is an almost holomorphic fibration.
- b. $h : X \dashrightarrow T$ is the T -quotient of X .

Proof (of 12.11:) The second assertion is an immediate consequence of the first, because h is then holomorphic near each of its regular fibres (the ones not meeting the indeterminacy locus I_h of h). So that each such fibre is then an equivalence class for $R(T) = R$.

Everything is clear from the definitions and assumptions in the first assertion, except for the fact that h is almost holomorphic. We check this by contradiction, assuming that each fibre $F = V_t$ of h meets I_h . This implies that F also meets J , the indeterminacy locus of p^{-1} . Let $x \in J \cap F$. Because X is normal, $p^{-1}(x') := W'$ is positive dimensional at each of its points; in particular at $x' := (t, x)$. Let W be an irreducible component of W' containing x' .

Let $S := p(q^{-1}(q(W))) \subset X$, and let $y \in F = V_t$ be generic. Then $S \times y \subset R^2(T)$ (because (x, y) and (s, x) are in $R^1(T)$, if $s \in S$, by construction. More precisely, if $F = V_t$ is irreducible reduced, then $S \times y \subset R_2(T)$, because S is then irreducible and $t \in T$ generic.

We now get that $d_2 \geq d + m$, if $d := \dim(S)$, and $m := \dim(X)$, while $d_1 = f + m$, if $f := \dim(F) = (\dim(V) - m)$. To get a contradiction, it is thus sufficient to show that $d > f$. But this is obvious because: $\dim(S) = \dim(p(q(q^{-1}(W))) = \dim(q(q^{-1}(W))) = \dim(W') + f > f$. The second equality comes from the fact that p is a modification, and that F is not contained in J , but is contained in $q(q^{-1}(W))$. $\square$.

We now start the:

Proof (of 12.10): We shall denote, for any $A \subset X \times X$ by r' (resp. r'') the restriction to A of the first (resp. second) projection of $X \times X$ to X , and by $A'(x)$ (resp. $A''(x)$) the fibre of r' (resp. r'') above $x \in X$.

Let $s : X \times X \rightarrow X \times X$ be the involution exchanging the factors. We say that A is **symmetric** if $s(A) = A$. In this case, $A'(x) = A''(x), \forall x \in X$, and is simply denoted $A(x)$.

The main step in the proof of 12.10 is:

Lemma 12.12. *Let $n \geq 0$ be such that $d_n = d_{2n}$. Then:*

- a. R_n is symmetric and irreducible.
- b. $R_n(x)$ is irreducible and locally irreducible at (x, x) , for x generic in X .

Before we show 12.12, let us prove that it implies proposition 12.10:

Let $r' : R_n \rightarrow X$ be the first projection. By 12.12, it is a fibration. Let $g : X \dashrightarrow \mathcal{C}(X)$ be the associated map, sending $x \in X$ generic to $R_n(x)$. Let $T' \subset \mathcal{C}(X)$ be the image of g . This is an irreducible covering family of x .

It thus only remains to observe the following two properties:

- a. $R(T) = R(T')$
- b. The family T' is stationary

Let us show [a.]: For x generic in X , $R_n(x) = (R')^1(x)$, by definition, with $R' := R(T')$. Because $(R')^1$ is irreducible (by 12.1), we conclude that $R^1(T') = R_n(T) \subset R(T)$. Thus $R(T') \subset R(T)$.

In the opposite direction, because $R^1(T)$ is irreducible and contains the diagonal, $R^1(T) \subset R_n(T) = R^1(T')$. Thus: $R(T) \subset R(T')$.

(Observe this argument still works if T consists of finitely (or infinitely!) many components T_i , each of which is covering (so that $R^1(T_i)$ contains the diagonal. But it does not work in general).

Let us next show [b.]: Because $R^1(T') = R_n(T)$, we have: $R^2(T') \subset R^{2n}(T)$. Thus also: $R_2(T') \subset R_{2n}(T)$. So: $d_2(T') \leq d_{2n} = d_n = d_1(T')$. Hence [b.]. $\square$

To conclude the proof of Theorem 12.4, it thus only remains to give the:

Proof (of Lemma 12.12): We first prove by contradiction that $R_n(x)$ is irreducible at (x, x) , for $x \in X$ generic.

Otherwise, R_n had two distinct local irreducible components S_1, S_2 at (x, x) , with S_1 say of dimension d_n ; and $S_1(x), S_2(x)$ are irreducible at (x', x') , for x' close to X , by Lemma 12.13 below (applied to each local irreducible component of R_n).

We have a natural holomorphic map: $h : W := S_1(x) \times_X s(S_2) \rightarrow R_{2n}(x)$ defined by sending $((x, y), (y, z))$ to (x, z) . Shrinking suitably the domain W of this map, we can assume that it is irreducible, because the generic fibres of W over X are irreducible. But now $h(W)$ contains

$S_1(x) \cap U$ (take $z = x$ and $S_2(x) \cap U$ (take $y = x$), if U is a sufficiently small open neighborhood of x in X .

Because $S_2(x)$ is not contained in $S_1(x)$, by hypothesis, the rank of h at the generic point of W is thus strictly larger than: $\dim(S_1) - m = d_n - m$, if $m := \dim(X)$. Because this rank is at most $d_{2n} - m$, this implies that $d_{2n} > d_n$, contradicting our assumptions.

The others assertions of 12.12 are immediate consequences of lemma 12.13 below. $\square$.

To complete the proof of Theorem 12.4 for T irreducible, it thus simply remain to show the:

Lemma 12.13. *Let $g : V \rightarrow W$ be a surjective holomorphic map between complex spaces, with $\sigma : W \rightarrow V$ a holomorphic section. Let $W' := \sigma(W)$.*

If V is locally irreducible at the generic point of W' , then for $w \in W$ generic:

- a. *The g -fibre V_w over w is locally irreducible at $w' := \sigma(w)$.*
- b. *V_w is irreducible, if g is proper.*

Proof: We shall give, by lack of suitable reference a proof based on desingularisation. The only advantage is its shortness. Elementary algebro-geometric proofs were longer (see [C81] for an analytic proof).

Proof of [a.]: let $d : V' \rightarrow V$ be a smooth model of V , and $g' := g \circ d : V' \rightarrow W$. We can assume W to be smooth connected. The fibres of d over W' are thus connected, by assumption (and Zariski's Main theorem). Let $W'' := d^{-1}(W') \subset V'$. The rank of g' is then maximal, equal to $m := \dim(W)$, at each $w'' \in W^* \subset W''$, where W^* is Zariski open and contains $W'' \cap ((g')^{-1}(U))$, where U is a dense Zariski open subset of W . This holds true by standard Zariski lower semi-continuity properties of the rank, and Bertini theorem.

For $w \in W$ generic, we conclude that the fibre V'_w of g' over w is connected and smooth near W_w'' . Thus $V_w = d(V'_w)$ is irreducible near w' . Hence [a.].

Proof of [b.]: Let V'_w be a smooth generic fibre of g' . We saw that $W_w'' := d^{-1}(w')$ is connected. It is thus contained in a unique connected component $\bar{w}$ of V'_w . Let $g' := h \circ \bar{g}$ be the Stein factorisation of g' , with $\bar{g} : V' \rightarrow \bar{W}$ connected and $h : \bar{W} \rightarrow W$ finite. Now the previous observation defines a natural meromorphic section $\bar{h}$ of h , which sends $w \in W$ generic to $\bar{w}$. This shows [b.]. $\square$

The above proof of Theorem 12.4 easily gives an estimate of the length of the T -chains connecting two general T -equivalent points of X (see [Ko-Mi-Mo], and [De] for similar results):

Proposition 12.14. *Let T be an irreducible covering family of X , normal. Let $q_T : X \rightarrow X_T$ be the T -quotient of X , and F a regular (ie: contained in the regular locus of q_T) fibre of q_T . Then any two points of F are joined by a T -chain of length at most $l := 2^{(f-r)}$, where $r := \dim(V_t)$, and $f := \dim(X) - \dim(X_T)$ is the dimension of a generic fibre of q_T .*

In particular: $l \leq 2^{f-1}$, unless T is the family of points of X .

Actually, the statement holds for **any** Chow-theoretic fibre F of q_T , as the proof shows.

Proof: Assume the sequence $d_1, d_2, d_4, \dots, d_{2^k}$ is strictly increasing, while $d_{2^k} = d_{2^{k+1}}$. Thus: $d_{2^k} \geq d_1 + k$. From the proof of 12.4, we see that $d_1 = m + r$, while $d_{2^k} = m + f$, if $m := \dim(X)$. Thus: $m + f = d_{2^k} \geq d_1 + k = m + r + k$, and $k \leq (f - r)$. Hence the claim, since by construction, two points of F are connected by a T -chain of length 2^k . $\square$.

Remark 12.15.

The proofs given here work also when T has all of its components T_i covering, if r denotes the infimum of the corresponding r_i 's. So, in this case, we always get: $l \leq 2^{f-1}$. The estimate is then in this case slightly sharper than the one ($l \leq 2^f - 1$) given in loc.cit.

The estimate $l \leq 2^f - 1$ can easily be deduced but apparently not improved (see section 12.3 below) in the general case from 12.14, and the reduction process described in the next section.

12.3. Reduction to the case where T is irreducible. We show how to reduce the proof of Theorem 12.4 to the case where T is irreducible.

We shall consider the set $\mathcal{F}(X, T)$ of almost holomorphic fibrations $f : X \dashrightarrow Y$ which are **subordinate to T** . This means, by definition, that $R(f) := R(Y) \subset R(T)$. Here Y is seen as the covering family parametrised by the family of fibres of f .

Notice that the identity map of X is in $\mathcal{F}(X, T)$.

Let $T' := T_i$ be a component of T , and V', p', q' the data associated to T' , as in 12.1.

Let $f \in \mathcal{F}(X, T)$. We say that T_i is **f -covering** if $f \circ p' : V' \rightarrow Y$ is surjective.

The main step of the reduction process is contained in:

Lemma 12.16. *Assume that T' is f -covering for some f subordinate to T . Then either:*

1. $R(T') \subset R(f)$, or:
2. *There exists a fibration $g : Y \dashrightarrow Y'$ with $\dim(Y') < \dim(Y)$, and such that $f' := g \circ f : X \dashrightarrow Y'$ is almost holomorphic and subordinate to T . (Moreover: $R(T') \subset R(f')$).*

Before proving 12.16, we show how it implies the existence of q_T in the general case:

□ Take $f : X \dashrightarrow Y$ be in $\mathcal{F}(X, T)$, and such that $\dim(Y)$ is minimum. Let J be the set of i 's such that T_i is not f -covering. Let T' be the union of all components T_i of T which are f -covering.

From the preceding lemma, we see that $R(T_i) \subset R(f)$ for each $i \notin J$. For a general fibre F of f , we thus see that F is saturated for $R(T')$, and moreover that $R(T') = R(T)$ on this fibre, because the T_i 's are not f -covering if $i \in J$. Thus F is saturated for $R(T)$. Finally, because f is subordinate to T , we see that F is a single $R(T)$ equivalence class □

Proof of 12.16: Let thus T' be f -covering, and assume that $R(T')$ is not contained in $R(f)$. So $\dim(V_{t'}) > 0$ if t' is generic in T' .

In this case, define for $t' \in T'$ generic the new reduced cycle $V_{t'}' \subset X$ by: $V_{t'}' := (f)^{-1}(f(V_{t'}))$. This cycle is irreducible, because so are $V_{t'}$ and the generic fibre of f . Let now $T'' \subset \mathcal{C}(X)$ be the closure in $\mathcal{C}(X)$ of the points parametrising the cycles $V_{t'}'$ so obtained.

It is obvious to check that T'' is an irreducible covering family of X , and that we have a natural factorisation map: $g : Y \rightarrow Y' := X_{T''}$ such that $f' = g \circ f$ if $f' : X \rightarrow Y'$ is the T'' -quotient of X . Obviously: $\dim(Y') < \dim(Y)$ because: $\dim(V_{t'}) > 0$ if t' is generic in T' .

Moreover, we have an inclusion: $R^1(T'') \subset R^1(T)$, because $R(T'')$ is generated by $R(f)$ and $R(T')$. This inclusion extends to $R(T'') \subset R(T)$, so that f' is subordinate to T , as claimed.

Let us show that $R(T'')$ is generated by $R(f)$ and $R(T')$: The inclusions of $R^1(f)$ and $R^1(T')$ in $R^1(T'')$ are obvious, by construction. So that $R(f)$ and $R(T')$ are contained in $R(f')$. The reverse direction is immediately given by the:

Lemma 12.17. 1. $R^1(T'')$ is the main component of $(f \times f)^{-1}((f \times f)(R^1(T')))$.

2. $R^1(T) \subset R^2(S)$ if S is the covering family with the two irreducible components Y and T' .

Proof: The first assertion follows immediately from the definition of the generic V'_μ and the fact that $R^1(T''')$ is irreducible, while the second follows from the first and Lemma 12.7. $\square$.

Remark 12.18. ([De],[Ko-Mi-Mo]) Any two points of any Chow-theoretic fibre F' of $q : X \dashrightarrow X_T$ can be connected by a T -chain of length at most: $l' \leq 2^b - 1$, if $b = \dim(X) - \dim(X_T)$ is the dimension of F' .

Proof: By the reduction process just described above, and using the same notations, we just need to show that if this holds for f in place of q_T , this holds also for f' , so that $R(f')$ is generated by $R(f)$ and $R(T')$, T' an irreducible f -covering family of X .

By 12.14, to points of any fibre (Chow-theoretic, as always) F' of f' are connected by a T''' -chain of length $l'' \leq 2^{b'-r'} \leq 2^{b'-b-1}$, where $b' := \dim(F')$, $r' := \dim(V'_\mu) \geq b + 1$, with $b := \dim(F)$, F any fibre of f . By the definition of V'_μ , we immediately see that any two of its points are connected by a T -chain of length at most $(2l + 1)$, if two points of any fibre of can be connected by a T -chain of length l .

By the hypothesis on f , we can take $l := (2^b - 1)$. So that any two points of any fibre of f' can be connected by a T -chain of length at most $l' \leq l'' \cdot (2l + 1) \leq 2^{b'-b-1} \cdot (2 \cdot (2^b - 1) + 1) \leq (2^{b'} - 1)$, as claimed $\square$.

12.4. Zariski Regularity.

Definition 12.19. Let $A \subset S$, where S is a complex space. We say that A is **Z-regular (in S)** if for each irreducible Zariski closed subset T of S , $A \cap T$ either contains the general point of T , or is contained in a countable union of Zariski closed proper subsets of T . (In which case we say that A is of first Zariski category in T).

There are many examples of this situation (actually it rather seems that counterexamples are unnatural in algebraic or analytic geometry):

Theorem 12.20. Let $X \in \mathcal{C}$. Let $A_P(X) = A_P \subset \mathcal{C}(X)$ be the set of points t parametrising the set of irreducible reduced subvarieties V_t of X possessing a certain property P . (By convention, points of X are always supposed to posses P).

Then $A_P(X) = A_P$ is Z-regular in $\mathcal{C}(X)$ if P is any one of the following properties.

1. V_t is special;
2. V_t is a (possibly singular) rational curve.
3. V_t has algebraic dimension at least d , a given nonnegative integer.
4. $\pi_1(\hat{V}_t)_X \in \mathcal{G}$, where $\mathcal{G}$ is a class of (isomorphism classes) of groups, and the first term is the image in $\pi_1(X)$ of the fundamental group of the normalisation of V_t .

Proof: For 1 (resp. 3 ; 4) see corollary 2.37 and proposition 12.21 below (resp. [C80]; [C01]). Assertion (2) is easy.

Proposition 12.21. Let P be a property such that if $f : X \rightarrow Y$ is any fibration such that $X \in \mathcal{C}$, then the set of irreducible fibres of f having P either contains the general point of Y , or is of first Zariski category in Y .

Then: for any $X \in \mathcal{C}$, $A_P(X)$ is Z -regular.

Proof: Let $T \subset \mathcal{C}(X)$ be irreducible Zariski closed. Let $V = V_T$ be the incidence graph of the family T , with projections $p : V \rightarrow X$ and $q : V \rightarrow T$. By [C80], $V \in \mathcal{C}$. Thus: $A_P(X) \cap T$ either contains the general point of T , or is of first Zariski category in T , by the hypothesis made on P $\square$.

Proposition 12.22. *Let $A \subset S$ be Z -regular. There exists a countable or finite family of Zariski closed irreducible subsets S_i of S such that:*

1. *For each i , $A_i := A \cap S_i$ contains the general point of S_i .*
2. *A is the union of the A_i 's. The family S_i is moreover unique if it is **irredundant**, that is: if there is no inclusion among the S_i 's. The S_i 's are called the **components** of A .*

Proof: Let $\overline{A}$ be the Zariski closure of A in S , and let A' be an irreducible component of $\overline{A}$. Consider $A \cap A'$: either it contains the general point of A' , or it is contained in a countable union B of (Zariski) closed subsets of A' .

In the first case, A' is a component of A . Proceed the $\overline{A}$, after removing its component A' .

In the second case, remove A' from $\overline{A}$, and replace it by B .

Repeat this process, observing that for each component of $\overline{A}$, we let at each step decrease the dimension of the remaining components. So the process ends in finitely many steps for each such component.

The uniqueness statement is easy to check: assume we have two irredundant families S_i and S'_j of components of A . It is sufficient to show that any S_i is contained in some S'_j . If not, $(A \cap S_i)$ is contained in the countable union of the $(A \cap S'_j)$. It is then of first Zariski category in S_i . This is a contradiction to the definition of a component of A . $\square$

The notion of Z -regularity has application to the construction of meromorphic quotients:

Theorem 12.23. *Let $X \in \mathcal{C}$ be normal. Let $A \subset \mathcal{C}(X)$ be Z -regular. Let $T := T(A)$ be the family of components of A , as defined in 12.22. (If T is not covering, we add the family of points of X). Let $q_A : X \dashrightarrow X_{T(A)} = X_A$ be the T -quotient of X .*

Let $t \in A$ such that V_t meets some general fibre F of q_T . Then V_t is contained in F .

(We shall call also q_T the A -reduction of X).

Proof: By the defining property of q_T , its general fibre F is an equivalence class for $R(T(A))$. In particular, any V_t , $t \in A$, is either disjoint from, or contained in F .

Remark 12.24. *Two points in such a fibre are joined by a $T(A)$ -chain. Such a $T(A)$ -chain turns out to be an A -chain if A is stable under specialisation, in the sense defined just below. So, it is important to know if the property P defining A is stable under specialisation (ie: if each component of V_t has the property P when $t \in S_i$, if S_i is a component of A_P , and A_P is Z -regular).*

For example, this is easily seen to be the case for properties 2,3 in 12.20 above. The case of 1. is not presently known, but a positive answer to the following question is expected:

Question 12.25. *Let $f : X \rightarrow Y$ be a special fibration, with $X \in \mathcal{C}$. Is each irreducible component of X_y special, for each $y \in Y$?*

An application of 12.23 above (obtained explicitly in [C92], and in [Ko-Mi-Mo] in the algebraic case) is:

Theorem 12.26. *Let $X \in \mathcal{C}$ be normal. There exists an almost holomorphic map $r_X : X \rightarrow R(X)$ called the **rational quotient** of X such that:*

1. *The fibres of r_X are rationally connected (ie: any two of their points are contained in a connected union of finitely many rational curves)*
2. *If a rational curve C inside X meets some general fibre F of r_X , then C is contained in F .*

Moreover, A -reductions can be constructed in a relative situation as well.

Theorem 12.27. *Let $X \in \mathcal{C}$ be normal. Let $f : X \rightarrow Y$ be a holomorphic fibration. Let $A \subset \mathcal{C}(X)$ be Z -regular. Let $A_f \subset A$ be the set of $t \in A$ such that V_t is contained in some fibre of f . Then $A_f \subset \mathcal{C}(X)$ is also Z -regular.*

If $q_{A_f} : X \dashrightarrow X_{A_f}$ is the A_f -reduction of X , there exists a factorisation $h_{A_f} : X_{A_f} \dashrightarrow Y$ such that $f = h_{A_f} \circ q_{A_f}$.

Moreover, for $y \in Y$ general, the restriction of q_{A_f} to X_y is the A_y -reduction of X_y , if $A_y := A_f \cap \mathcal{C}(X_y)$.

*The map q_{A_f} is called the **A -reduction** of f .*

Proof: The fact that A_f is Z -regular is obvious, because A is Z -regular, and A_f is the intersection of A with its Zariski closed subset $\mathcal{C}(X/Y)$ consisting of t 's such that V_t is contained in some fibre of f . We thus have a quotient q_{A_f} for A_f .

The natural map $f_* : \mathcal{C}(X/Y) \rightarrow Y$ now defines the factorisation h_{A_f} , because the equivalence classes for $T(A_f)$ are obviously contained in fibres of f . If one prefers: $R(f)$ contains obviously $R(T(A_f))$.

We shall now show that, for y general in Y , $(T(A_f))_y = T(A_y)$. This will immediately imply that $(X_y)_{A_y} = (X_y)_{(A_f)_y}$, and so the claim.

Notice first that, by definition, $A_y = (A_f)_y$, for any y . Next, if $T'(A_f)$ is the union of all components of $T(A_f)$ surjectively mapped to Y by f_* , then $(T'(A_f))_y = (T(A_f))_y$ for y general in Y . It is thus sufficient to show that $(T'(A_f))_y = T(A_y)$ for y general in Y .

This we now show: first, we have, for y general: $A_y = (A_f)_y \subset (T'(A_f))_y$. This gives the first inclusion: $T(A_y) \subset (T'(A_f))_y$. To show the reverse, let T be a component of $T'(A_f)$. Then: $T_y \subset T(A_y)$ for general y . Indeed: $T/(A_f \cap T)$ is contained in a countable union of Zariski closed proper subsets B_j of T . For each j , the set C_j of $y \in Y$ such that $B_j \cap T_y$ has empty interior in T_y is Zariski closed proper in Y .

If C is the union of the C_j 's, and y is not in C , we conclude that $(T'(A_f))_y \subset T(A_y)$, as claimed $\square$

The main applications of 12.27 are to construct relative versions of the rational quotient, of the core, and other natural fibrations. For example: applying 12.27 to the family of rational curves on $X \in \mathcal{C}$, we get:

Proposition 12.28. *Let $f : X \rightarrow Y$ be a fibration, $X \in \mathcal{C}$. Then f has a **rational quotient**, that is: a factorisation $f = R_f \circ r_f$, where $r_f : X \rightarrow R(f)$ is a fibration which induces the rational quotient $r_{X_y} : X_y \rightarrow R(X_y) = (R(f))_y$ for y general in Y .*

12.5. Stability.

Definition 12.29. Let $A \subset \mathcal{C}(X)$ be Z -regular, for $X \in \mathcal{C}$. We say that A is **stable** if it has the following two properties:

- (stab1) The general fibre of $q_T : X \rightarrow X_T$ is in A if T is any **covering** irreducible component of A (in the sense of Definition 12.22 above).
- (stab2) $t \in A$ if $V_t = V \subset X$ is a subvariety and $f : V \dashrightarrow W$ is a fibration with general fibre V_w in A , such that a subvariety $W' \subset V$ exists, with $[W'] \in A$, and $f(W') = W$. (By abuse of language, we say that a subvariety Z of X is in A if the point $[Z] \in \mathcal{C}(X)$ which parametrises this subvariety is in A .)

Let us give some examples:

Theorem 12.30. Let $A_P(X) \subset \mathcal{C}(X)$ be Z -regular, for $X \in \mathcal{C}$ be as in theorem 12.20 above. Then A_P is stable in the following two cases:

1. P is the property of being special, if X is smooth.
2. A_P is the set of t 's such that $\pi_1(\hat{V}_t)_X \in \mathcal{G}$, where $\mathcal{G}$ is a class of (isomorphism classes) of groups which is stable in the sense of [C98]. (Such classes are the class of finite, or solvable groups, for example).

The proof is given in section 3 for [1.], and in [C98] for [2.]. The basic result about stability is the following:

Theorem 12.31. Let $A \subset \mathcal{C}(X)$ be Z -regular and stable, for $X \in \mathcal{C}$ normal. Let $q_A : X \dashrightarrow X_A$ be the $T(A)$ -quotient of X . Then:

1. The general fibre of q_A is in A .
2. The general fibre F of q_A contains V , If V is a subvariety of X which is in A and meets F .

Proof: The second property is just one of the defining properties of q_A . To show the first one, we observe that, because of [stab1.], it is obviously satisfied for the quotient of X by any of the covering irreducible components of $T(A)$, and so for the first step of the construction of q_A , as exposed in section 12.3.

We now go to the induction step (“the general fibre of q_{n+1} is in A if q_n has this same property”). We use here this same section 12.3 to which we refer, with the same notations (and $T := T(A)$). Then the subvarieties V'_t constructed there from the varieties V_t are also in A , by [stab2.] applied to q' and $V_t \in A$, by the hypothesis $T_{n+1} \subset T(A)$, using the induction hypothesis (“the general fibre of q' is in A ”).

And so, by [stab1.] again, we see that the general fibre of q_{n+1} is indeed in A $\square$

RÉFÉRENCES

- [Ab97] D.Abramovich. Langs map and Harris conjecture. *Isr.J.Math.* 101 (1997), 85-91.
- [A-N99] D.Arapura-M.Nori.Solvable Fundamental Groups of Algebraic Varieties and Kähler Manifolds. *Comp.Math.* 116(1999), 173-193.
- [Ba75] D.Barlet. Espace analytique réduit des cycles analytiques complexes compacts d'un espace analytique de dimension finie. *LNM* 482 (1975), 1-158.

- [B-P-V84] W.Barth-C.Peters-A.Van de Ven. Compact Complex Surfaces. *Erg. der Math. u. Ihrer Grenzgebiete. 3.Folge, Band 4. Springer Verlag. (1984)*
- [Be89] A.Beuville. Annulation du H^1 pour les fibrés en droites plats. Proceedings Bayreuth 1989. *LNM 1507(1992)*, 1-15.
- [Blo26] S.Bloch. Sur les systemes de fonctions uniformes satisfaisant l'équation d'une variété algébrique dont l'irrégularité dépasse la dimension. *J.Math.pures et Appliquées. 5 (1926)*, 19-66.
- [Bo79] F.A.Bogomolov. Holomorphic Tensors and vector Bundles on Projective Varieties. *Math.USSR.Izv.13 (1979)*, 499-555.
- [Bo-T00] F.A.Bogomolov-Y.Tschinkel. Density of rational points on elliptic K3 surfaces. *Asian Math.J.4 (2000)*, 351-368.
- [Br00] M.Brunella. Courbes entières dans les surfaces algébriques complexes. *Séminaire Bourbaki. Novembre 2000. Exposé 881*.
- [B-L00] G.Buzzard-S.Lu. Algebraic surfaces holomorphically dominable by $\mathbb{C}^2$. *Inv.Math.139. (2000)*, 617-59.
- [C80] F.Campana. Réduction algébrique d'un morphisme faiblement Kählérien propre. et applications. *Math. Ann. 256 (1980)*, 157-189.
- [C81] F.Campana. Coréduction algébrique d'un espace analytique faiblement Kählérien compact. *Inv.Math. (1981)*
- [C85] F.Campana. Réduction d'Albanese d'un morphisme faiblement Kählérien propre et applications I,II. *Comp.Math.54(1985)*, 373-416.
- [C92] F.Campana. An application of twistor theory to the non-hyperbolicity of certain compact symplectic Kähler manifolds. *J.Reine.Angew.Math.425 (1992)*, 1-7.
- [C92'] F.Campana. Connexité rationnelle des variétés de Fano. *Ann.Sc.ENS.25(1992)*, 539-45.
- [C94] F.Campana. Remarques sur le revêtement universel des variétés Kählériennes compactes. *Bull.S.M.F (1994)*
- [C95] F.Campana. Fundamental Group and Positivity Properties of Cotangent Bundles of Compact Kähler Manifolds. *J.Alg.Geom.4 (1995)*, 487-505.
- [C98] F.Campana. Negativity of compact curves in infinite étale covers of projective surfaces. *J.Alg.Geom.7(1998)*, 673-693.
- [C01] F.Campana. Ensembles de Green-Lazarsfeld et quotients résolubles des groupes de Kähler. *J.Alg.Geom. 10 (2001)*, 599-622.
See also: $\mathcal{G}$ -connectedness of compact Kähler manifolds I. *Contemp.Math. 241 (1999)*, 85-97 for a survey and extension.
- [C-P00] F.Campana-T.Peternell. Complex Threefolds with Non-trivial holomorphic 2-Forms. *J.Alg.Geom.9 (2000)*, 223-264.
- [C-Z99] F.Campana-Q.Zhang. Kähler threefolds covered by $\mathbb{C}^3$. Preprint (1999)
- [CT86] J.L.Colliot-Thélène. Arithmétique des variétés rationnelles et problèmes birationnels. *Proc. ICM. Berkeley (1986)*, 641-653.
- [CT-S-S97] J.L.Colliot-Thélène-A.Skorobogatov-P.Swinnerton-Dyer. Double fibers and double covers: paucity in rational points. *Acta Arithm. 79 (1997)*, 113-135.
- [Cox82] D.Cox. Mordell-Weil groups of elliptic curves over $C(t)$ with $p_g = 0$, or 1. *Duke Math.J. 49(1982)*, 677-689.
- [D-G95] H.Darmon-A.Granville. On the equations $z^m = F(x, y)$ and $A.x^p + B.y^q = C.z^r$. *Bull.London Math.Soc. 27 (1995)*, 513-543.
- [Db01] O.Debarre. Higher-Dimensional Algebraic Geometry. *Universitext. Springer Verlag (2001)*.
- [D-E00] J.P.Demailly-J.El Goul. Hyperbolicity of generic hypersurfaces in the projective 3-space. *Amer.J.Math. 122(2000)*, 515-546.
- [D-L-S94] J.P.Demailly-L.Lempert-B.Shiffman. Algebraic approximation of holomorphic maps from Stein domains to projective manifolds. *Duke Math.J.76 (1994)*, 333-63.
- [D-P-S93] J.P.Demailly-T.Peternell-M.Schneider. Kähler manifolds with numerically effective Ricci class. *Comp.Math.89 (1993)*, 217-40.
- [D-P-S96] J.P.Demailly-T.Peternell-M.Schneider. Compact Kähler Manifolds with hermitian semipositive anti-canonical bundle. *Comp.Math. 101.(1996)*, 217-224.
- [E80] H.Esnault. Classification des variétés de dimension 3 et plus. *Séminaire Bourbaki. Exposé 568. (1980/81)*.

- [F94] G.Faltings. The general case of S.Lang's Conjecture. The Barsotti Symposium. W.Messing and V.Cristante Eds. *Academic Press. Cambridge Mass. (1994)*, 175-182.
- [F-MK92] L.Fong-J.McKernan. Log Abundance For Surfaces,in: Flips And Abundance for Algebraic Threefolds, J.Kollar Ed. *Astérisque 211.(1992). Soc.Math. de France.pp. 127-137.*
- [F-M94] R.Friedman-J.Morgan. Smooth Four-Manifolds and Complex Surfaces. *Springer Verlag. Erg.der Mathem.27.(1994).*
- [Fu82] A.Fujiki. On the Douady Space of a complex space in $\mathcal{C}$. *Publ. RIMS (1982).*
- [Fuj78] T.Fujita. On Kähler fibre spaces over curves. *J.Math.Soc.jap. 30 (1978), 779-794.*
- [G-H-S01] T.Grabner-M.Harris-J.Starr. Families of rationally connected varieties. Preprint 2001.
- [Gr89] H.Grauert. Jetmetriken und hyperbolische geometrie.*Math.Z.200 (1989),149-68.*
- [Gr60] H.Grauert.Ein Theorem der analytischen Garbentheorie und die Modulräume komplexer Strukturen. *Publ.Math.IHES 5 (1960),1-64 and vol.16 (1963),35-36.*
- [Gr65] H.Grauert. Mordell's Vermutung über rationale Punkte auf algebraischen Kurven und Funktionenkörpern. *Publ. Math. IHES. 25 (1965), 131-149.*
- [H-S00] M.Hindry-J.Silverman. Diophantine Geometry: an Introduction. *GTM 201. Springer-Verlag. (2000).*
- [H-T00] J.Harris-Y.Tschinkel. Rational points on quartics. *Duke Math.J. 104 (2000), 477-500.*
- [I72] S.Iitaka.Genera and Classification of Algebraic Varieties.*Sugaku 24 (1972),14-27.*
- [I73] S.Iitaka.On Algebraic Varieties whose Universal covering Manifolds are Complex Affine $\mathbb{C}^3$ -space.Number Theory... *Volume in honour of Y.Akizuki.(1973),95-146.Tokyo Kunikuniya.*
- [K86] V.A.Kaimanovitch. Brownian Motion and Harmonic Functions on Covering Manifolds. An Entropy Approach. *Sov.Math.Dokl. 33. (1986), 812-816.*
- [K81] Y.Kawamata.Characterisation of Abelian varieties.*Comp.Math. (1981),253-76.*
- [K99] Y.Kawamata. Pluricanonical Forms.*Contemp.Math.241(1999),193-209.*
- [K80] Y.Kawamata. On Bloch's Conjecture.*Inv.Math.57 (1980), 97-100.*
- [K-V80] Y.Kawamata-E.Viehweg. On a characterisation of an Abelian Variety. *Comp.Math. 41 (1980), 355-359.*
- [K-M-M87] Y.Kawamata,K.Matsuda,K.Matsuki.Introduction to the Minimal Model Problem.*Adv.Studies in Pure Mathematics 10 (1987),283-360.*
- [K-MK97] S.Keel-S.McKiernan. Rational curves on quasi-projective surfaces. *Memoirs AMS 669 (1999).*
- [Kob98] S.Kobayashi. Hyperbolic complex spaces.*Grundlehren der Math.Wiss. 318.Springer Verlag.(1998).*
- [Kob76] S.Kobayashi. Intrinsic distances, measures, and geometric function theory.*Bull.AMS 82 (1976), 357-416.*
- [K-O75] S.Kobayashi-T.Ochiai.Meromorphic mappings into compact complex spaces of general type.*Inv.Math.31 (1975),7-16.*
- [Ko86] J.Kollar. Higher direct image sheaves of dualising sheaves.*Ann. Math. 123 (1986), 11-42.*
- [Ko96] J.Kollar.Rational curves on Algebraic varieties.*Erg.der Math.und ihrer Grenz.32.Springer Verlag.(1996)*
- [Ko93] J.Kollar.Shafarevitch maps and plurigenera of Algebraic Varieties.*Inv.Math.113 (1993), 177-215.*
- [Ko-Mi-Mo92] J.Kollar-Y.Myiaoka-S.Mori. Rationally connected Varieties. *J.Alg.Geom.1 (1992),429-48.*
- [Ko-Mo98] J.Kollar-S.Mori. Birational Geometry of Algebraic Varieties. *Cambridge Tracts in Mathematics 134.(1998).*
- [La86] S.Lang.Hyperbolic and Diophantine Analysis.*Bull.AMS.14(1986),159-205.*
- [La91] S.Lang. Survey of Diophantine Geometry.(Number Theory III).EMS vol.60. *Springer Verlag 1991.*
- [L-N59] S.Lang-A.Néron. Rational points of Abelian Varieties over Function Fields. *Amer.J.Math. 81 (1959), 95-118.*
- [Lieb75] D.Liebermann.Compactness of the Chow Scheme :Applications to automorphisms and deformations of Kähler Manifolds. *Lecture Notes in Mathematics 670. F.Norguet Ed.(1975),140-186.*
- [L-S75] D.Liebermann-E.Sernesi. Semi-continuity of Kodaira Dimension. *Bull.Amer.Math. Soc. 81 (1975), 459-460.*
- [L01] S.Lu.Multiply marked Riemann surfaces and the Kobayashi pseudometric on Algebraic manifolds.Preprint (2001)
- [Mae83] K.Maehara. A finiteness Property of Varieties of General Type. *Math.Ann. 262 (1983), 101-123.*
- [Man63] Y.Manin. Rational Points of Algebraic Curves over Function Fields. *Izv.Akad.Nauk. SSSR 27 (1963), 737-756.*

- [Miy88] Y.Miyaoka. On the Kodaira Dimension of Minimal Threefolds. *Math. Ann.* 281 (1988), 325-332.
- [Mois70] B.Moishezon. Algebraic Varieties and Compact Complex Spaces. *Actes du Congrès International des Mathématiciens, Nice 1970. Vol.2*, 643-648.
- [Mok92] N.Mok. Factorisation of Semi-Simple Discrete Representations of Kähler Groups. *Inv.Math.* 110 (1992), 557-614.
- [Mo88] S.Mori. Flip Theorem and the existence of Minimal Models for Threefolds. *J.AMS.* 1(1988), 117-253.
- [Mw95] A.Moriwaki. Geometric Height inequality on Varieties with ample cotangent bundle. *J.Alg.Geom.* 4 (1995), 385-96.
- [Na87] M.Namba. Branched Coverings and Algebraic Functions. *Pitman Research Notes* 161. Longman Sc. and Tech. (1987).
- [Nak98] N.Nakayama. Invariance of Plurigeners of algebraic Varieties. RIMS Preprint (1998).
- [Oc77] T.Ochiai. On holomorphic curves in algebraic varieties with ample irregularity. *Inv.Math.* 43 (1977), 83-96.
- [Pa98] M.Paun. Sur les variétés Kählériennes compactes à classe de Ricci nef. *Bull. Sci. Math.* 122 (1998), 83-92.
- [Par68] A.N.Parshin. Algebraic Curves over Function Fields. *Math.USSR Izv.* 2. (1968), 1145-1170.
- [Pey] E.Peyre. Oral communication. July 2001.
- [SD69] H.Swinnerton-Dyer. Applications of Algebraic Geometry to number Theory. *Proc. Symp. Pure Math.* vol. XX (1969), 1-52.
- [Sim93] C.Simpson. Subspaces of Moduli Spaces of Rank One Local Systems. *Ann.Sc.ENS.* 26 (1993), 361-401.
- [Siu82] Y.T.Siu. Complex Analyticity of Harmonic Maps. *J.Diff.Geom.* 17 (1982), 55-138.
- [Siu02] Y.T.Siu. Extension of Pluricanonical Sections with Plurisubharmonic Weights... Preprint 2001.
- [Sk92] V.Shokurov. 3-fold log-flips. *Izv.Akad.Nauk.Ser.Mat.* 56 (1992), 105-203.
- [S-V86] A.Sommese-A.Van de Ven. Homotopy of Pullback of Varieties. *Nagoya Math. J.* 102 (1986), 79-90.
- [U75] K.Ueno. Classification Theory of Algebraic Varieties and Compact Complex Manifolds. *LNM* 439(1975), Springer Verlag.
- [Vi82] E.Viehweg. Die Additivität der Kodaira Dimension für Projektive Faserräume über Varietäten des Allgemeinen Typs. *J.Reine.Angew.Math.* 330(1982), 132-142.
- [Vi83] E.Viehweg. Weak positivity and the additivity of the Kodaira dimension for certain fibre spaces. *Adv.Stud.Pure Math. Vol I. North-Holland.* (1983), 329-353.
- [Vi86] E.Viehweg. Vanishing theorems and positivity of Algebraic fibre spaces. *J.Proc. Int. Congr. Math. Berkeley* (1986).
- [Zh96] Q.Zhang. On projective manifolds with nef anticanonical bundles. *J.Reine.Angew.Math.* 478 (1996), 57-60.
- [Z97] K.Zuo. Factorisation theorems for representations of fundamental groups of algebraic varieties. Preprint Kaiserslautern. (1997). See also: Representation of fundamental groups of algebraic varieties. Lecture Notes in Mathematics 1708 (1999). Springer-Verlag.

F.Campana

Département de Mathématiques.

Université Nancy 1

BP 239.

F-54506.Vandoeuvre-les-Nancy.Cedex.